

Antifungal Effects of *Lactobacillus* spp. Bacteria on *Candida* Yeast

Ergin KARİPTAŞ *  Şener TULUMOĞLU ** Belgin ERDEM *

* Department of Biology, Faculty of Sciences and Arts, Ahi Evran University, 40200 Kırşehir - TURKEY

** Doctor Behçet Uz Hospital for Child Diseases Education and Research, 35210 Konak, İzmir - TURKEY

Makale Kodu (Article Code): KVFD-2010-2363

Summary

Thirty *Lactobacillus* were isolated from the stool of 5-15 year-old children using MRS agar and identified by API 50CHL. Also, 50 *Candida* were isolated from blood cultures using SD agar and identified by Vitek 2 instrument with YST kit. Antifungal effects of *Lactobacillus* bacteria on *Candida* were explored using well diffusion method in SD agar. *Lactobacillus* had the most prominent antifungal effect against on *C. albicans* (M29, M36), *C. parapsilosis* (M25, M26), *C. famata* (M28) and *C. guilliermondii* (M38) yeasts. This study shows the presence of antifungal effect of *Lactobacillus* on *Candida* isolated from blood cultures.

Keywords: Antifungal effect, *Lactobacillus*, *Candida*, Blood culture

Lactobacillus Bakterilerinin *Candida*'lar Üzerine Antifungal Etkileri

Özet

Otuz adet *Lactobacillus*, 5-15 yaş grubu çocuk gaitasından MRS agar'da izole edilerek API 50CHL cihazı tanımlanmaları yapılmıştır. Ayrıca 50 adet kan kültürü kaynaklı *Candida* ise SD agar'da izole edilerek Vitek 2 cihazında YST kiti ile tanımlanmaları yapılmıştır. *Lactobacillus*'ların *Candida*'lar üzerindeki antifungal etkileri SD agar'da kuyu difüzyon yöntemi ile tespit edilmiştir. *Lactobacillus*'lar en fazla *C. albicans* (M29 ve M36), *C. parapsilosis* (M25, M26), *C. famata* (M28) ve *C. guilliermondii* (M38) mayaları üzerinde antifungal etki göstermiştir. Bu çalışma *Lactobacillus*'ların, kan kültürlerinden elde edilen *Candida* izolatları üzerine antifungal etkisinin varlığını göstermektedir.

Anahtar sözcükler: Antifungal etki, *Lactobacillus*, *Candida*, Kan kültürü

INTRODUCTION

Yeasts are microorganisms commonly found in nature. They are present in the normal flora (in moist places like the intestinal system, mouth etc.) in a human body. Hospital-acquired fungal infections cause serious morbidity and mortality and it is clear that fungal diseases have emerged as important public health problems recently ¹. *Candida* constitutes 80% of hospital-acquired fungal infections. In recent years, there has been a significant increase in non-albicans *Candida* species, especially in blood cultures. *C. albicans* and non-albicans species pose a high risk of infection in the blood circulation system of children ². It has been reported that lactic acid bacteria (LAB) can produce

antimicrobial substances with the capacity to inhibit the growth of pathogenic and spoilage microorganisms. Organic acids, hydrogen peroxide, diacetyl and bacteriocins are included among these compounds ³. Recent years have witnessed an increase in the studies exploring the antifungal effects of *Lactobacillus* strains. These studies have primarily focused on protection against urogenital candidiasis and prevention of fungal contamination of food ^{4,5}. Atanasova et al. ⁶ established that *Lactobacillus paracasei* subsp *paracasei* M3 bacterium had an anti-fungal effect on *C. albicans*, *C. pseudointermedia* and *C. blankii*. Antifungal resistance of *Candida* species is an increasing problem ⁷. Despite the wellfounded



İletişim (Correspondence)



+90 386 2114542



ekariptas@ahievran.edu.tr

enthusiasm for new antifungal agents, it is very important to continue to assess the activity of newer agents ⁷.

Studies about the antifungal effect of *Lactobacillus* on *Candida* isolated from blood cultures are scarce. To our knowledge, this is the first report about the antifungal effect of *Lactobacillus* on *Candida* isolated from blood cultures. The aim of the present study is to examine the antifungal effect of *Lactobacillus* of human intestinal origin on *Candida* yeasts isolated from blood cultures. Besides, we have studied the pattern of *Candida* spp. to common usable antifungals in Turkey.

MATERIAL and METHODS

Microorganisms

The study used 30 *Lactobacillus* bacteria isolated from the stool of children in the 5-15 age group at the Microbiology Laboratory of Dr. Behçet Uz Children's Hospital. *Lactobacillus* was kept in bead tubes (Cryobank, Mast Diagnostics, France) at -20°C.

Bottles of blood cultures collected from the units were placed in BD Bactec 9240 (BD, USA) instrument. The bottles were incubated in the instrument for 7 days at 37°C and the bottles which were reproduction positive were transferred to SD agar (BD, USA) culture medium. Cultures that showed growth were microscopically examined (direct examination and gram staining) and those containing yeasts were separated. Each *Candida*

strain was isolated from individual samples. Yeasts were sorted out using Vitek 2 (BioMérieux, France) kit. Isolates were stored in bead tubes (Cryobank, Mast Diagnostics, France) at -20°C ⁸.

Antifungal Assay

Antifungal effect of *Lactobacillus* on *Candida* was examined using well agar diffusion method ⁹. *Lactobacillus* cultures were incubated in MRS liquid culture medium (BD, USA) under anaerobic conditions at 37°C for 24 h. The cultures were then centrifuged at 10.000 cfu for 10 min. The floating supernatant was recovered with an injector and filtered through 0.45 µ pore filter (Millipore, Molsheim, France) ⁹. *Candida* were activated in SD agar and planted into 0.5 McF SDA (BD) culture media. In all media, 10 mm wells were opened and added 100 µl supernatant. The plaques were kept at room temperature for 2 h and left to incubate at 37°C for 24 h. Then, the zones around the wells were measured in mm ⁹.

RESULTS

Table 1 presents species of *Candida* yeasts isolated from blood cultures and their percentages. Of the 50 species isolated, 16 (32%) were *C. albicans* and 34 (68%) were non-albicans species. Yeasts isolated from blood cultures were 16 *C. albicans* (32%), 22 *C. parapsilosis* (44%), 4 *C. famata* (8%), 3 *C. guilliermondii* (6%), and 1 *C. lusitanae* (2%). Our study shows that pathogenicity

Table 1. *Candida* strains isolated from blood cultures and their distributions

Tablo 1. Kan kültürlerinden izole edilen *Candida* suşları ve onların dağılımları

Services	<i>C. albicans</i>	<i>C. parapsilosis</i>	<i>C. famata</i>	<i>C. pelliculosa</i>	<i>C. guilliermondii</i>	<i>C. lusitanae</i>
Emergency - ICU *	2	15	-	-	-	1
Infectious diseases	-	1	-	-	-	-
New-born unit	4	-	1	-	-	-
Oncology - Hematology	2	1	1	-	-	-
Burn	3	-	-	-	-	-
Surgical - ICU	3	2	2	2	2	-
Premature - ICU	1	2	-	2	-	-
Nursing infant	-	-	-	-	1	-
Cardiology	-	1	-	-	-	-
Big boy	1	-	-	-	-	-
Total (%)	16 (%32)	22 (%44)	4 (%8)	4 (%8)	3 (%6)	1 (%2)

* ICU : Intensive Care Unit

of non-albicans species is much higher than that of *C. albicans*. Nearly similar findings have been reported by Levy et al.². Various studies carried out in Turkey have revealed that non-albicans isolated from blood cultures are as common as or more common than *C. albicans*¹⁰⁻¹².

Table 2 shows the antifungal effects of *Lactobacillus* on *Candida* yeasts. *Lactobacillus* was seen to have antifungal effects against *C. albicans* (M29, M36), *C. parapsilosis* (M25, M26 and M44), *C. famata* (M28) and *C. guilliermondii* (M38) strains. However, the effect of *Lactobacillus* against *Candida* was observed as weak.

Table 2. Antifungal effects of *Lactobacillus* on *Candida* yeasts

Table 2. *Lactobacillus*'ların *Candida* mayaları üzerine antifungal etkileri

<i>Lactobacillus</i> spp.	Diameter of the Zones of Inhibition in mm								
	<i>C. albicans</i> M29	<i>C. albicans</i> M36	<i>C. parapsilosis</i> M25	<i>C. parapsilosis</i> M26	<i>C. parapsilosis</i> M44	<i>C. famata</i> M28	<i>C. guilliermondii</i> M38	<i>C. guilliermondii</i> M40	<i>C. guilliermondii</i> M41
L1	2	4	2		2	2			
L2	2	3.5	2		2	2			
L3	2	3		2	2	2			
L4		4			2		2		
L5		3	4						
L6							1.5		
L7	4		4		2	2	1.5		
L8		2	4		4				
L9		2				2			
L10						2.5			
L11		2				2			
L12					1.3				
L13			4		3.5	2	1.5		
L14									
L15			4						
L16		4							
L17	4	4	4	2			4	2	4
L18	4	3.5	4	2			3.5	2.5	
L19		1.5	5			2.5	4		
L20	2	1	5	2		2		2	
L21	2	6	4	2			3.4		
L22		3.5	5				3		
L23	4	4	4		3.5		2.5	2	8
L24	4	4	4				3.5		
L25	4	3.5	4				3		
L26		2.5	5	2	3.5		3		
L27	3.5	4	4				3		
L28	4	4	4				3		
L20	2	3	4						
L30	3.5	2	4		1.5		2		2

DISCUSSION

Various studies point out that some molecules with protein structure produced by bacteria have antifungal characteristics ⁶. Özkaya et al.¹³ studied the antifungal activity of *Lactobacillus* on various yeasts. This study, which used 19 *Lactobacillus* strains and 12 yeast strains, demonstrated that *Lactobacillus* bacteria had varying degrees of inhibitive effects against *C. krusei* 4B, *C. lipolytica* 5A, *C. lusitaniae* 7, *C. ciferrii* 10, *S. cerevisiae* 12, 13, *C. pseudotropicalis* 16 and *C. glabrata* 18 yeasts. An examination of *L. plantarum* and *C. albicans* mixed cultures under anaerobic conditions at 37°C for 48 h showed growth in *L. plantarum*, but not in *C. albicans* ¹⁴.

Studies about the strains isolated from the human intestinal system are scarce. It was found that *L. rhamnosus* and *L. paracasei* probiotic bacteria isolated from human stool had a high degree of antifungal effect against *C. albicans* ATCC 10291 ¹⁵. In our study, however, *Lactobacillus* had a poor antifungal effect.

The literature does not contain studies about the antifungal effects of *Lactobacillus* in human intestinal system origin on yeasts (*Candida*) isolated from the human blood circulation system. Our study could be considered the first of its kind in this respect. Presence of yeasts in the blood circulation system in children poses a great risk, which needs to be eliminated. We know that over time resistance develops against antifungal drugs used against yeasts. *Lactobacillus* spp. of human origin, which has probiotic characteristics, may be used to reduce fungus through their antifungal effects or to eliminate yeasts, when combined with antifungal drugs. In conclusion, it is believed that outputs of this study will help to identify to find out new antifungal agents against resistance problem. Therefore, further and more detailed studies are needed in this field.

REFERENCES

1. Chen KW, Chen YC, Lo HJ, Odds FC, Wang TH, Lin CY, Li SY: Multilocus sequence typing for analyses of clonality of *Candida albicans* strains in Taiwan. *J Clin Microbiol*, 44, 2171-2178, 2006.
2. Levy I, Rubin LG, Vasistha S, Tucci V, Sood SK: Emergence of *Candida parapsilosis* as the predominant species causing Candidemia in children. *Clin Infect Dis*, 26, 1086-1088, 1998.
3. Sezer Ç, Güven A: Investigation of bacteriocin production capability of lactic acid bacteria isolated from foods. *Kafkas Univ Vet Fak Derg*, 15, 45-50, 2009.
4. Corsetti, A, Gobbetti M, Rossi J, Damiani P: Antimould activity of sourdough lactic acid bacteria: Identification of a mixture of organic acids produced by *Lactobacillus sanfrancisco* CB1. *Appl Microbiol Biotechnol*, 50, 253-256, 1998.
5. Barousse MM, van der Pol BJ, Fortenberry D, Orr D, Fidel Jr PL: Vaginal yeast colonisation, prevalence of vaginitis, and associated local immunity in adolescents. *Sex Transm Infect*, 80, 48-53, 2004.
6. Atanassova M, Choiset Y, Dalgalarondo M, Chobert JM, Dousset X, Ivanova I, Haertle T: Isolation and partial biochemical characterization of proteinaceous anti-bacteria and anti-yeast compound produced by *Lactobacillus paracasei* subsp. *paracasei* strain M3. *I J Food Microbiol*, 87, 63-73, 2003.
7. Berry V, Badyal DK: Sensitivity of clinical isolates of *Candida* species to antifungal drugs. *J Med Education Res*, 8, 214-216, 2006.
8. Warren NG, Hazen KC: Cryptococcus and other yeasts of medical importance. In *Manual of Clinical Microbiology*, ed. Murray, P.R., Baron, E.J.O., Tenover, F.C. and Tenover, R.H. pp. 1184-1199. Washington: ASM Pres, 1999.
9. Toba T, Samant SK, Itoh T: Assay system for detecting bacteriocin in microdilution wells. *Lett Appl Microbiol*, 13, 102-104, 1991.
10. Ermertcan Ş, İnci R, Hilmioğlu S, Tümbay E: Kan kültürlerinden soyutlanan *Candida* kökenlerinin flukonazole in vitro duyarlılığı. *İnfek D*, 12, 531-533, 1998.
11. Koç AN, Erdem F, Çetin N: Kan kültürlerinde üreyen mayaların retrospektif olarak değerlendirilmesi ve antifungal duyarlılıkları. *T Mikrobiyol Cem Der*, 29, 177-182, 1999.
12. Tulunoğlu Ş, Kariptaş E, Erdem B: Identification and antifungal susceptibility of *Candida* isolates from various clinical specimens in Doctor Behçet Uz Hospital. *Anatol J Clin Investig* 3, 170-173, 2009.
13. Özkaya FD, Karabıçak N, Kayalı R, Esen B: Inhibition of yeasts isolated from traditional Turkish cheeses by *Lactobacillus* spp. *Int J Dairy Technol*, 58, 111-114, 2005.
14. Wynne AG, McCartney AL, Brostoff J, Hudspith BN, Gibson GR: An in vitro assessment of the effects of broad-spectrum antibiotics on the human gut microflora and concomitant isolation of a *Lactobacillus plantarum* with anti-*Candida* activities. *Anaerobe*, 10, 165-169, 2004.
15. Verdenelli MC, Ghelfi F, Silvi S, Orpianesi C, Cecchini C, Cresci A: Probiotic properties of *Lactobacillus rhamnosus* and *Lactobacillus paracasei* isolated from human faeces. *Eur J Nutr*, 48, 355-363, 2009.