

---

# MYCOTAXON

ISSN (print) 0093-4666 (online) 2154-8889 © 2016. Mycotaxon, Ltd.

April–June 2016—Volume 131, pp. 331–335

<http://dx.doi.org/10.5248/131.331>

---

## ***Acrodictys*, *Corynespora*, *Karstenula*, *Oncopodium*, and *Sporocadus*: new genera for Turkey**

ELŞAD HÜSEYİN\*, FARUK SELÇUK & KADRIYE EKICI

*Department of Biology, Arts and Sciences Faculty, Ahi Evran University, Kırşehir, Turkey*

\* CORRESPONDENCE TO: [elsadhuseyin@hotmail.com](mailto:elsadhuseyin@hotmail.com)

**ABSTRACT**—Five microfungial genera are recorded for the first time in Turkey. Brief morphological descriptions based on collections from Kırşehir province are provided for *Sporocadus pezizoides* (Phlogicylindriaceae), *Corynespora cambrensis* (Corynesporascaceae), *Karstenula rhodostoma* (Montagnulaceae), *Acrodictys elliptica* (Pezizomycotina), and *Oncopodium elaeagni* (Pezizomycotina).

**KEY WORDS**—anamorphic fungi, Anatolian Peninsula, Ascomycota, biodiversity

### **Introduction**

Most studies on micromycetes in Turkey have focused on powdery mildews (*Erysiphales*) and rusts (*Pucciniales*). Other microfungi, especially the asexual states of saprobic ascomycetes that develop on forest trees and bushes, are insufficiently studied. However, in the last fifteen years the research on these fungi has increased (Akgul et al. 2015; Hüseyin 2004; Hüseyinov & Selçuk 1999, 2001; Hüseyinov 2000; Hüseyin & Selçuk 2001, 2002a,b, 2014; Selçuk et al. 2003, 2009, 2012a,b, 2014; Mel'nik et al. 2004; Selçuk & Ekici 2014; Selçuk & Hüseyin 2014; Hüseyin et al. 2005, 2014; Vasighzadeh et al. 2014).

Kırşehir province (38°50'–39°50'N 33°30'–34°50'E) is located in the middle of Anatolian Peninsula. This area lies within the Irano-Turanian phytogeographic region. According to the data obtained from the Kırşehir meteorological station, annual precipitation is 340–400 mm and the annual average temperature is 11°C.

This study contributes to the list of mycobiota of Turkey by adding five new genera.

## Materials & methods

The specimens were collected during periodical mycological excursions in Boztepe district, Kırşehir Province, between 2012 and 2014. Microscopic studies were carried out on slides, with the samples prepared in distilled water. Morphological features were examined using an Olympus BX 53 research microscope (Japan) and Axio Imager 2 microscope equipped with Nomarski differential interference contrast optics. Hyphomycetous genera were identified via the key by Seifert et al. (2011). Species were identified by consulting handbooks and other publications (Ellis 1960, Sutton 1980, Dennis 1981, Morelet 1985, Ellis & Ellis 1987, Manoharachary et al. 2006, Magyar & Révay 2009). The host plants were identified using “Flora of Turkey and East Aegean Islands” (Davis 1965–85). Taxonomic names and citations follow Kirk et al. (2008) and Kirk (2015). All specimens are deposited in the Mycological Collection, Department of Biology, Arts and Sciences Faculty, Ahi Evran University, Kırşehir, Turkey (AEUT).

## Taxonomy

*Sporocadus pezizoides* (Ellis & Everh.) M. Morelet, Ann. Soc. Sci. Nat. Archéol.

Toulon Var 37(4): 234 (1985)

Conidiomata dark brown, sporodochial to disciform, 300–400 µm wide, 80–100 µm deep. Conidiophores hyaline, simple, cylindrical, 17.5–25 × 1.5–3 µm. Conidia brown, dark brown, straight or curved, clavate to fusiform with rounded ends, 25–30 × 7–8(–10) µm, 3-septate, septa prominent and very dark brown.

SPECIMENS EXAMINED: TURKEY, KIRŞEHİR PROVINCE, Boztepe district, around Kervansaray mountain, 39°13'10"N 34°13'55"E, 1300 m a.s.l., on dead wood of *Salix alba* L., 20.IV.2013, Kadriye Ekici 203 (AEUT); 30.III.2014, Kadriye Ekici 203a (AEUT).

COMMENTS: The Turkish specimens of *Sporocadus pezizoides* have more slender conidiophores and narrower conidia than those described by Morelet (1985; conidiophores 2.5–3 µm wide, conidia 8.5–10.5 µm wide).

*Corynespora cambrensis* M.B. Ellis, Mycol. Pap. 76: 28 (1960)

Colonies on the natural substratum effuse, hairy, dark blackish brown. Mycelium superficial and immersed. Hyphae branched, 4–5 µm diam. Conidiophores simple, septate, dark brown, (45–)50–55(–65) × 5–7 µm. Conidia dark brown, smooth, (5–)8–12-pseudoseptate, cylindric, (23–)25–57(–75) × (7.5–)8–10 µm.

SPECIMENS EXAMINED: TURKEY, KIRŞEHİR PROVINCE, Boztepe district, around Kervansaray mountain, 39°12'50" N 34°13'01" E, 1440 m a.s.l., on dead wood of *Salix cinerea* L. trunk, 27.IV.2013, Kadriye Ekici 207 (AEUT).

COMMENTS: The Turkish specimen of *Corynespora cambrensis* has wider conidia with more numerous pseudosepta than described for the type specimen (Ellis 1960; 5–10 µm wide, with 2–8 pseudosepta).

*Karstenula rhodostoma* (Alb. & Schwein.) Speg., Dec. Mycol. Ital. 7–12: no. 94 (1879)

Pseudothecia gregarious beneath the bark, blackish or black, globose, globose-depressed, 450–550 µm diam. Hymenium paraphysate. Asci 8-spored, cylindrical, apically rounded, (115–)130–135(–160) × 12.5–18 µm, with thick and short stalk. Ascospores dark brown, guttulate, uniseriate, 3-septate, constricted at the septa, cylindrical, fusiform, rounded at both ends, 22.5–25 × 8–9.5 µm.

SPECIMEN EXAMINED: TURKEY, KIRŞEHİR PROVINCE, Boztepe district, around Kervansaray mountain, 39°13'30" N 34°14'10" E, 1320 m a.s.l., on dead branches of *Rhamnus petiolaris* Boiss. & Balansa, 02.XII.2012, Kadriye Ekici 191 (AEUT).

COMMENTS: The Turkish specimen of *Karstenula rhodostoma* has wider asci and ascospores than those described by Dennis (1981; asci 12 µm wide, ascospores 7–9 µm wide).

*Acrodictys elliptica* Manohar., N.K. Rao, D.K. Agarwal & Kunwar, Indian Phytopath. 59: 91 (2006)

Colonies on the natural substratum effuse, hairy, brownish black, black, 5–10 × 1–2 mm. Mycelium mostly immersed. Hyphae smooth, brown, 3–3.5 µm diam. Conidiophores simple, septate, brown to black, erect, usually straight or slightly flexuous, 30–70 × 4–8 µm. Conidia dark brown, smooth, with 3–4 transverse and 1–2 longitudinal septa, ellipsoidal, ovoid, slightly constricted at the septa, (16–)17.5–22.5(–25) × (8–)9–12.5 µm.

SPECIMEN EXAMINED: TURKEY, KIRŞEHİR PROVINCE, Boztepe district, around Kervansaray mountain, 39°12'05" N 34°13'08" E, 1380 m a.s. l., on dead wood of *Cotoneaster nummularius* Fisch. & C.A. Mey., 04.XI.2012, Kadriye Ekici 62 (AEUT).

COMMENTS: The Turkish specimen of *Acrodictys elliptica* has conidia with more transverse septa and fewer longitudinal septa but otherwise morphologically close to the type description (Manoharachary et al. 2006; 2–3 transverse and 2–3 longitudinal septa).

*Oncopodium elaeagni* D. Magyar, Nova Hedwigia 88: 477 (2009)

Colonies on the natural substratum brown. Mycelium immersed. Hyphae septate, smooth, brown, 4–5 µm diam. Conidiomata sporodochial, black. Conidiophores caespitose, septate, hyaline, 15–40 × 2.5–4.5 µm. Conidia fusiform, limoniform, brown, smooth, with 6–7 transverse and 3–4 longitudinal septa, 25–31 × 12–13(–14) µm, with 2 lateral and 1 central horn-like cells.

SPECIMEN EXAMINED: TURKEY, KIRŞEHİR PROVINCE, Boztepe district, around Kervansaray mountain, 39°13'10"N 34°13'40" E, 1320 m a.s.l., on dead bark of *Elaeagnus angustifolia* L., 27.VII.2013, Kadriye Ekici 134 (AEUT).

COMMENTS: The Turkish specimen of *Oncopodium elaeagni* has conidia that do not differ significantly in size from that cited by Magyar & Révay (2009: 24–30 × 9–18 µm).

#### Acknowledgements

We thank Dr. Eugene Yurchenko (Minsk, Belarus) and Dr. Sevda Kirbag (Elazig, Turkey) for serving as presubmission reviewers. Especially we are grateful to Dr. Shaun Pennycook (Auckland, New Zealand) for detailed linguistic help and nomenclatural review of the manuscript.

#### Literature cited

- Akgul H, Ergul CC, Yılmazkaya D, Akata I, Selçuk F, Huseyin E. 2015. Diversity of microfungi on *Fagaceae* in Uludag forests. *Oxidation Communications* 38(3): 1529–1538.
- Davis PH (ed.). 1965–85. *Flora of Turkey and East Aegean Islands*. Vols 1–9. Edinburgh University Press, Edinburgh.
- Dennis RWG. 1981. *British Ascomycetes*. J. Cramer, Vaduz.
- Ellis MB. 1960. Dematiaceous hyphomycetes. I. *Mycological Papers* 76. 36 p.
- Ellis MB, Ellis JP. 1987. *Microfungi on land plants: an identification handbook*. Croom Helm, London-Sydney.
- Hüseyin E. 2004. *Cumminsella mirabilis* on *Mahonia aquifolium* in Turkey. *Mycotaxon* 90(1): 125–127.
- Hüseyin E, Selçuk F. 2001. New and poorly known genera of microfungi for Turkey. *Turkish Journal of Botany* 25: 437–438.
- Hüseyin E, Selçuk F. 2002a. A new species of *Septoria*. *Pakistan Journal of Botany* 34(2): 113–115.
- Hüseyin E, Selçuk F. 2002b. A new species of *Colletotrichum*. *Israel Journal of Plant Sciences* 50(2): 161–163. <http://dx.doi.org/10.1560/1NUH-YTR6-NBGG-W038>
- Hüseyin E, Selçuk F. 2014. Coelomycetous fungi in several forest ecosystems of Black Sea provinces of Turkey. *Agriculture and Forestry* 60(2): 19–32.
- Hüseyin E, Selçuk F, Şahin A. 2005. The world's second record of *Neoheteroceras flageoletii* reported from Turkey. *Mycotaxon* 94: 241–244.
- Hüseyin E, Selçuk F, Churakov BP. 2014. Microfungi on forest trees and shrubs of Duzce Province (Turkey) and Ulyanovsk Region (Russia). *Ulyanovsk Medical and Biological Journal* 4: 127–139. [In Russian].
- Hüseyinov E. 2000. New records of microfungi for Turkey. *Israel Journal of Plant Sciences* 48(1): 75–78. <http://dx.doi.org/10.1560/9FCD-2N2D-M0C7-8FE5>
- Hüseyinov E, Selçuk F. 1999. New records of phytopathogenic microfungi for Turkey. *Plant Disease Research* 14: 175–176.
- Hüseyinov E, Selçuk F. 2001. Contribution to study of mycoflora of Turkey. I. Coelomycetes of orders *Melanconiales* and *Sphaeropsidales* on forest trees and shrubs in the Black Sea coast (Rize and Trabzon provinces). *Mikologiya i Fitopatologiya* 35(1): 28–33.
- Kirk P. 2015. Index Fungorum. <http://www.indexfungorum.org>. Accessed 10 August 2015.
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. *Ainsworth & Bisby's dictionary of the fungi*, 10th ed.. CABI International, Wallingford, UK. 771 p.
- Magyar D, Révay Á. 2009. *Oncopodium elaeagni*, a new hyphomycete from Hungary. *Nova Hedwigia* 88(3-4): 475–481. <http://dx.doi.org/10.1127/0029-5035/2009/0088-0475>

- Manoharachary C, Rao NK, Agarwal DK, Kunwar IK. 2006. Two new species of *Acrodactys* M.B. Ellis from India. *Indian Phytopathology* 59(1): 91–93.
- Mel'nik V, Hüseyin E, Selçuk F. 2004. Contribution to the studying of micromycetes in several Black Sea provinces of Turkey. *Novitates Systematicae Plantarum non Vascularum* 37: 133–148.
- Morelet M. 1985. Nouveaux binomes chez les Deuteromycotina. *Annales de la Société des Sciences Naturelles et d'Archéologie de Toulon et du Var* 37(4): 233–234.
- Seifert K, Morgan-Jones G, Gams W, Kendrick B. 2011. The genera of hyphomycetes. *CBS Biodiversity Series* 9. 997 p.
- Selçuk F, Ekici K. 2014. A new species of *Manoharachariella* (hyphomycetes) from Central Anatolia, Turkey. *Mycosphere* 5: 419–423. <http://dx.doi.org/10.5943/mycosphere/5/3/4>
- Selçuk F, Hüseyin E. 2014. New records of microfungi from Mountain Strandzha in Turkey (South-Eastern Europe). II. *Mikologiya i Fitopatologiya* 48(3): 202–208.
- Selçuk F, Hüseyin E, Bitmiş K. 2003. Some materials on mitosporic fungi from Turkey. II. *Coelomycetes*. *Botanica Lithuanica* 9(2): 161–170.
- Selçuk F, Erdoğan M, Akgül H, Hüseyin E. 2009. The genus *Septoria* Sacc. in Turkey. *Mycopath* 7: 21–28.
- Selçuk F, Hüseyin E, Bulbul AS. 2012a. Second record of *Ramularia hypericicola* – collected in Turkey on a new host. *Mycotaxon* 119: 369–372. <http://dx.doi.org/10.5248/119.369>
- Selçuk F, Hüseyin E, Gündoğan T, Özkan E. 2012b. Microfungi of genus *Phyllosticta* Pers. determined in Turkey ecosystems. 184, in: *Proceedings of the Ecological Symposium 2012*. 03–05 May 2012. Kilis 7 Aralık University. Kilis, Turkey. [In Turkish].
- Selçuk F, Hüseyin E, Şahin A, Cebeci CC. 2014. Hyphomycetous fungi in several forest ecosystems of Black Sea provinces of Turkey. *Mycosphere* 5: 334–344. <http://dx.doi.org/10.5943/mycosphere/5/2/7>
- Sutton BC. 1980. *The coelomycetes: fungi imperfecti with pycnidia, acervuli and stromata*. CABI Publishing, Kew, UK.
- Vasighzadeh A, Zafari D, Selçuk F, Hüseyin E, Kurşat M, Lutz M, Piątek M. 2014. Discovery of *Thecaphora schwarzmaniana* on *Rheum ribes* in Iran and Turkey: implications for the diversity and phylogeny of leaf smut on rhubarbs. *Mycol Progress* 13: 881–892. <http://dx.doi.org/10.1007/s11557-014-0972-4>