

---

# MYCOTAXON

<http://dx.doi.org/10.5248/130.11>

Volume 130, pp. 11–15

January–March 2015

---

## Additions to rust and chytrid pathogens of Turkey

CUMALI ÖZASLAN<sup>1\*</sup>, MAKBULE ERDOĞDU<sup>2</sup>,  
ELŞAD HÜSEYİN<sup>2</sup> & ZEKIYE SULUDERE<sup>3</sup>

<sup>1</sup>Dicle University, Faculty of Agriculture, Department of Plant Protection, Diyarbakır-Turkey

<sup>2</sup>Ahi Evran University, Faculty of Science and Literature, Department of Biology,  
Kırşehir-Turkey

<sup>3</sup>Gazi University, Faculty of Science, Department of Biology, Ankara-Turkey

\*CORRESPONDENCE TO: [cumali.ozaslan@dicle.edu.tr](mailto:cumali.ozaslan@dicle.edu.tr)


**ABSTRACT** — *Uromyces bornmuelleri* on *Bongardia chrysogonum* and *Physoderma maculare* on *Alisma plantago-aquatica* have been recently discovered in Turkey. Morphological data obtained by light and scanning electron microscopy of identified fungi are presented.

**KEY WORDS** — new records, *Chytridiomycota*, *Basidiomycota*

### Introduction

The *Pucciniaceae* are the largest family within the *Uredinales*, parasitising almost all major angiosperm orders, with some primarily herbaceous plant families such as *Asteraceae* and *Poaceae* being most commonly infected (van der Merwe et al. 2007). *Uromyces* (Link) Unger was raised to generic rank by Unger (1832), and is typified by *U. appendiculatus* on *Phaseolus*. Seventy-four species of *Uromyces* have been registered on 227 species of higher plants in Turkey (Bahçecioğlu & Kabaktepe 2012).

All known representatives of the genus *Physoderma* Wallr. (*Blastocladiomycetes*, *Physodermataceae*) are obligate parasites of vascular plants (Olson et al. 1980). The generalized life cycle of *Physoderma* is composed of two distinct and separate phases, and the ephemeral epibiotic sporangium with an endobiotic rhizoidal system is confined to a single host cell (Sparrow & Johns 1965). *Physoderma maculare*, the type species of the genus, on *Alisma* confirms Clinton's (1902) account of the epibiotic stage being produced from resting zoospores. The latter on mature host tissue may also give rise to the endobiotic stage, which bears resting spores. On seedlings, however, resting zoospores produce only epibiotic sporangia (Sparrow 1964).



FIGS 1–7. *Uromyces bornmuelleri*. 1: Deformation of host plant. 2: Telia on leaf, in situ. 3: Teliospores. 4–6: Telia and teliospores (SEM). 7: Teliospores (SEM).

### Materials & methods

Plant specimens infected with microfungi were collected from Adiyaman and Siirt provinces, Turkey and prepared according to established herbarium techniques. Host plants were identified using the Flora of Turkey and East Aegean Islands (Davis 1965–85). The fungal specimens were prepared from the host plants by obtaining thin sections. Measurements were made from tissues mounted in 5% KOH or tap water. Microscopical features were examined and microphotographs were made using a Leica

DM E light microscope. Thirty spores were measured for each sample. The microfungi were identified using relevant literature (De Toni 1888, Saccardo & Sydow 1902, Gonzalez Fragoso 1918, Kuprevich & Ulijanishchev 1975, van der Merwe et al. 2007). Species names follow Index Fungorum (2014). The examined specimens have been deposited in the mycological collection of the Department of Plant Protection, Faculty of Agriculture, Dicle University, Diyarbakır, Turkey (DUF-M).

For scanning electron microscopy (SEM), 8–10 mm squares of infected leaves were mounted on SEM stubs with double-sided adhesive tape, coated with gold using a Polaron SC 502 Sputter Coater, and examined with a Jeol JSM 6060 scanning electron microscope at 5–10 kV in the Electron Microscopy Unit, Faculty of Science, Gazi University (Turkey).

### Taxonomy

*Uromyces bornmuelleri* Magnus, Verh. Ges. Deutsch. Naturf. 65: 151.1893. Figs 1–7

SPERMAGONIA, AECIDIA and UREDINIA unknown. TELIA amphigenous, chestnut brown, in dense groups or scattered, 0.2–3 mm diam., causing leaf deformation, at first covered by the epidermis, later becoming erumpent, pulverulent. TELIOSPORES yellow or yellowish-brown, ovoid, globoid, sometimes oblong, 21–30 × 18–21.5 µm, rounded at the apex, rounded or sometimes attenuate at the base; wall 2–4 µm thick, pedicels hyaline, short, fragile.

SPECIMEN EXAMINED — TURKEY, ADIYAMAN PROVINCE, Center, Kahta, Gerger, Sincik, Çelikhan, Samsat, in wheat field ecosystem, on the leaves of *Bongardia chrysogonum* (L.) Spach (*Berberidaceae*), 12.V.2009, C. Öztaşlan CÖ2029 (DUF-M).


*Uromyces bornmuelleri* has been reported from Azerbaijan on *Leontice chrysogonum* [≡ *Bongardia chrysogonum*] (Tranzschel 1939), and from Iraq and Cyprus on *Bongardia chrysogonum* (Georghiou 1957, Mathur 1972). *Uromyces bornmuelleri* is reported for the first time from Turkey.

*Physoderma maculare* Wallr., Fl. Crypt. Germ. 2: 192. 1833. Figs 8–13

RESTING SPORANGIA amphigenous, mostly hypophyllous and on leaf petioles, concentrated on leaves along the nerves, scattered, covered by the unbroken epidermis, ellipsoid, ovoid, sometimes punctiformis, 1–2 mm diam., chestnut brown. RESTING SPORES chestnut brown, globoid, ovoid to ellipsoid, content granular with more refractive globules, (23.5–)25.5–32 × 21.5–29 µm; with two distinct walls, the inner wall thin and colorless, the outer wall chestnut brown, 1–2 µm thick, smooth. ZOOSPORES not seen.

SPECIMEN EXAMINED — TURKEY, SIIRT PROVINCE, Karaca Village, 38°07 09 N 42°01 59 E, in rice field ecosystem, on the leaves of *Alisma plantago-aquatica* L. (*Alismataceae*), 8.VI.2012, C. Öztaşlan CÖ2035 (DUF-M).

The genus *Physoderma* and *P. maculare* are reported for the first time from Turkey. This fungus has been reported from Canada (Connors 1967), Ireland



FIGS 8–13. *Physoderma maculare*. 8: Resting sporangia on leaf, in situ. 9: Resting spores. 10: Resting sporangia on leaf (SEM). 11–13: Resting spores (SEM).

(Muskett & Malone 1984, as *Cladochytrium alismatis*), Poland (Czczuga et al. 2007), Siberia (De Toni 1888, as *Uredo alismatis*), Spain (Gonzalez Fragoso 1918, as *U. alismatis*), and U.S.A. (Farr et al. 1989).

#### Acknowledgements

The author thanks Prof. Dr. Z.M. Azbukina (Vladivostok, Russia) and Dr. Yuri Tykhonenko (Kiev, Ukraine) for critically reading the manuscript and serving as presubmission reviewers. Especially we are grateful to Dr. Shaun Pennycook (Auckland,

New Zealand) for detailed linguistic help and nomenclatural review of the manuscript. We would like to thank DÜBAP (Dicle University Research Projects Coordinator) for financial support of this Project (DÜBAP 12ZF71). We also thank Prof. Dr. Mecit Vural (Gazi University, Ankara) and Prof. Dr. A. Selçuk Ertekin (Dicle University, Diyarbakır) for his help with the identification of the hosts.

#### Literature cited

- Bahçecioglu Z, Kabaktepe Ş. 2012. Checklist of rust fungi in Turkey. [Mycotaxon 119: 494.] <http://www.mycotaxon.com/resources/checklists/Bahcecioglu-v119-checklist.pdf>
- Clinton GP. 1902. *Cladochytrium alismatis*. Botanical Gazette 33: 49–61.
- Connors IL. 1967. An annotated index of plant diseases in Canada and fungi recorded on plants in Alaska, Canada and Greenland. Research Branch, Canada Department of Agriculture, Publication 1251. 381 p.
- Czczuga B, Muszynska E, Godlewska A, Mazalska B. 2007. Aquatic fungi and straminipilous organisms on decomposing fragments of wetland plants. Mycologia Balcanica 4: 31–44.
- Davis PH (ed.). 1965–85. Flora of Turkey and East Aegean Islands. Vols 1–9. Edinburgh University Press, Edinburgh.
- De Toni GB. 1888. Sylloge ustilaginearum et uredinearum. Sylloge Fungorum 7(2): 449–882.
- Farr DF, Bills GF, Chamuris GP, Rossman AY. 1989. Fungi on plants and plant products in the United States. APS Press, St Paul MN. 1252 p.
- Georghiou GP, Papadopoulos C. 1957. A second list of Cyprus fungi. Government of Cyprus, Department of Agriculture.
- Gonzalez Fragoso R. 1918. La roya de los vegetales. Enumeracion y distribucion geografica de los Uredales. Conocidos hasta hoy en la Peninsula Iberica e Islas Baleares. Trabajos del Museo Nacional de Ciencias Naturales, Serie Botánica 15: 1–267.
- Index Fungorum. 2014. <http://www.indexfungorum.org/names/names.asp>. Accessed 26 May 2014.
- Kuprevich VF, Ulijanishchev V. 1975. Key to the rust fungi in SSSR. Minsk, Belarus: Nauka i Tekhnika.
- Mathur RS. 1972. Checklist of Iraqi *Uredinales*. Iraq Natural History Museum, Publication 29: 1–32.
- Muskett A, Malone J. 1984. Catalogue of Irish fungi – V. *Mastigomycotina* and *Zygomycotina*. Proceedings of the Royal Irish Academy, B. 84: 83–102.
- Olson LW, Edén UM, Lange L. 1980. The endobiotic thallus of *Physoderma maydis*, the causal agent of *Physoderma* disease of maize. Protoplasma 103: 1–16.
- Saccardo PA, Sydow P. 1902. Supplementum universale, pars V. Sylloge Fungorum 16. 1291 p.
- Sparrow FK. 1964. Observations on chytridiaceous parasites of phanerogams. XIII. *Physoderma maculare* Wallroth. Archiv für Mikrobiologie 48: 136–149.
- Sparrow FK, Johns RM. 1965. Observation on chytridiaceous parasites of phanerogams XVI. Notes on *Physoderma* from *Scirpeae*. Archiv für Mikrobiologie 51: 351–364.
- Tranzschel VG. 1939. Compendium of rusts of the USSR. Botanicheskii Institut Akademii Nauk SSSR. Moscow. 426 p.
- Unger F. 1832 [“1833”]. Die Exantheme der Pflanzen. Wien. 422 p.
- van der Merwe M, Ericson L, Walker J, Thrall PH, Burdon JJ. 2007. Evolutionary relationships among species of *Puccinia* and *Uromyces* (*Pucciniaceae*, *Uredinales*) inferred from protein coding gene phylogenies. Mycological Research 111: 163–175.