

GENERALIZED MORREY REGULARITY FOR PARABOLIC EQUATIONS WITH DISCONTINUOUS DATA

VAGIF S. GULIYEV^{1,2} AND LUBOMIRA G. SOFTOVA³

¹*Department of Mathematics, Ahi Evran University, Kirsehir, Turkey*

²*Institute of Mathematics and Mechanics, National Academy of Sciences of Azerbaijan,
Baku, Azerbaijan (vagif@guliyev.com)*

³*Department of Civil Engineering, Design, Construction and Environment,
Second University of Naples, Piazza Luigi Miraglia 2, 80138 Naples, Italy
(luba.softova@unina2.it)*

(Received 8 October 2012)

Abstract We prove continuity in generalized parabolic Morrey spaces $M_{p,\varphi}$ of sublinear operators generated by the parabolic Calderón–Zygmund operator and by the commutator of this operator with bounded mean oscillation (BMO) functions. As a consequence, we obtain a global $M_{p,\varphi}$ -regularity result for the Cauchy–Dirichlet problem for linear uniformly parabolic equations with vanishing mean oscillation (VMO) coefficients.

Keywords: generalized Morrey spaces; sublinear operators; Calderón–Zygmund integrals; BMO; VMO; Cauchy–Dirichlet problem

2010 Mathematics subject classification: Primary 35K20; 35B45; 35D35; 35R05; 42B20; 47B38

1. Introduction

The classical Morrey spaces $L_{p,\lambda}$ were originally introduced in [17] in order to prove local Hölder continuity of solutions to certain systems of partial differential equations (PDEs). A real-valued function f is said to belong to the Morrey space $L_{p,\lambda}(\mathbb{R}^n)$ with $p \in [1, \infty)$, $\lambda \in (0, n)$, provided that the norm

$$\|f\|_{L_{p,\lambda}(\mathbb{R}^n)} = \left(\sup_{(x,r) \in \mathbb{R}^n \times \mathbb{R}_+} \frac{1}{r^\lambda} \int_{\mathcal{B}_r(x)} |f(y)|^p dy \right)^{1/p}.$$

is finite. The main result connected with these spaces is the following celebrated lemma: let $|Df| \in L_{p,\lambda}$ even locally, with $\lambda < p$; u is then Hölder continuous of exponent $\alpha = 1 - \lambda/p$. This result has applications in the study of the regularity of the strong solutions to elliptic and parabolic PDEs and systems. In [5] Chiarenza and Frasca showed boundedness of the Hardy–Littlewood maximal operator, which allowed them to prove continuity in $L_{p,\lambda}(\mathbb{R}^n)$ of some classical integral operators. These operators appear in the representation formulae of the solutions of various linear PDEs. Thus, the results in [5]

Full addresses OK?
If not, please supply a full alternative.

Spelled out in abstract – OK? The term will also be spelled out at first instance in the body of the text. And similarly for ‘VMO’. Can the in-full versions go in the keywords too?

Please note: the subject classifications must be correct according to the 2010 classification – please confirm that they are OK as they are, and indicate any that should be secondary rather than primary classifications.

Please check that all differential (upright) ‘d’ and variable (italic) ‘d’ have been typeset correctly throughout and mark any for amendment.

Change OK?

allow us to study the regularity of the solutions of these operators in $L_{p,\lambda}$ (see [20, 21, 24] and the references therein). In [16] Mizuhara extended the concept of Morrey of integral average over a ball with a certain growth, taking a weight function $\omega(x, r): \mathbb{R}^n \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ instead of r^λ . Thus, he set various conditions on the weight function at the start of the study of the generalized Morrey spaces $L_{p,\omega}$. In [18] Nakai studied continuity of some classical integral operators in $L_{p,\omega}$ imposing the following doubling and integral conditions on ω :

$$C_1 \leq \frac{\omega(x, s)}{\omega(x, r)} \leq C_2, \quad r \leq s \leq 2r, \quad \int_r^\infty \frac{\omega(x, s)}{s^{n+1}} ds \leq C_3 \frac{\omega(x, r)}{r^n},$$

where the constants do not depend on s , r and x . Furthermore, in [19] Nakai extended the theory of Morrey spaces on homogeneous spaces (X, d, μ) endowed with a quasi-distance d and a non-negative measure μ . The generalized Morrey space is then defined to be the set of all $f \in L^1_{\text{loc}}(X)$ such that

$$\|f\|_{L_{p,\phi}(X)} = \sup_B \frac{1}{\phi(B)} \left(\frac{1}{\mu(B)} \int_B |f(x)|^p d\mu(x) \right)^{1/p},$$

where the suprema is taken over all balls $B = B(a, r)$ with respect to the quasi-distance d . The weight function $\phi(B) = \phi(a, r)$ satisfies the integral condition

$$\int_r^\infty \frac{\phi(a, s)}{s} ds \leq C\phi(a, r) \quad \forall a \in X, \quad r > 0. \quad (1.1)$$

As a consequence, the boundedness of the Hardy–Littlewood maximal function and the Calderón–Zygmund singular integral operators in Morrey-type spaces on spaces of homogeneous type hold. Some applications of these operators in the regularity theory of partial differential equations are presented in [23, 25, 26]. Therein, the second author obtained global $L_{p,\omega}$ -regularity for elliptic and parabolic boundary-value problems. The approach is based on explicit representation formulae for the higher-order derivatives of the solution and estimates for the above-mentioned operators.

Other generalizations of the Morrey spaces are considered in [2, 8, 9, 12]. In these works Guliyev *et al.* studied the continuity of known integral operators acting from one Morrey-type space, $M_{p,\varphi_1}(\mathbb{R}^n)$, to another, $M_{p,\varphi_2}(\mathbb{R}^n)$, where the couple (φ_1, φ_2) satisfies the condition

$$\int_r^\infty \frac{\text{ess inf}_{s < \zeta < \infty} \varphi_1(x, \zeta) \zeta^{n/p}}{s^{n/p+1}} ds \leq C\varphi_2(x, r), \quad (x, r) \in \mathbb{R}^n \times \mathbb{R}_+. \quad (1.2)$$

The technique consists of obtaining some estimates for sublinear operators generated by various classical integral operators such as the Calderón–Zygmund operator, the commutator with bounded mean oscillation (BMO) functions, the Riesz potential, the Hardy–Littlewood maximal operator and others. We note that the condition (1.2) is weaker than (1.1). Indeed, if (1.1) holds and $\varphi_1 = \varphi_2 = \varphi$, then

$$\int_r^\infty \frac{\text{ess inf}_{s < \zeta < \infty} \varphi(x, \zeta) \zeta^{n/p}}{s^{n/p+1}} ds \leq \int_r^\infty \frac{\varphi(x, s)}{s} ds \leq C\varphi(x, r).$$

The following example shows that there exist functions satisfying (1.2) but not (1.1).

‘Morrey spaces’?
Please clarify this sentence.

Changes to sentence OK?

Changes to sentence OK?

Example 1.1. For $\beta \in (0, n/p)$, consider the weight function

$$\varphi(r) = r^{\beta-n/p} \left| \sin \left(\max \left\{ 1, \frac{\pi}{r} \right\} \right) \right|.$$

If $r \in (0, \pi)$, then $\text{ess inf}_{r < \zeta < \infty} \varphi(\zeta) \zeta^{n/p} = 0$. If $r \in (\pi, \infty)$, then $\text{ess inf}_{r < \zeta < \infty} \varphi(\zeta) \zeta^{n/p} = r^\beta \sin 1$. Then,

Slight change to wording of these sentences to avoid a bad line break – OK?

$$\int_r^\infty \frac{\text{ess inf}_{s < \zeta < \infty} \varphi(\zeta) \zeta^{n/p}}{s^{n/p+1}} ds = \begin{cases} 0, & r \in (0, \pi), \\ r^{\beta-n/p} \sin 1, & r \in (\pi, \infty), \end{cases} \leq C\varphi(r).$$

The function φ does not satisfy the condition (1.1).

In [11] we applied the results obtained in [2] to the study of the global $M_{p,\varphi}$ -regularity for the Dirichlet problem for linear uniformly elliptic equations with vanishing mean oscillation (VMO) coefficients. In the present work we extend our study over singular integral operators with a parabolic-type kernel. As a by-product we obtain regularity result for strong solutions of parabolic boundary-value problems. The unique strong solvability of the problem under consideration is guaranteed by [3]. Furthermore, its regularity has been studied in the frameworks of the Morrey, Morrey-type and weighted Lebesgue spaces, in [21], [25] and [10], respectively, while in [27] we deal with the oblique derivative problem in $M_{p,\varphi}$ spaces. Our approach is based on estimates for sublinear operators generated by singular integrals with parabolic kernels (see § 3). The last ones enter in the interior representation formula of the derivatives $D_{ij}u$ of the solution of (2.1). In § 4 we establish continuity of sublinear operators generated by non-singular integral operators. A similar result also holds for the commutators of these operators with BMO functions. The last integrals are included in the boundary representation formula for $D_{ij}u$, and allow us to obtain a local *a priori* estimate near the boundary. The global *a priori* estimate for u is obtained in § 6.

Word added – OK? ‘a regularity result’ or ‘regularity results’?

This sentence is unclear. Do you mean ‘The last estimates in § 3 are included in the interior representation...’? Please clarify.

Changes to sentence OK?

The following notation is used throughout this paper:

- $x = (x', t), y = (y', \tau) \in \mathbb{R}^{n+1} = \mathbb{R}^n \times \mathbb{R}, \mathbb{R}_+^{n+1} = \mathbb{R}^n \times \mathbb{R}_+$;
- $x = (x'', x_n, t) \in \mathbb{D}_+^{n+1} = \mathbb{R}^{n-1} \times \mathbb{R}_+ \times \mathbb{R}_+, \mathbb{D}_-^{n+1} = \mathbb{R}^{n-1} \times \mathbb{R}_- \times \mathbb{R}_+$;
- $|\cdot|$ is the Euclidean metric, $|x| = (\sum_{i=1}^n x_i^2 + t^2)^{1/2}$;
- $\mathcal{B}_r(x') = \{y' \in \mathbb{R}^n : |x' - y'| < r\}, |\mathcal{B}_r| = Cr^n$;
- $\mathcal{I}_r(x) = \{y \in \mathbb{R}^{n+1} : |x' - y'| < r, |t - \tau| < r^2\}, |\mathcal{I}_r| = Cr^{n+2}$;
- \mathbb{S}^n is the unit sphere in \mathbb{R}^{n+1} ;
- $D_i u = \partial u / \partial x_i, Du = (D_1 u, \dots, D_n u), u_t = \partial u / \partial t$;
- $D_{ij} u = \partial^2 u / \partial x_i \partial x_j, D^2 u = \{D_{ij} u\}_{i,j=1}^n$ denotes the Hessian matrix of u ;

- for any $f \in L_p(A)$, $A \subset \mathbb{R}^{n+1}$, we write

$$\|f\|_{p,A} \equiv \|f\|_{L_p(A)} = \left(\int_A |f(y)|^p dy \right)^{1/p};$$

- the standard summation convention on repeated upper and lower indices is adopted;
- the symbol C is used for various positive constants and may change from one occurrence to another.

2. Definitions and statement of the problem

In the following, besides the standard parabolic metric $\varrho(x) = \max(|x'|, |t|^{1/2})$ we use the equivalent one $\rho(x) = ((|x'|^2 + \sqrt{|x'|^4 + 4t^2})/2)^{1/2}$ considered by Fabes and Riviere in [7] (see also [28]). The topology this induces consists of the ellipsoids

$$\mathcal{E}_r(x) = \left\{ y \in \mathbb{R}^{n+1} : \frac{|x' - y'|^2}{r^2} + \frac{|t - \tau|^2}{r^4} < 1 \right\}, \quad |\mathcal{E}_r| = Cr^{n+2}, \quad \mathcal{E}_1(x) \equiv \mathcal{B}_1(x).$$

It is easy to see that the metrics $\rho(\cdot)$ and $\varrho(\cdot)$ are equivalent. In fact, for each \mathcal{E}_r there exist parabolic cylinders $\underline{\mathcal{I}}$ and $\overline{\mathcal{I}}$ with measure comparable to r^{n+2} such that $\underline{\mathcal{I}} \subset \mathcal{E}_r \subset \overline{\mathcal{I}}$. In what follows, all estimates obtained over ellipsoids also hold true over parabolic cylinders, and we use this property without explicit references (see [28]).

Let $\Omega \subset \mathbb{R}^n$ be a bounded $C^{1,1}$ -domain, let $Q = \Omega \times (0, T)$, and let $T > 0$ be a cylinder in \mathbb{R}_+^{n+1} . We give the definitions of the functional spaces that we are going to use.

Definition 2.1. Let $a \in L_1^{\text{loc}}(\mathbb{R}^{n+1})$ and let $a_{\mathcal{E}_r} = |\mathcal{E}_r|^{-1} \int_{\mathcal{E}_r} a(y) dy$ be the mean integral of a . Define

$$\eta_a(R) = \sup_{r \leq R} \frac{1}{|\mathcal{E}_r|} \int_{\mathcal{E}_r} |f(y) - f_{\mathcal{E}_r}| dy \quad \text{for every } R > 0,$$

where \mathcal{E}_r ranges over all ellipsoids in \mathbb{R}^{n+1} . We say that the following hold.

- $a \in \text{BMO}$ (*bounded mean oscillation* [13]) provided that

$$\|a\|_* = \sup_{R>0} \eta_a(R)$$

is finite. The quantity $\|\cdot\|_*$ is a norm in a BMO modulo constant function under which BMO is a Banach space.

- $a \in \text{VMO}$ (*vanishing mean oscillation* [22]) if $a \in \text{BMO}$ and

$$\lim_{R \rightarrow 0} \eta_a(R) = 0.$$

The quantity $\eta_a(R)$ is called the VMO-modulus of a .

Changes to sentence OK?

I have changed 'cf.' to 'see' throughout, in keeping with journal style. Please mark where it actually means 'compare with'.

Changes to sentence OK?

Changes to sentence OK?

Word added – OK?

Word added – OK?

For any bounded cylinder Q we define $\text{BMO}(Q)$ and $\text{VMO}(Q)$ taking $a \in L_1(Q)$ and $Q_r(x) = Q \cap \mathcal{E}_r(x)$, $x \in Q$, instead of \mathcal{E}_r in the definition above.

According to [1, 14], having a function $a \in \text{BMO}(Q)$ or $\text{VMO}(Q)$, it is possible to extend the function in the whole of \mathbb{R}^{n+1} preserving its BMO-norm or VMO-modulus, respectively. In the following we use this property without explicit references. Any bounded uniformly continuous (BUC) function f with modulus of continuity $\omega_f(R)$ belongs to VMO with $\eta_f(R) = \omega_f(R)$. Besides that, BMO and VMO also contain discontinuous functions, and the following example shows the inclusion $W_{1,n+2}(\mathbb{R}^{n+1}) \subset \text{VMO} \subset \text{BMO}$.

Changes to sentence OK?

Example 2.2. We have that $f(x) = |\log \rho(x)| \in \text{BMO} \setminus \text{VMO}$, $\sin f(x) \in \text{BMO} \cap L_\infty(\mathbb{R}^{n+1})$, $f_\alpha(x) = |\log \rho(x)|^\alpha \in \text{VMO}$ for any $\alpha \in (0, 1)$,

Changes to sentence OK?

$$\begin{aligned} f_\alpha &\in W_{1,n+2}(\mathbb{R}^{n+1}) \quad \text{for } \alpha \in (0, 1 - 1/(n+2)), \\ f_\alpha &\notin W_{1,n+2}(\mathbb{R}^{n+1}) \quad \text{for } \alpha \in [1 - 1/(n+2), 1). \end{aligned}$$

Definition 2.3. Let $\varphi: \mathbb{R}^{n+1} \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be a measurable function and let $p \in [1, \infty)$. The generalized parabolic Morrey space $M_{p,\varphi}(\mathbb{R}^{n+1})$ consists of all $f \in L_p^{\text{loc}}(\mathbb{R}^{n+1})$ such that

$$\|f\|_{p,\varphi;\mathbb{R}^{n+1}} = \sup_{(x,r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+} \varphi(x,r)^{-1} \left(r^{-(n+2)} \int_{\mathcal{E}_r(x)} |f(y)|^p dy \right)^{1/p} < \infty.$$

The space $M_{p,\varphi}(Q)$ consists of $L_p(Q)$ functions provided that the following norm is finite:

$$\|f\|_{p,\varphi;Q} = \sup_{(x,r) \in Q \times \mathbb{R}_+} \varphi(x,r)^{-1} \left(r^{-(n+2)} \int_{Q_r(x)} |f(y)|^p dy \right)^{1/p}.$$

The generalized weak parabolic Morrey space $WM_{1,\varphi}(\mathbb{R}^{n+1})$ consists of all measurable functions such that

$$\|f\|_{WM_{1,\varphi}(\mathbb{R}^{n+1})} = \sup_{(x,r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+} \varphi(x,r)^{-1} r^{-(n+2)} \|f\|_{WL_1(\mathcal{E}_r(x))},$$

where WL_1 denotes the weak L_1 space.

The generalized Sobolev–Morrey space $W_{p,\varphi}^{2,1}(Q)$, $p \in [1, \infty)$, consists of all Sobolev functions $u \in W_p^{2,1}(Q)$ with distributional derivatives $D_t^l D_x^s u \in M_{p,\varphi}(Q)$, $0 \leq 2l + |s| \leq 2$, endowed by the norm

$$\|u\|_{W_{p,\varphi}^{2,1}(Q)} = \|u_t\|_{p,\varphi;Q} + \sum_{|s| \leq 2} \|D^s u\|_{p,\varphi;Q}.$$

We also define the space

$$\overset{\circ}{W}_{p,\varphi}^{2,1}(Q) = \{u \in W_{p,\varphi}^{2,1}(Q) : u(x) = 0, x \in \partial Q\}, \quad \|u\|_{\overset{\circ}{W}_{p,\varphi}^{2,1}(Q)} = \|u\|_{W_{p,\varphi}^{2,1}(Q)},$$

where ∂Q means the parabolic boundary $\Omega \cup (\partial\Omega \times (0, T))$.

'a.a.' spelled out at first instance as 'almost all' here, and assumed to stand for that throughout. Please mark if it should be 'almost anywhere'.

We consider the linear Cauchy–Dirichlet problem

$$u_t - a^{ij}(x)D_{ij}u(x) = f(x) \text{ for almost all (a.a.) } x \in Q, \quad u \in \mathring{W}_{p,\varphi}^{2,1}(Q), \quad (2.1)$$

where the coefficient matrix $\mathbf{a}(x) = \{a^{ij}(x)\}_{i,j=1}^n$ satisfies

$$\left. \begin{aligned} \exists \Lambda > 0: \Lambda^{-1}|\xi|^2 \leq a^{ij}(x)\xi_i\xi_j \leq \Lambda|\xi|^2 \text{ for a.a. } x \in Q, \forall \xi \in \mathbb{R}^n, \\ a^{ij}(x) = a^{ji}(x), \text{ which implies that } a^{ij} \in L_\infty(Q). \end{aligned} \right\} \quad (2.2)$$

Changes to the following display OK?

Theorem 2.4 (main result). *Let $\mathbf{a} \in \text{VMO}(Q)$ with $\eta_{\mathbf{a}} = \sum_{i,j=1}^n \eta_{a_{ij}}$ satisfy (2.2), and, for each $p \in (1, \infty)$, let $u \in \mathring{W}_{p,\varphi}^{2,1}(Q)$ be a strong solution of (2.1). If $f \in M_{p,\varphi}(Q)$ with $\varphi(x, r)$ being a measurable positive function satisfying*

Word added – OK?

$$\int_r^\infty \left(1 + \ln \frac{s}{r}\right) \frac{\text{ess inf}_{s < \zeta < \infty} \varphi(x, \zeta) \zeta^{(n+2)/p}}{s^{(n+2)/p+1}} ds \leq C\varphi(x, r), \quad (x, r) \in Q \times \mathbb{R}_+, \quad (2.3)$$

then $u \in \mathring{W}_{p,\varphi}^{2,1}(Q)$ and

$$\|u\|_{\mathring{W}_{p,\varphi}^{2,1}(Q)} \leq C\|f\|_{p,\varphi;Q}, \quad (2.4)$$

with $C = C(n, p, \Lambda, \partial\Omega, T, \eta_{\mathbf{a}}, \|\mathbf{a}\|_{\infty;Q})$.

3. Sublinear operators generated by parabolic singular integrals in generalized Morrey spaces

Let $f \in L_1(\mathbb{R}^{n+1})$ be a function with a compact support and let $a \in \text{BMO}$. For any $x \notin \text{supp } f$ define the sublinear operators T and T_a such that

Word added – OK?

$$|Tf(x)| \leq C \int_{\mathbb{R}^{n+1}} \frac{|f(y)|}{\rho(x-y)^{n+2}} dy, \quad (3.1)$$

$$|T_a f(x)| \leq C \int_{\mathbb{R}^{n+1}} |a(x) - a(y)| \frac{|f(y)|}{\rho(x-y)^{n+2}} dy. \quad (3.2)$$

Suppose, in addition, that both the operators are bounded in $L_p(\mathbb{R}^{n+1})_2$, satisfying the estimates

Change OK?

$$\|Tf\|_{p;\mathbb{R}^{n+1}} \leq C\|f\|_{p;\mathbb{R}^{n+1}}, \quad \|T_a f\|_{p;\mathbb{R}^{n+1}} \leq C\|a\|_* \|f\|_{p;\mathbb{R}^{n+1}}, \quad (3.3)$$

with constants independent of a and f . The following known result concerns the Hardy operator $\text{Hg}(r) = (1/r) \int_0^r g(s) ds$, $r > 0$.

Operator name changed to roman – OK?

Theorem 3.1 (Carro *et al.* [4]). *The inequality*

$$\text{ess sup}_{r>0} w(r) \text{Hg}(r) \leq A \text{ess sup}_{r>0} v(r)g(r) \quad (3.4)$$

holds for all non-increasing functions $g: \mathbb{R}_+ \rightarrow \mathbb{R}_+$ if and only if

$$A = C \sup_{r>0} \frac{w(r)}{r} \int_0^r \frac{ds}{\text{ess sup}_{0<\zeta<s} v(\zeta)} < \infty. \quad (3.5)$$

Lemma 3.2. Let $f \in L_p^{\text{loc}}(\mathbb{R}^{n+1})$, $p \in [1, \infty)$, be such that

$$\int_r^\infty s^{-(n+2)/p-1} \|f\|_{p; \mathcal{E}_s(x_0)} ds < \infty \quad \forall (x_0, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+, \quad (3.6)$$

and let T be a sublinear operator satisfying (3.1).

(i) If $p > 1$ and T is bounded on $L_p(\mathbb{R}^{n+1})$, then

$$\|Tf\|_{p; \mathcal{E}_r(x_0)} \leq Cr^{(n+2)/p} \int_{2r}^\infty s^{-(n+2)/p-1} \|f\|_{p; \mathcal{E}_s(x_0)} ds. \quad (3.7)$$

(ii) If $p = 1$ and T is bounded from $L_1(\mathbb{R}^{n+1})$ on $WL_1(\mathbb{R}^{n+1})$, then

$$\|Tf\|_{WL_1(\mathcal{E}_r(x_0))} \leq Cr^{n+2} \int_{2r}^\infty s^{-(n+3)} \|f\|_{1, \mathcal{E}_s(x_0)} ds, \quad (3.8)$$

where the constants are independent of r , x_0 and f .

Proof. (i) Fix a point $x_0 \in \mathbb{R}^{n+1}$ and consider an ellipsoid $\mathcal{E}_r(x_0)$. Define $2\mathcal{E}_r(x_0) = \mathcal{E}_{2r}(x_0)$, $\mathcal{E}_r^c(x_0) = \mathbb{R}^{n+1} \setminus \mathcal{E}_r(x_0)$ and consider the decomposition of f ,

$$f = f\chi_{2\mathcal{E}_r(x_0)} + f\chi_{2\mathcal{E}_r^c(x_0)} = f_1 + f_2.$$

Because of the (p, p) -boundedness of the operator T and $f_1 \in L_p(\mathbb{R}^{n+1})$ we have that

$$\|Tf_1\|_{p; \mathcal{E}_r(x_0)} \leq \|Tf_1\|_{p; \mathbb{R}^{n+1}} \leq C\|f_1\|_{p; \mathbb{R}^{n+1}} = C\|f\|_{p; 2\mathcal{E}_r(x_0)}.$$

It is easy to see that for arbitrary points $x \in \mathcal{E}_r(x_0)$ and $y \in 2\mathcal{E}_r^c(x_0)$ it holds that

$$\frac{1}{2}\rho(x_0 - y) \leq \rho(x - y) \leq \frac{3}{2}\rho(x_0 - y). \quad (3.9)$$

Applying (3.1), (3.9), the Fubini theorem and the Hölder inequality to Tf_2 , we get that

$$\begin{aligned} |Tf_2(x)| &\leq C \int_{2\mathcal{E}_r^c(x_0)} \frac{|f(y)|}{\rho(x_0 - y)^{n+2}} dy \\ &\leq C \int_{2\mathcal{E}_r^c(x_0)} |f(y)| \left(\int_{\rho(x_0 - y)}^\infty \frac{ds}{s^{n+3}} \right) dy \\ &\leq C \int_{2r}^\infty \left(\int_{2r \leq \rho(x_0 - y) < s} |f(y)| dy \right) \frac{ds}{s^{n+3}} \\ &\leq C \int_{2r}^\infty \left(\int_{\mathcal{E}_s(x_0)} |f(y)| dy \right) \frac{ds}{s^{n+3}} \\ &\leq C \int_{2r}^\infty \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}}. \end{aligned}$$

Direct calculations give

$$\|Tf_2\|_{p; \mathcal{E}_r(x_0)} \leq Cr^{(n+2)/p} \int_{2r}^\infty \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}}, \quad (3.10)$$

Changes to the style of the following itemized list – OK?

Word added – OK?

Word added – OK?

Changes to sentence OK?

which holds for all $p \in [1, \infty)$. Thus,

$$\|Tf\|_{p; \mathcal{E}_r(x_0)} \leq C \left(\|f\|_{p; 2\mathcal{E}_r(x_0)} + r^{(n+2)/p} \int_{2r}^{\infty} \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}} \right). \quad (3.11)$$

On the other hand,

$$\|f\|_{p, 2\mathcal{E}_r(x_0)} \leq Cr^{(n+2)/p} \int_{2r}^{\infty} \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}}, \quad (3.12)$$

which, unified with (3.11), gives (3.7).

(ii) Let $f \in L_1(\mathbb{R}^{n+1})$; the weak $(1, 1)$ -boundedness of T implies that

Changes to sentence OK?

$$\begin{aligned} \|Tf_1\|_{WL_1(\mathcal{E}_r(x_0))} &\leq \|Tf_1\|_{WL_1(\mathbb{R}^{n+1})} \\ &\leq C\|f_1\|_{1, \mathbb{R}^{n+1}} \\ &= C\|f\|_{1, 2\mathcal{E}_r(x_0)} \\ &\leq Cr^{n+2} \int_{2r}^{+\infty} \|f\|_{1, \mathcal{E}_s(x_0)} \frac{ds}{s^{n+3}}, \end{aligned}$$

which, unified with (3.10), gives (3.8). \square

Theorem 3.3. *Let $p \in [1, \infty)$, let $\varphi(x, r)$ be a measurable positive function satisfying*

$$\int_r^{\infty} \frac{\text{ess inf}_{s < \zeta < \infty} \varphi(x, \zeta) \zeta^{(n+2)/p}}{s^{(n+2)/p+1}} ds \leq C\varphi(x, r) \quad \forall (x, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+, \quad (3.13)$$

and let T be a sublinear operator satisfying (3.1).

Word added – OK?

(i) If $p > 1$ and T is bounded on $L_p(\mathbb{R}^{n+1})$, then T is bounded on $M_{p, \varphi}(\mathbb{R}^{n+1})$, and

Changes to sentence OK?

$$\|Tf\|_{p, \varphi; \mathbb{R}^{n+1}} \leq C\|f\|_{p, \varphi; \mathbb{R}^{n+1}}. \quad (3.14)$$

(ii) If $p = 1$ and T is bounded from $L_1(\mathbb{R}^{n+1})$ to $WL_1(\mathbb{R}^{n+1})$, then it is bounded from $M_{1, \varphi}(\mathbb{R}^{n+1})$ to $WM_{1, \varphi}(\mathbb{R}^{n+1})$, and

Changes to sentence OK?

$$\|Tf\|_{WM_{1, \varphi}(\mathbb{R}^{n+1})} \leq C\|f\|_{1, \varphi; \mathbb{R}^{n+1}} \quad (3.15)$$

with constants independent on f .

Proof. (i) By Lemma 3.2, we have that

$$\begin{aligned} \|Tf\|_{p, \varphi; \mathbb{R}^{n+1}} &\leq C \sup_{(x, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+} \varphi(x, r)^{-1} \int_r^{\infty} \|f\|_{p; \mathcal{E}_s(x)} \frac{ds}{s^{(n+2)/p+1}} \\ &= C \sup_{(x, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+} \varphi(x, r)^{-1} \int_0^{r^{-(n+2)/p}} \|f\|_{p; \mathcal{E}_{s^{-p/(n+2)}}(x)} ds \\ &= C \sup_{(x, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+} \varphi(x, r^{-p/(n+2)})^{-1} \int_0^r \|f\|_{p; \mathcal{E}_{s^{-p/(n+2)}}(x)} ds. \end{aligned}$$

Applying Theorem 3.1 with

$$w(r) = v(r) = r\varphi(x, r^{-p/(n+2)})^{-1}, \quad g(r) = \|f\|_{p; \mathcal{E}_{r^{-p/(n+2)}}(x)},$$

$$\text{Hg}(r) = r^{-1} \int_0^r \|f\|_{p; \mathcal{E}_{s^{-p/(n+2)}}(x)} \, ds,$$

where the condition (3.5) is equivalent to (3.13), we obtain (3.14).

(ii) The estimate follows after using (3.8) and (3.4):

$$\begin{aligned} \|Tf\|_{WM_{1,\varphi}(\mathbb{R}^{n+1})} &\leq C \sup_{(x_0, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+} \varphi(x_0, r)^{-1} \int_r^\infty \|f\|_{1, \mathcal{E}_s(x_0)} \frac{ds}{s^{n+3}} \\ &= C \sup_{(x_0, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+} \varphi(x_0, r^{-1/(n+2)})^{-1} \int_0^r \|f\|_{1, \mathcal{E}_{s^{-1/(n+2)}}(x_0)} \, ds \\ &\leq C \sup_{(x_0, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+} \varphi(x_0, r^{-1/(n+2)})^{-1} r \|f\|_{1, \mathcal{E}_{r^{-1/(n+2)}}(x_0)} \\ &= C \|f\|_{1, \varphi; \mathbb{R}^{n+1}}. \end{aligned}$$

□

Our next step is to show boundedness of T_a in $M_{p,\varphi}(\mathbb{R}^{n+1})$. For this we recall some properties of the BMO functions.

Lemma 3.4 (John–Nirenberg lemma [3, Lemma 2.8]). *Let $a \in \text{BMO}$ and let $p \in [1, \infty)$. Then, for any \mathcal{E}_r ,*

$$\left(\frac{1}{|\mathcal{E}_r|} \int_{\mathcal{E}_r} |a(y) - a_{\mathcal{E}_r}|^p \, dy \right)^{1/p} \leq C(p) \|a\|_*$$

Changes to sentence OK?

holds.

As an immediate consequence of Lemma 3.4 we get the following property.

Corollary 3.5. *Let $a \in \text{BMO}$; then, for all $0 < 2r < s$,*

$$|a_{\mathcal{E}_r} - a_{\mathcal{E}_s}| \leq C(n) \left(1 + \ln \frac{s}{r} \right) \|a\|_* \quad (3.16)$$

Changes to sentence OK?

holds.

Proof. Since $s > 2r$, there exists $k \in \mathbb{N}$, $k \geq 1$, such that $2^k r < s \leq 2^{k+1} r$ and, hence, $k \ln 2 < \ln(s/r) \leq (k+1) \ln 2$. By [3, Lemma 2.9] we have that

Change OK?

$$\begin{aligned} |a_{\mathcal{E}_s} - a_{\mathcal{E}_r}| &\leq |a_{2^k \mathcal{E}_r} - a_{\mathcal{E}_r}| + |a_{2^k \mathcal{E}_r} - a_{\mathcal{E}_s}| \\ &\leq C(n)k \|a\|_* + \frac{1}{|2^k \mathcal{E}_r|} \int_{2^k \mathcal{E}_r} |a(y) - a_{\mathcal{E}_s}| \, dy \\ &\leq C(n) \left(k \|a\|_* + \frac{1}{|\mathcal{E}_s|} \int_{\mathcal{E}_s} |a(y) - a_{\mathcal{E}_s}| \, dy \right) \\ &< C(n) \left(\ln \frac{s}{r} + 1 \right) \|a\|_*. \end{aligned}$$

□

To estimate the norm of T_a we employ the same idea that we have used in the proof of Lemma 3.2.

Lemma 3.6. *Let $a \in \text{BMO}$ and let T_a be a bounded operator in $L_p(\mathbb{R}^{n+1})$, $p \in (1, \infty)_2$ satisfying (3.2) and (3.3). Suppose that, for any $f \in L_p^{\text{loc}}(\mathbb{R}^{n+1})$,* Change OK?

$$\int_r^\infty \left(1 + \ln \frac{s}{r}\right) \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}} < \infty \quad \forall (x_0, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+. \quad (3.17)$$

Then,

$$\|T_a f\|_{p; \mathcal{E}_r(x_0)} \leq C \|a\|_* r^{(n+2)/p} \int_{2r}^\infty \left(1 + \ln \frac{s}{r}\right) \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}}, \quad (3.18)$$

where C is independent of a , f , x_0 and r .

Proof. Fix a point $x_0 \in \mathbb{R}^{n+1}$ and consider the decomposition

$$f = f \chi_{2\mathcal{E}_r(x_0)} + f \chi_{2\mathcal{E}_r^c(x_0)} = f_1 + f_2.$$

Displayed to avoid
a bad line break -
OK?

Hence,

$$\|T_a f\|_{p; \mathcal{E}_r(x_0)} \leq \|T_a f_1\|_{p; \mathcal{E}_r(x_0)} + \|T_a f_2\|_{p; \mathcal{E}_r(x_0)},$$

and by (3.3) as in Lemma 3.2 we have that

$$\|T_a f_1\|_{p; \mathcal{E}_r(x_0)} \leq C \|a\|_* \|f\|_{p; 2\mathcal{E}_r(x_0)}. \quad (3.19)$$

On the other hand, because of (3.9) we can write

$$\begin{aligned} \|T_a f_2\|_{p; \mathcal{E}_r(x_0)} &\leq C \left(\int_{\mathcal{E}_r(x_0)} \left(\int_{2\mathcal{E}_r^c(x_0)} \frac{|a(x) - a(y)| |f(y)|}{\rho(x_0 - y)^{n+2}} dy \right)^p dx \right)^{1/p} \\ &\leq C \left(\int_{\mathcal{E}_r(x_0)} \left(\int_{2\mathcal{E}_r^c(x_0)} \frac{|a(y) - a_{\mathcal{E}_r(x_0)}| |f(y)|}{\rho(x_0 - y)^{n+2}} dy \right)^p dx \right)^{1/p} \\ &\quad + C \left(\int_{\mathcal{E}_r(x_0)} \left(\int_{2\mathcal{E}_r^c(x_0)} \frac{|a(x) - a_{\mathcal{E}_r(x_0)}| |f(y)|}{\rho(x_0 - y)^{n+2}} dy \right)^p dx \right)^{1/p} \\ &= I_1 + I_2. \end{aligned}$$

Applying (3.2), the Fubini theorem and the Hölder inequality as in Lemmas 3.2 and 3.4, we get

$$\begin{aligned} I_1 &\leq C r^{(n+2)/p} \left(\int_{2r}^\infty \int_{\mathcal{E}_s(x_0)} |a(y) - a_{\mathcal{E}_r(x_0)}| |f(y)| dy \right) \frac{ds}{s^{n+3}} \\ &\leq C r^{(n+2)/p} \left(\int_{2r}^\infty \int_{\mathcal{E}_s(x_0)} |a(y) - a_{\mathcal{E}_s(x_0)}| |f(y)| dy \right) \frac{ds}{s^{n+3}} \\ &\quad + C r^{(n+2)/p} \int_{2r}^\infty |a_{\mathcal{E}_r(x_0)} - a_{\mathcal{E}_s(x_0)}| \left(\int_{\mathcal{E}_s(x_0)} |f(y)| dy \right) \frac{ds}{s^{n+3}} \end{aligned}$$

$$\begin{aligned}
 &\leq C r^{(n+2)/p} \int_{2r}^{\infty} \left(\int_{\mathcal{E}_s(x_0)} |a(y) - a_{\mathcal{E}_s(x_0)}|^{p/(p-1)} dy \right)^{(p-1)/p} \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{n+3}} \\
 &\quad + C r^{(n+2)/p} \int_{2r}^{\infty} |a_{\mathcal{E}_r(x_0)} - a_{\mathcal{E}_s(x_0)}| \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}} \\
 &\leq C \|a\|_* r^{(n+2)/p} \int_{2r}^{\infty} \left(1 + \ln \frac{s}{r} \right) \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}}.
 \end{aligned}$$

In order to estimate I_2 we note that

$$I_2 = \left(\int_{\mathcal{E}_r(x_0)} |a(x) - a_{\mathcal{E}_r(x_0)}|^p dx \right)^{1/p} \int_{2\mathcal{E}_r^c(x_0)} \frac{|f(y)|}{\rho(x_0 - y)^{n+2}} dy.$$

By Lemma 3.4 and (3.10) we get that

$$\begin{aligned}
 I_2 &\leq C \|a\|_* r^{(n+2)/p} \int_{2\mathcal{E}_r^c(x_0)} \frac{|f(y)|}{\rho(x_0 - y)^{n+2}} dy \\
 &\leq C \|a\|_* r^{(n+2)/p} \int_{2r}^{\infty} \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}}.
 \end{aligned}$$

Summing up (3.19), I_1 and I_2 we get that

$$\|T_a f\|_{p; \mathcal{E}_r(x_0)} \leq C \|a\|_* \left(\|f\|_{p; 2\mathcal{E}_r(x_0)} + r^{(n+2)/p} \int_{2r}^{\infty} \left(1 + \ln \frac{s}{r} \right) \|f\|_{p; \mathcal{E}_s(x_0)} \frac{ds}{s^{(n+2)/p+1}} \right),$$

and the statement follows after applying (3.12). \square

Theorem 3.7. *Let $p \in (1, \infty)$ and $\varphi(x, r)$ be measurable positive functions such that*

$$\int_r^{\infty} \left(1 + \ln \frac{s}{r} \right) \frac{\text{ess inf}_{s < \zeta < \infty} \varphi(x, \zeta) \zeta^{(n+2)/p}}{s^{(n+2)/p+1}} ds \leq C \varphi(x, r) \quad \forall (x, r) \in \mathbb{R}^{n+1} \times \mathbb{R}_+, \quad (3.20)$$

where C is independent of x and r . Suppose that $a \in \text{BMO}$ and let T_a be a sublinear operator satisfying (3.2). If T_a is bounded in $L_p(\mathbb{R}^{n+1})$, then it is bounded in $M_{p, \varphi}(\mathbb{R}^{n+1})$, and

$$\|T_a f\|_{p, \varphi; \mathbb{R}^{n+1}} \leq C \|a\|_* \|f\|_{p, \varphi; \mathbb{R}^{n+1}}, \quad (3.21)$$

with a constant independent of a and f .

The statement of the theorem follows by Lemma 3.6 and Theorem 3.1 in the same manner as for Theorem 3.3.

Example 3.8. We have the functions

$$\varphi(x, r) = r^{\beta - (n+2)/p} \quad \text{and} \quad \varphi(x, r) = r^{\beta - (n+2)/p} \log^m(e + r)$$

with $0 < \beta < (n+2)/p$, and $m \geq 1$ are weight functions satisfying the condition (3.20).

Changes to this sentence and the next OK?

Change OK?

Changes to sentence OK?

4. Sublinear operators generated by non-singular integrals in generalized Morrey spaces

For any $x \in \mathbb{D}_+^{n+1}$, define $\tilde{x} = (x'', -x_n, t) \in \mathbb{D}_+^{n+1}$ and $x^0 = (x'', 0, 0) \in \mathbb{R}^{n-1}$. Consider the semi-ellipsoids $\mathcal{E}_r^+(x^0) = \mathcal{E}_r(x^0) \cap \mathbb{D}_+^{n+1}$. Let $f \in L_1(\mathbb{D}_+^{n+1})$, let $a \in \text{BMO}(\mathbb{D}_+^{n+1})_2$ and let \tilde{T} and \tilde{T}_a be sublinear operators such that

Changes to sentence OK?

$$|\tilde{T}f(x)| \leq C \int_{\mathbb{D}_+^{n+1}} \frac{|f(y)|}{\rho(\tilde{x} - y)^{n+2}} dy, \quad (4.1)$$

$$|\tilde{T}_a f(x)| \leq C \int_{\mathbb{D}_+^{n+1}} |a(x) - a(y)| \frac{|f(y)|}{\rho(\tilde{x} - y)^{n+2}} dy. \quad (4.2)$$

Suppose, in addition, that both the operators are bounded in $L_p(\mathbb{D}_+^{n+1})_2$ satisfying the estimates

Change OK?

$$\|\tilde{T}f\|_{p; \mathbb{D}_+^{n+1}} \leq C \|f\|_{p; \mathbb{D}_+^{n+1}}, \quad \|\tilde{T}_a f\|_{p; \mathbb{D}_+^{n+1}} \leq C \|a\|_* \|f\|_{p; \mathbb{D}_+^{n+1}}, \quad (4.3)$$

with constants independent of a and f . The following assertions can be proved in the same manner as in §3.

Lemma 4.1. *Let $f \in L_p^{\text{loc}}(\mathbb{D}_+^{n+1})$, $p \in (1, \infty)$, and, for all $(x^0, r) \in \mathbb{R}^{n-1} \times \mathbb{R}_+$,*

$$\int_r^\infty s^{-(n+2)/p-1} \|f\|_{p; \mathcal{E}_s^+(x^0)} ds < \infty. \quad (4.4)$$

If \tilde{T} is bounded on $L_p(\mathbb{D}_+^{n+1})$, then

$$\|\tilde{T}f\|_{p; \mathcal{E}_r^+(x^0)} \leq C r^{(n+2)/p} \int_{2r}^\infty s^{-(n+2)/p-1} \|f\|_{p; \mathcal{E}_s^+(x^0)} ds, \quad (4.5)$$

where the constant C is independent of r , x^0 and f .

Theorem 4.2. *Let φ be a weight function satisfying (3.13), and let \tilde{T} be a sublinear operator satisfying (4.1) and (4.3). It is bounded in $M_{p, \varphi}(\mathbb{D}_+^{n+1})$, $p \in (1, \infty)$, and*

Please clarify what 'it' refers to here.

$$\|\tilde{T}f\|_{p, \varphi; \mathbb{D}_+^{n+1}} \leq C \|f\|_{p, \varphi; \mathbb{D}_+^{n+1}}, \quad (4.6)$$

with a constant C independent of f .

Lemma 4.3. *Let $p \in (1, \infty)$, let $a \in \text{BMO}(\mathbb{D}_+^{n+1})$, and let \tilde{T}_a satisfy (4.2) and (4.3). Suppose that, for all $f \in L_p^{\text{loc}}(\mathbb{D}_+^{n+1})$,*

Changes to sentence OK?

$$\int_r^\infty \left(1 + \ln \frac{s}{r}\right) s^{-(n+2)/p-1} \|f\|_{p; \mathcal{E}_s^+(x^0)} ds < \infty \quad \forall (x^0, r) \in \mathbb{R}^{n-1} \times \mathbb{R}_+. \quad (4.7)$$

Then,

$$\|\tilde{T}_a f\|_{p; \mathcal{E}_r^+(x^0)} \leq C \|a\|_* r^{(n+2)/p} \int_{2r}^\infty \left(1 + \ln \frac{s}{r}\right) \|f\|_{p; \mathcal{E}_s^+(x^0)} \frac{ds}{s^{(n+2)/p+1}},$$

with a constant C independent of a , f , x^0 and r .

Theorem 4.4. *Let $p \in (1, \infty)$, let $a \in \text{BMO}(\mathbb{D}_+^{n+1})$, let $\varphi(x^0, r)$ be a weight function satisfying (3.20), and let \tilde{T}_a be a sublinear operator satisfying (3.2) and (3.3). Then, \tilde{T}_a is bounded in $M_{p,\varphi}(\mathbb{D}_+^{n+1})$, and*

Changes to sentence OK?

$$\|\tilde{T}_a f\|_{p,\varphi;\mathbb{D}_+^{n+1}} \leq C \|a\|_* \|f\|_{p,\varphi;\mathbb{D}_+^{n+1}}, \quad (4.8)$$

with a constant C independent of a and f .

5. Singular and non-singular integrals in generalized Morrey spaces

In the present section we apply the above results to Calderón–Zygmund-type operators with parabolic kernel. Since these operators are sublinear and bounded in $L_p(\mathbb{R}^{n+1})$, their continuity in $M_{p,\varphi}$ follows immediately.

Definition 5.1. A measurable function $\mathcal{K}(x, \xi): \mathbb{R}^{n+1} \times \mathbb{R}^{n+1} \setminus \{0\} \rightarrow \mathbb{R}$ is called a variable parabolic Calderón–Zygmund kernel if the following hold.

Changes to sentence OK?

(i) $\mathcal{K}(x, \cdot)$ is a parabolic Calderón–Zygmund kernel for a.a. $x \in \mathbb{R}^{n+1}$:

- (a) $\mathcal{K}(x, \cdot) \in C^\infty(\mathbb{R}^{n+1} \setminus \{0\})$,
- (b) $\mathcal{K}(x, \mu\xi) = \mu^{-(n+2)}\mathcal{K}(x, \xi)$ for all $\mu > 0$,
- (c) $\int_{\mathbb{S}^n} \mathcal{K}(x, \xi) \, d\sigma_\xi = 0$, $\int_{\mathbb{S}^n} |\mathcal{K}(x, \xi)| \, d\sigma_\xi < +\infty$.

(ii) $\|D_\xi^\beta \mathcal{K}\|_{L_\infty(\mathbb{R}^{n+1} \times \mathbb{S}^n)} \leq M(\beta) < \infty$ for every multi-index β .

Moreover,

$$|\mathcal{K}(x, x-y)| \leq \rho(x-y)^{-(n+2)} \left| \mathcal{K}\left(x, \frac{x-y}{\rho(x-y)}\right) \right| \leq \frac{M}{\rho(x-y)^{n+2}},$$

which means that the singular integrals

$$\left. \begin{aligned} \mathfrak{R}f(x) &= \text{PV} \int_{\mathbb{R}^{n+1}} \mathcal{K}(x, x-y) f(y) \, dy, \\ \mathfrak{C}[a, f](x) &= \text{PV} \int_{\mathbb{R}^{n+1}} \mathcal{K}(x, x-y) [a(y) - a(x)] f(y) \, dy \end{aligned} \right\} \quad (5.1)$$

are sublinear and bounded in $L_p(\mathbb{R}^{n+1})$ according to the results in [3, 7]. We note that any weight function φ satisfying (3.20) also satisfies (3.13) and, hence, the following holds as a simple application of the estimates proved in § 3.

Theorem 5.2. *For any $f \in M_{p,\varphi}(\mathbb{R}^{n+1})$ with (p, φ) as in Theorem 3.7 and $a \in \text{BMO}$, there exist constants depending on n, p and the kernel such that*

$$\|\mathfrak{R}f\|_{p,\varphi;\mathbb{R}^{n+1}} \leq C \|f\|_{p,\varphi;\mathbb{R}^{n+1}}, \quad \|\mathfrak{C}[a, f]\|_{p,\varphi;\mathbb{R}^{n+1}} \leq C \|a\|_* \|f\|_{p,\varphi;\mathbb{R}^{n+1}}. \quad (5.2)$$

Corollary 5.3. *Let Q be a cylinder in \mathbb{R}_+^{n+1} , let $f \in M_{p,\varphi}(Q)$, let $a \in \text{BMO}(Q)$, and let $\mathcal{K}(x, \xi): Q \times \mathbb{R}_+^{n+1} \setminus \{0\} \rightarrow \mathbb{R}$. The operators (5.1) are then bounded in $M_{p,\varphi}(Q)$, and*

Changes to this sentence and the next OK?

$$\|\bar{\mathcal{K}}f\|_{p,\varphi;Q} \leq C\|f\|_{p,\varphi;Q}, \quad \|\mathfrak{C}[a, f]\|_{p,\varphi;Q} \leq C\|a\|_*\|f\|_{p,\varphi;Q}, \quad (5.3)$$

with C independent of a and f .

Proof. Define the extensions

$$\bar{\mathcal{K}}(x, \xi) = \begin{cases} \mathcal{K}(x, \xi), & (x, \xi) \in Q \times \mathbb{R}_+^{n+1} \setminus \{0\}, \\ 0 & \text{elsewhere,} \end{cases} \quad \bar{f}(x) = \begin{cases} f(x), & x \in Q, \\ 0, & x \notin Q. \end{cases}$$

Denote by $\bar{\mathcal{K}}f$ the singular integral with a kernel $\bar{\mathcal{K}}$ and potential \bar{f} . Then,

$$|\bar{\mathcal{K}}f| \leq |\bar{\mathcal{K}}\bar{f}| \leq C \int_{\mathbb{R}^{n+1}} \frac{|\bar{f}(y)|}{\rho(x-y)^{n+2}} dy$$

and

$$\|\bar{\mathcal{K}}f\|_{p,\varphi;Q} \leq \|\bar{\mathcal{K}}\bar{f}\|_{p,\varphi;\mathbb{R}^{n+1}} \leq C\|\bar{f}\|_{p,\varphi;\mathbb{R}^{n+1}} = C\|f\|_{p,\varphi;Q}.$$

The estimate for the commutator follows in a similar way. \square

Corollary 5.4. *Let $a \in \text{VMO}$ and let (p, φ) be as in Theorem 3.7. For any $\varepsilon > 0$ there then exists a positive number $r_0 = r_0(\varepsilon, \eta_a)$ such that, for any $\mathcal{E}_r(x_0)$ with a radius $r \in (0, r_0)$ and all $f \in M_{p,\varphi}(\mathcal{E}_r(x_0))$,*

Word added – OK?

$$\|\mathfrak{C}[a, f]\|_{p,\varphi;\mathcal{E}_r(x_0)} \leq C\varepsilon\|f\|_{p,\varphi;\mathcal{E}_r(x_0)}, \quad (5.4)$$

where C is independent of ε , f , r and x_0 .

Proof. Since any VMO function can be approximated by BUC functions (see [6, 22]), for each $\varepsilon > 0$ there exist $r_0(\varepsilon, \eta_a)$ and $g \in \text{BUC}$ with modulus of continuity $\omega_g(r_0) < \varepsilon/2$ such that $\|a - g\|_* < \varepsilon/2$. Fixing $\mathcal{E}_r(x_0)$ with $r \in (0, r_0)$, we define the function

Changes to sentence OK?

$$h(x) = \begin{cases} g(x), & x \in \mathcal{E}_r(x_0), \\ g\left(x_0 + r\frac{x' - x'_0}{\rho(x - x_0)}, t_0 + r^2\frac{t - t_0}{\rho^2(x - x_0)}\right), & x \in \mathcal{E}_r^c(x_0), \end{cases}$$

such that $h \in \text{BUC}(\mathbb{R}^{n+1})$ and $\omega_h(r_0) \leq \omega_g(r_0) < \varepsilon/2$. Hence,

$$\begin{aligned} \|\mathfrak{C}[a, f]\|_{p,\varphi;\mathcal{E}_r(x_0)} &\leq \|\mathfrak{C}[a - g, f]\|_{p,\varphi;\mathcal{E}_r(x_0)} + \|\mathfrak{C}[g, f]\|_{p,\varphi;\mathcal{E}_r(x_0)} \\ &\leq C\|a - g\|_*\|f\|_{p,\varphi;\mathcal{E}_r(x_0)} + \|\mathfrak{C}[h, f]\|_{p,\varphi;\mathcal{E}_r(x_0)} \\ &< C\varepsilon\|f\|_{p,\varphi;\mathcal{E}_r(x_0)}. \end{aligned}$$

\square

For any $x' \in \mathbb{R}_+^n$ and any fixed $t > 0$, define the *generalized reflection*

$$\mathcal{T}(x) = (\mathcal{T}'(x), t), \quad \mathcal{T}'(x) = x' - 2x_n \frac{\mathbf{a}^n(x', t)}{a^{nn}(x', t)}, \quad (5.5)$$

where $\mathbf{a}^n(x)$ is the last row of the coefficients matrix $\mathbf{a}(x)$ of (2.1). The function $\mathcal{T}'(x)$ maps \mathbb{R}_+^n into \mathbb{R}_-^n , and the kernel $\mathcal{K}(x, \mathcal{T}(x) - y) = \mathcal{K}(x, \mathcal{T}'(x) - y', t - \tau)$ is non-singular one for any $x, y \in \mathbb{D}_+^{n+1}$. Taking $\tilde{x} \in \mathbb{D}_-^{n+1}$, there exist positive constants κ_1 and κ_2 such that

$$\kappa_1 \rho(\tilde{x} - y) \leq \rho(\mathcal{T}(x) - y) \leq \kappa_2 \rho(\tilde{x} - y). \quad (5.6)$$

For any $f \in M_{p,\varphi}(\mathbb{D}_+^{n+1})$ and $a \in \text{BMO}(\mathbb{D}_+^{n+1})$ define the non-singular integral operators

$$\left. \begin{aligned} \tilde{\mathfrak{R}}f(x) &= \int_{\mathbb{D}_+^{n+1}} \mathcal{K}(x, \mathcal{T}(x) - y) f(y) \, dy, \\ \tilde{\mathfrak{C}}[a, f](x) &= \int_{\mathbb{D}_+^{n+1}} \mathcal{K}(x, \mathcal{T}(x) - y) [a(y) - a(x)] f(y) \, dy. \end{aligned} \right\} \quad (5.7)$$

Since $\mathcal{K}(x, \mathcal{T}(x) - y)$ is still homogeneous one and satisfies Definition 5.1 (b), we have that

$$|\mathcal{K}(x, \mathcal{T}(x) - y)| \leq \frac{M}{\rho(\mathcal{T}(x) - y)^{n+2}} \leq \frac{C}{\rho(\tilde{x} - y)^{n+2}}.$$

Hence, the operators (5.7) are sublinear and bounded in $L_p(\mathbb{D}_+^{n+1})$, $p \in (1, \infty)$ (see [3]). The following estimates are simple consequences of the results in § 4.

Theorem 5.5. *Let $a \in \text{BMO}(\mathbb{D}_+^{n+1})$ and let $f \in M_{p,\varphi}(\mathbb{D}_+^{n+1})$ with (p, φ) as in Theorem 3.7. The operators $\tilde{\mathfrak{R}}f$ and $\tilde{\mathfrak{C}}[a, f]$ are then continuous in $M_{p,\varphi}(\mathbb{D}_+^{n+1})_2$ and*

$$\|\tilde{\mathfrak{R}}f\|_{p,\varphi;\mathbb{D}_+^{n+1}} \leq C \|f\|_{p,\varphi;\mathbb{D}_+^{n+1}}, \quad \|\tilde{\mathfrak{C}}[a, f]\|_{p,\varphi;\mathbb{D}_+^{n+1}} \leq C \|a\|_* \|f\|_{p,\varphi;\mathbb{D}_+^{n+1}}, \quad (5.8)$$

with a constant independent of a and f .

Corollary 5.6. *Let $a \in \text{VMO}$ and let (p, φ) be as above. For any $\varepsilon > 0$ there then exists a positive number $r_0 = r_0(\varepsilon, \eta_a)$ such that, for any $\mathcal{E}_r^+(x^0)$ with a radius $r \in (0, r_0)$ and all $f \in M_{p,\varphi}(\mathcal{E}_r^+(x^0))$,*

$$\|\mathfrak{C}[a, f]\|_{p,\varphi;\mathcal{E}_r^+(x^0)} \leq C\varepsilon \|f\|_{p,\varphi;\mathcal{E}_r^+(x^0)}, \quad (5.9)$$

where C is independent of ε, f, r and x^0 .

6. Proof of the main result

Consider (2.1) with $f \in M_{p,\varphi}(Q)$, (p, φ) as in Theorem 3.7. Since $M_{p,\varphi}(Q)$ is a proper subset of $L_p(Q)$, (2.1) is uniquely solvable and the solution u belongs at least to $W_p^{2,1}(Q)$. Our aim is to show that this solution also belongs to $W_{p,\varphi}^{2,1}(Q)$. For this we need an *a priori* estimate of u , which we prove in two steps.

Can this word be removed? Otherwise, please clarify the sentence.

Just 'still homogeneous'?

Change OK?

Word added - OK?

Interior estimate. For any $x_0 \in \mathbb{R}_+^{n+1}$ define the parabolic semi-cylinders $\mathcal{C}_r(x_0) = \mathcal{B}_r(x'_0) \times (t_0 - r^2, t_0)$. Let $v \in C_0^\infty(\mathcal{C}_r)$ and suppose that $v(x, t) = 0$ for $t \leq 0$. According to [3, Theorem 1.4], for any $x \in \text{supp} v$ the following representation formula for the second derivatives of v holds true:

$$\begin{aligned} D_{ij}v(x) &= \text{PV} \int_{\mathbb{R}^{n+1}} \Gamma_{ij}(x, x-y)[a^{hk}(y) - a^{hk}(x)]D_{hk}v(y) \, dy \\ &\quad + \text{PV} \int_{\mathbb{R}^{n+1}} \Gamma_{ij}(x, x-y)\mathcal{P}v(y) \, dy + \mathcal{P}v(x) \int_{\mathbb{S}^n} \Gamma_j(x, y)\nu_i \, d\sigma_y, \end{aligned} \quad (6.1)$$

where $\nu(\nu_1, \dots, \nu_{n+1})$ is the outward normal to \mathbb{S}^n . Here, $\Gamma(x, \xi)$ is the fundamental solution of the operator \mathcal{P}_ξ and $\Gamma_{ij}(x, \xi) = \partial^2 \Gamma(x, \xi) / \partial \xi_i \partial \xi_j$. Since any function $v \in W_p^{2,1}$ can be approximated by C_0^∞ functions, the representation formula (6.1) still holds for any $v \in W_p^{2,1}(\mathcal{C}_r(x_0))$. The properties of the fundamental solution (see [3, 15, 24]) imply that Γ_{ij} are variable Calderón–Zygmund kernels in the sense of Definition 5.1. Using the notation of (5.1), we can write

$$\begin{aligned} D_{ij}v(x) &= \mathfrak{C}_{ij}[a^{hk}, D_{hk}v](x) \\ &\quad + \mathfrak{K}_{ij}(\mathcal{P}v)(x) + \mathcal{P}v(x) \int_{\mathbb{S}^n} \Gamma_j(x, y)\nu_i \, d\sigma_y, \quad i, j = 1, \dots, n. \end{aligned} \quad (6.2)$$

The operators \mathfrak{K}_{ij} and \mathfrak{C}_{ij} are defined by (5.1) with $\mathcal{K}(x, x-y) = \Gamma_{ij}(x, x-y)$. Due to Corollaries 5.3 and 5.4 and the equivalence of the metrics, we get that

$$\|D^2v\|_{p, \varphi; \mathcal{C}_r(x_0)} \leq C(\varepsilon \|D^2v\|_{p, \varphi; \mathcal{C}_r(x_0)} + \|\mathcal{P}u\|_{p, \varphi; \mathcal{C}_r(x_0)}) \quad (6.3)$$

for some r small enough. Moving the norm of D^2v on the left-hand side, we get that

$$\|D^2v\|_{p, \varphi; \mathcal{C}_r(x_0)} \leq C(n, p, \eta_{\mathbf{a}}, \|D\Gamma\|_{\infty, Q}) \|\mathcal{P}v\|_{p, \varphi; \mathcal{C}_r(x_0)}.$$

Define a cut-off function $\phi(x) = \phi_1(x')\phi_2(t)$, with $\phi_1 \in C_0^\infty(\mathcal{B}_r(x'_0))$, $\phi_2 \in C_0^\infty(\mathbb{R})$ such that

$$\phi_1(x') = \begin{cases} 1, & x' \in \mathcal{B}_{\theta r}(x'_0), \\ 0, & x' \notin \mathcal{B}_{\theta' r}(x'_0), \end{cases} \quad \phi_2(t) = \begin{cases} 1, & t \in (t_0 - (\theta r)^2, t_0], \\ 0, & t < t_0 - (\theta' r)^2, \end{cases}$$

with $\theta \in (0, 1)$, $\theta' = \theta(3 - \theta)/2 > \theta$ and $|D^s \phi| \leq C[\theta(1 - \theta)r]^{-s}$, $s = 0, 1, 2$, $|\phi_t| \sim |D^2 \phi|$. For any solution $u \in W_p^{2,1}(Q)$ of (2.1) define $v(x) = \phi(x)u(x) \in W_p^{2,1}(\mathcal{C}_r)$. Hence,

$$\begin{aligned} \|D^2u\|_{p, \varphi; \mathcal{C}_{\theta r}(x_0)} &\leq \|D^2v\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)} \\ &\leq C \|\mathcal{P}v\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)} \\ &\leq C \left(\|f\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)} + \frac{\|Du\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)}}{\theta(1 - \theta)r} + \frac{\|u\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)}}{[\theta(1 - \theta)r]^2} \right). \end{aligned}$$

Hence,

$$\begin{aligned} &[\theta(1 - \theta)r]^2 \|D^2u\|_{p, \varphi; \mathcal{C}_{\theta r}(x_0)} \\ &\leq ([\theta(1 - \theta)r]^2 \|f\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)} + \theta(1 - \theta)r \|Du\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)} + \|u\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)}) \\ &\quad \text{(by the choice of } \theta' \text{ it follows that } \theta(1 - \theta) \leq 2\theta'(1 - \theta')) \\ &\leq C(r^2 \|f\|_{p, \varphi; Q} + \theta'(1 - \theta')r \|Du\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)} + \|u\|_{p, \varphi; \mathcal{C}_{\theta' r}(x_0)}). \end{aligned}$$

Introducing the semi-norms

$$\Theta_s = \sup_{0 < \theta < 1} [\theta(1 - \theta)r]^s \|D^s u\|_{p, \varphi; \mathcal{C}_{\theta r}(x_0)}, \quad s = 0, 1, 2,$$

the above inequality becomes

$$[\theta(1 - \theta)r]^2 \|D^2 u\|_{p, \varphi; \mathcal{C}_{\theta r}(x_0)} \leq \Theta_2 \leq C(r^2 \|f\|_{p, \varphi; Q} + \Theta_1 + \Theta_0). \quad (6.4)$$

The interpolation inequality [25, Lemma 4.2] gives that there exists a positive constant C independent of r such that

$$\Theta_1 \leq \varepsilon \Theta_2 + \frac{C}{\varepsilon} \Theta_0 \quad \text{for any } \varepsilon \in (0, 2).$$

Thus, (6.4) becomes

$$[\theta(1 - \theta)r]^2 \|D^2 u\|_{p, \varphi; \mathcal{C}_{\theta r}(x_0)} \leq \Theta_2 \leq C(r^2 \|f\|_{p, \varphi; Q} + \Theta_0) \quad \forall \theta \in (0, 1).$$

Taking $\theta = \frac{1}{2}$ we get the Caccioppoli-type estimate

$$\|D^2 u\|_{p, \varphi; \mathcal{C}_{r/2}(x_0)} \leq C \left(\|f\|_{p, \varphi; Q} + \frac{1}{r^2} \|u\|_{p, \varphi; \mathcal{C}_r(x_0)} \right).$$

To estimate u_t we exploit the parabolic structure of the equation and the boundedness of the coefficients

$$\begin{aligned} \|u_t\|_{p, \varphi; \mathcal{C}_{r/2}(x_0)} &\leq \|\mathbf{a}\|_{\infty; Q} \|D^2 u\|_{p, \varphi; \mathcal{C}_{r/2}(x_0)} + \|f\|_{p, \varphi; \mathcal{C}_{r/2}(x_0)} \\ &\leq C \left(\|f\|_{p, \varphi; Q} + \frac{1}{r^2} \|u\|_{p, \varphi; \mathcal{C}_r(x_0)} \right). \end{aligned}$$

Consider the cylinders $Q' = \Omega' \times (0, T)$ and $Q'' = \Omega'' \times (0, T)$ with $\Omega' \Subset \Omega'' \Subset \Omega_{\underline{t}}$ by the standard covering procedure and partition of the unity we get that Change OK?

$$\|u\|_{W_{p, \varphi}^{2,1}(Q')} \leq C(\|f\|_{p, \varphi; Q} + \|u\|_{p, \varphi; Q''}), \quad (6.5)$$

where C depends on $n, p, \Lambda, T, \|D\Gamma\|_{\infty; Q}, \eta_{\mathbf{a}}, \|\mathbf{a}\|_{\infty, Q}$ and $\text{dist}(\Omega', \partial\Omega'')$.

Boundary estimates. For any fixed $(x^0, r) \in \mathbb{R}^{n-1} \times \mathbb{R}_+$ define the semi-cylinders

$$\mathcal{C}_r^+(x^0) = \mathcal{B}_r^+(x^{0'}) \times (0, r^2) = \{|x^0 - x'| < r, x_n > 0, 0 < t < r^2\}$$

with $\mathcal{S}_r^+ = \{(x'', 0, t) : |x^0 - x''| < r, 0 < t < r^2\}$. For any solution $u \in W_p^{2,1}(\mathcal{C}_r^+(x^0))$ with $\text{supp } u \in \mathcal{C}_r^+(x^0)$, the following boundary representation formula holds (see [3]):

$$D_{ij}u(x) = \mathfrak{C}_{ij}[a^{hk}, D_{hk}u](x) + \mathfrak{K}_{ij}(\mathcal{P}u)(x) + \mathcal{P}u(x) \int_{\mathbb{S}^n} \Gamma_j(x, y) \nu_i d\sigma_y - \mathfrak{J}_{ij}(x),$$

where

$$\begin{aligned}\mathfrak{J}_{ij}(x) &= \tilde{\mathfrak{K}}_{ij}(\mathcal{P}u)(x) + \tilde{\mathfrak{C}}_{ij}[a^{hk}, D_{hk}u](x), & i, j = 1, \dots, n-1, \\ \mathfrak{J}_{in}(x) &= \mathfrak{J}_{ni}(x) = \sum_{l=1}^n \left(\frac{\partial \mathcal{T}(x)}{\partial x_n} \right)^l [\tilde{\mathfrak{C}}_{il}[a^{hk}, D_{hk}u](x) + \tilde{\mathfrak{K}}_{il}(\mathcal{P}u)(x)], & i = 1, \dots, n-1, \\ \mathfrak{J}_{nm}(x) &= \sum_{r,l=1}^n \left(\frac{\partial \mathcal{T}(x)}{\partial x_n} \right)^r \left(\frac{\partial \mathcal{T}(x)}{\partial x_n} \right)^l [\tilde{\mathfrak{C}}_{rl}[a^{hk}, D_{hk}u](x) + \tilde{\mathfrak{K}}_{rl}(\mathcal{P}u)(x)], \\ \frac{\partial \mathcal{T}(x)}{\partial x_n} &= \left(-2 \frac{a^{n1}(x)}{a^{nn}(x)}, \dots, -2 \frac{a^{nn-1}(x)}{a^{nn}(x)}, -1, 0 \right).\end{aligned}$$

Here, $\tilde{\mathfrak{K}}_{ij}$ and $\tilde{\mathfrak{C}}_{ij}$ are non-singular operators defined by (5.7) with $\mathcal{K}(x, \mathcal{T}(x) - y) = \Gamma_{ij}(x, \mathcal{T}(x) - y)$. Applying the estimates (5.8) and (5.9) and having in mind that the components of the vector $\partial \mathcal{T}(x)/\partial x_n$ are bounded, we get that

$$\|D^2 u\|_{p, \varphi; \mathcal{C}_r^+(x^0)} \leq C(\|\mathcal{P}u\|_{p, \varphi; \mathcal{C}_r^+(x^0)} + \|u\|_{p, \varphi; \mathcal{C}_r^+(x^0)}).$$

The Jensen inequality applied to $u(x) = \int_0^t u_s(x', s) ds$ and the parabolic structure of the equation give that

$$\|u\|_{p, \varphi; \mathcal{C}_r^+(x^0)} \leq Cr^2 \|u_t\|_{p, \varphi; \mathcal{C}_r^+(x^0)} \leq C(\|f\|_{p, \varphi; Q} + r^2 \|u\|_{p, \varphi; \mathcal{C}_r^+(x^0)}).$$

Taking r small enough we can move the norm of u on the left-hand side, obtaining that

$$\|u\|_{p, \varphi; \mathcal{C}_r^+} \leq C\|f\|_{p, \varphi; Q},$$

with a constant C depending on $n, p, A, T, \eta_{\mathbf{a}}, \|\mathbf{a}\|_{\infty, Q}$. By covering the boundary with small cylinders, partition of the unit subordinated of that covering and local flattering of $\partial \Omega$ we get that

$$\|u\|_{W_{p, \varphi}^{2,1}(Q \setminus Q')} \leq C\|f\|_{p, \varphi; Q}. \quad (6.6)$$

Unifying (6.5) and (6.6), we get (2.4).

Acknowledgements. The research of V.G. was partly supported by the Science Development Foundation under the President of the Republic of Azerbaijan (Grant EIF-2010-1(1)-40/06-1), by the 2011 Ahi Evran University Scientific Research Projects (Grant BAP FBA-11-13) and by the Scientific and Technological Research Council of Turkey (TUBITAK Project 110T695).

The authors are indebted to the referee for their valuable remarks.

References

1. P. ACQUISTAPACE, On *BMO* regularity for linear elliptic systems, *Annali Mat. Pura Appl.* **161** (1992), 231–270.
2. A. AKBULUT, V. S. GULIYEV AND R. MUSTAFAYEV, On the boundedness of the maximal operator and singular integral operators in generalized Morrey spaces, *Math. Bohem.* **137**(1) (2012), 27–43.

Change OK?
'to' ?
'flattening' ?

3. M. BRAMANTI AND M. C. CERUTTI, $W_p^{1,2}$ solvability for the Cauchy–Dirichlet problem for parabolic equations with VMO coefficients, *Commun. PDEs* **18** (1993), 1735–1763.
4. M. CARRO, L. PICK, J. SORIA AND V. D. STEPANOV, On embeddings between classical Lorentz spaces, *Math. Inequal. Appl.* **4**(3) (2001), 397–428.
5. F. CHIARENZA AND M. FRASCA, Morrey spaces and Hardy–Littlewood maximal function, *Rend. Mat.* **7** (1987), 273–279.
6. F. CHIARENZA, M. FRASCA AND M. P. LONGO, Interior $W^{2,p}$ -estimates for nondivergence elliptic equations with discontinuous coefficients, *Ric. Mat.* **40** (1991), 149–168.
7. E. B. FABES AND N. RIVIÈRE, Singular integrals with mixed homogeneity, *Studia Math.* **27** (1996), 19–38.
8. V. S. GULIYEV, Integral operators on function spaces on the homogeneous groups and on domains in \mathbb{R}^n , PhD Thesis, Steklov Institute of Mathematics (1994; in Russian).
9. V. S. GULIYEV, Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces, *J. Inequal. Applicat.* **2009** (2009), 503948.
10. V. S. GULIYEV AND F. M. MUSHTAGOV, Parabolic equations with VMO coefficients in weighted Lebesgue spaces, *Proc. Razmadze Math. Inst.* **137** (2005), 1–27.
11. V. S. GULIYEV AND L. G. SOFTOVA, Global regularity in generalized Morrey spaces of solutions to non-divergence elliptic equations with VMO coefficients, *Potent. Analysis* **38**(3) (2013), 843–862.
12. V. S. GULIYEV, S. S. ALIYEV, T. KARAMAN AND P. SHUKUROV, Boundedness of sublinear operators and commutators on generalized Morrey spaces, *Integ. Eqns Operat. Theory* **71**(3) (2011), 327–355.
13. F. JOHN AND L. NIRENBERG, On functions of bounded mean oscillation, *Commun. Pure Appl. Math.* **14** (1961), 415–426.
14. P. W. JONES, Extension theorems for BMO, *Indiana Univ. Math. J.* **29** (1980), 41–66.
15. O. A. LADYZHENSKAYA, V. A. SOLONNIKOV AND N. N. URAL'TSEVA, *Linear and quasi-linear equations of parabolic type*, Translations of Mathematical Monographs, Volume 23 (American Mathematical Society, Providence, RI, 1968).
16. T. MIZUHARA, Boundedness of some classical operators on generalized Morrey spaces, in *ICM-90 Satellite Conference Proceedings: Harmonic Analysis*, pp. 183–189 (Springer, 1991).
17. C. B. MORREY, On the solutions of quasi-linear elliptic partial differential equations, *Trans. Am. Math. Soc.* **43** (1938), 126–166.
18. E. NAKAI, Hardy–Littlewood maximal operator, singular integral operators and the Riesz potentials on generalized Morrey spaces, *Math. Nachr.* **166** (1994), 95–103.
19. E. NAKAI, The Campanato, Morrey and Hölder spaces on spaces of homogeneous type, *Studia Math.* **176**(1) (2006), 1–19.
20. D. K. PALAGACHEV AND L. G. SOFTOVA, Singular integral operators, Morrey spaces and fine regularity of solutions to PDEs, *Potent. Analysis* **20** (2004), 237–263.
21. D. K. PALAGACHEV, M. A. RAGUSA AND L. G. SOFTOVA, Cauchy–Dirichlet problem in Morrey spaces for parabolic equations with discontinuous coefficients, *Boll. UMI B* **8**(6) (2003), 667–683.
22. D. SARASON, On functions of vanishing mean oscillation, *Trans. Am. Math. Soc.* **207** (1975), 391–405.
23. L. G. SOFTOVA, Singular integrals and commutators in generalized Morrey spaces, *Acta Math. Sinica* **22** (2006), 757–766.
24. L. G. SOFTOVA, Singular integral operators in Morrey spaces and interior regularity of solutions to systems of linear PDE's, *J. Global Optim.* **40** (2008), 427–442.
25. L. G. SOFTOVA, Morrey-type regularity of solutions to parabolic problems with discontinuous data, *Manuscr. Math.* **136**(3) (2011), 365–382.

Definitely sloping
in original
publication? See
also two more
instances below.

Apostrophe
definitely in
original?

26. L. G. SOFTOVA, The Dirichlet problem for elliptic equations with *VMO* coefficients in generalized Morrey spaces, in *Advances in harmonic analysis and operator theory, the Stefan Samko anniversary volume*, Operator Theory: Advances and Applications, Volume 229, pp. 371–386 (Springer, 2013).
27. L. G. SOFTOVA, Parabolic oblique derivative problem with discontinuous coefficients in generalized Morrey spaces, *Ric. Mat.* **62**(2) (2013), 265–278.
28. E. STEIN, *Harmonic analysis: real-variable methods, orthogonality and oscillatory integrals*, Princeton Mathematical Series, Volume 43 (Princeton University Press, 1993).