

# Effects of Ellagic acid and Hesperetin on Levels of Some Elements in Livers of Aluminum-Induced Rats

[Aluminyuma Maruz Kalmış Sıçanların Karaciğerlerindeki Bazı Elementlerin Seviyeleri Üzerine Hesperetin ve Ellagik Asitin Etkileri]

Ahmet Özkaya<sup>1</sup>,  
Harun Çiftçi<sup>2</sup>,  
Alpaslan Dayangaç<sup>3</sup>,  
B.Sitki Çevrimli<sup>4</sup>,  
Ali Ölçücü<sup>5</sup>,  
Sait Çelik<sup>6</sup>

<sup>1</sup>Department of Chemistry, Faculty of Science and Art University of Adiyaman, Adiyaman;

Departments of <sup>2</sup>Chemistry and <sup>3</sup>Biology, Faculty of Science and Art University of Ahi Evran, Kirsehir;

<sup>4</sup>University of Gazi, Atatürk Vocational High School, Department of Chemical Technology, Ankara; <sup>5</sup>Department of Chemistry, Faculty of Science University of Firat, Elazığ;

<sup>6</sup>Department of Chemistry, Faculty of Science and Art University of Uşak, Uşak, Turkey

**Yazışma Adresi**  
[Correspondence Address]

**Dr. Ahmet Ozkaya**

Departments of Chemistry, Faculty of Science and Arts, University of Adiyaman, 02040, Adiyaman TURKEY,  
Phone: +90-416-2232210  
E-mail: aozkaya01@gmail.com ,

Registered: 25 April 2012; Accepted: 30 October 2012  
[Kayıt tarihi: 25 Nisan 2012; Kabul Tarihi: 30 Ekim 2012]

## ABSTRACT

**Objective:** Ellagic acid and hesperetin are well known as having free radical scavenging activity in some rat tissues. Aluminum is known to cause toxic effects on various organ systems. In this study, the protective role of ellagic acid and hesperetin has been investigated on macro and trace elements in liver of rats treated with AlCl<sub>3</sub>.

**Method:** Thirty-six healthy adult male *Sprague-dawley* rats (240±10 g body weight) were used in this study. The rats were randomly divided into six groups and Ellagic acid, Hesperetin, AlCl<sub>3</sub>, AlCl<sub>3</sub>+ Ellagic acid and AlCl<sub>3</sub>+Hesperetin were treated to each group except from the control group. Determinations of iron, calcium, copper and zinc were performed using a Flame Atomic Absorption Spectrometer and determination of aluminum and manganese were carried out with Inductively Coupled Plasma Atomic Emission Spectrometer.

**Results:** Levels of aluminum, calcium and iron in AlCl<sub>3</sub> treated group were found to be increased compared to the control group (p<0.01, p<0.001). In AlCl<sub>3</sub>+ Ellagic acid and AlCl<sub>3</sub>+Hesperetin treated groups, the levels of aluminum and calcium were found as decreased (p<0.05, p<0.001). When compared to the control group, level of manganese in liver were found to be increased in ellagic acid treated group (p<0.01). The zinc level of Hesperetin group increased when compared to the other groups (p<0.05).

**Conclusion:** Our results suggest that ellagic acid and hesperetin may have a protective role upon major and trace elements against AlCl<sub>3</sub>-induced stress conditions.

**Key Words:** Aluminum, Ellagic acid, Hesperetin, Trace elements, Calcium

**Conflict of Interest:** The authors declare no conflict of interest.

## ÖZET

**Amaç:** Ellagik asit ve hesperetin bazı sıçan dokularında serbest radikal seviyelerini azaltıcı aktiviteye sahip oldukları iyi bilinmektedir. Alüminyumun çeşitli organ sistemleri üzerine toksik etkisinin olduğu bilinmektedir. Bu çalışmada AlCl<sub>3</sub> uygulanmış sıçanların karaciğerlerinde makro ve eser elementlerin seviyeleri üzerine ellagik asit ve hesperetin koruyucu etkileri araştırılmıştır.

**Yöntem:** Bu çalışmada 36 adet sağlıklı erkek *Sprague-dawley* sıçanı (240±10 g ağırlığında) kullanılmıştır. Sıçanlar rastgele seçilerek 6 grup oluşturulmuş, bir grup kontrol grubu olarak kalmış, diğer grupların her birine Ellagik asit, Hesperetin, AlCl<sub>3</sub>, AlCl<sub>3</sub>+ Ellagik asit ve AlCl<sub>3</sub>+Hesperetin uygulanmıştır. Demir, kalsiyum, bakır ve çinko miktarları Alevli Atomik Absorbsiyon Spektrofotometresi ile alüminyum ve mangan miktarları ise indüklenmiş eşleşmiş plazma atomik emisyon spektrofotometresi ile tayin edilmiştir.

**Bulgular:** AlCl<sub>3</sub> uygulanan grupta alüminyum, kalsiyum ve demir düzeylerinin kontrol grubuna göre arttığı gözlenmiştir (p<0.01, p<0.001). AlCl<sub>3</sub>+ Ellagik asit ve AlCl<sub>3</sub>+Hesperetin gruplarında alüminyum ve kalsiyum seviyesi düşük bulunmuştur (p<0.05). Kontrol grubu ile kıyaslandığında ellagik asit karaciğerde mangan seviyesini artırdı (p<0.01). Diğer gruplarla kıyaslandığında Hesperetin grubunun çinko seviyesi artmıştır (p<0.05).

**Sonuçlar:** Sonuçlarımıza göre, ellagik asit ve hesperetin, AlCl<sub>3</sub> kaynaklı stress koşullarına karşı makro ve eser elementler üzerinde koruyucu bir etkisi olabileceğini düşündürmektedir.

**Anahtar Kelimeler:** Alüminyum, Ellagik asit, Hesperetin, Eser Elementler, Kalsiyum

**Çıkar Çatışması:** Yazarların mevcut olan konuyla ilgili herhangi bir çıkar çatışması bulunmamaktadır.

## Introduction

Elements are required for the proper functioning of the human system. Deficiency of those elements causes serious metabolic abnormalities and the increase of those creates toxicity. The levels of trace elements in some diseases, such as chronic kidney, liver and lung diseases, have been studied and important results have been obtained, as well. Aluminum is known to cause toxic effects in a variety of organ systems, including brain, bone, kidney and blood. It has been suggested that there has been a relationship between high level of aluminum and increased risk of a number of pathogenic disorders, such as microcytic anemia, neurodegenerative disorders including dialysis encephalopathy, Alzheimer's disease and Parkinson's disease. Aluminum ions have strong affinity to bio-membranes; they are capable of increasing the cellular oxidative milieu by potentiating the pro-oxidant properties of transition metals [1-3].

Hesperetin (HES) and Ellagic acid (EA) are polyphenolic compounds found in a wide variety of fruits. It has been reported that HES (3, 5, 7-trihydroxy-4-methoxy flavanone) shows a wide spectrum of pharmacological effects such as, anti-carcinogenic, anti-atherogenic and anti-hypertensive effects [4, 5].

EA(2,3,7,8-tetrahydroxy[1]benzopyrano[5,4,3-cde][1]benzopyran-5,10-dione) is known to increase antioxidant activity in lipid peroxidation and protect cells from oxidative damage. In addition, EA exhibits important health promoting effects via its antioxidant, anti-proliferative, chemo-preventive, anti-atherogenic, anti-apoptotic, anti-inflammatory activities and it also repairs DNA damage [6-9]. In our previous study, we observed positive effects of EA on some biochemical parameters in oxidative stress induced rats [10]. It has been stated that deficiency and redundancy of the elements is important for the metabolism [1-3]. We have determined in our literature review that effects of EA and HES upon trace element and major elements in aluminum induced rat tissues has not been researched before.

The aim of this study was to analyze effects of EA and HES on trace element and major element levels in  $AlCl_3$  treated rat liver.

## Materials and Methods

### Chemicals

$AlCl_3 \cdot 6H_2O$ , EA, HES and other chemicals were purchased from Merck Chemical Co (Germany).

### Animals and Treatment

Thirty-six healthy adult male *Sprague-dawley* rats (240±10 g body weight) were used in this study. The animals were obtained from Firat University Experimental Research Centre (FUDAM), Elazığ-Turkey. They have been kept at 21±1°C with a 12-h light/

dark cycle and have been given a commercial pellet diet (Elazığ Food Company, Elazığ-Turkey) and fresh drinking water *ad libitum*. Animal use protocol was approved by the National Institute of Health and Local Committee on Animal Research (Number:27.07.2006-3). The rats were randomly divided into six groups with each group containing six rats.

First group: Control (C)

Second group: EA alone administered group

Third group: HES alone administered group

Fourth group:  $AlCl_3$  (A) administered group

Fifth group: A+EA administered group

Sixth group: A+HES administered group

EA and HES were dissolved in corn oil and administered to animals by gavage at the dose of 12 mg/kg EA [10] and 75 mg/kg HES [11] body weight.  $AlCl_3$  was dissolved in distilled water and administered to the animals through intraperitoneal route at a dose of 8.3 mg/kg for 55 days [10]. The last dose was administered 12 h before the operation.

### Analysis of Tissue Samples

The rats were decapitated and their livers were removed by surgery. Liver samples (0.50-0.80 g) were washed with isotonic salt solution (0.9 % v/w) and homogenized. A 0.30 g homogenized sample was digested by microwave digestion system with closed system pots polytetrafluorethylen (PTFE) according to our previous study [12,13]. The same procedure was applied to the blank solution. Determinations of iron (Fe), calcium (Ca), copper (Cu) and zinc(Zn) were performed using a Flame Atomic Absorption Spectrometer (FAAS) (Perkin Elmer, Analysis 400) and determination of aluminum (Al) and manganese (Mn) were carried out with Inductively Coupled Plasma Atomic Emission Spectrometer (ICP-AES) (Perkin Elmer 3100).

### Statistical Analysis

The experimental results were reported as mean ± S.E. Statistical analysis was performed using SPSS 10.0 Software. Analysis of variance (ANOVA) and an LSD (Least Significant Difference) test were used to compare the experimental groups with the controls

## Results

The levels of elements in rat livers are shown in Table 1 and Table 2.

$AlCl_3$  has been increased in Fe, Ca and Al levels in liver ( $p < 0.01$ ,  $p < 0.001$ ) when compared to the control group. No statistically significant difference was observed in Fe, Ca and Al levels among EA, HES and control groups ( $p > 0.05$ ). Administration of ellagic acid and hesperetin to  $AlCl_3$ -treated rats has prevented the  $AlCl_3$ -induced increase in Al ( $p < 0.001$ ) and Ca ( $p < 0.05$ ) levels in liver tissues. Ellagic acid treatment has caused significant

**Table 1.** Concentrations ( $\mu\text{g/g}$ ) of Macro and Trace Elements in Liver of Rats Treated with  $\text{AlCl}_3$  and/or EA

Groups	Fe	Ca	Zn	Cu	Mn	Al
C	117,73 $\pm$ 3,15 <sup>a</sup>	91,53 $\pm$ 3,66 <sup>a</sup>	31,51 $\pm$ 1,21 <sup>a</sup>	2,22 $\pm$ 0,26 <sup>a</sup>	1,21 $\pm$ 0,08 <sup>a</sup>	4,73 $\pm$ 0,46 <sup>a</sup>
EA	131,65 $\pm$ 4,42 <sup>a</sup>	86,74 $\pm$ 8,92 <sup>at</sup>	32,13 $\pm$ 1,18 <sup>a</sup>	2,49 $\pm$ 0,31 <sup>a</sup>	1,56 $\pm$ 0,11 <sup>oz</sup>	7,78 $\pm$ 1,26 <sup>at</sup>
A	144,94 $\pm$ 5,70 <sup>c</sup>	135,26 $\pm$ 11,70 <sup>d</sup>	32,50 $\pm$ 1,80 <sup>a</sup>	1,87 $\pm$ 0,53 <sup>a</sup>	1,14 $\pm$ 0,08 <sup>a</sup>	41,63 $\pm$ 5,35 <sup>d</sup>
A+EA	139,94 $\pm$ 6,29 <sup>b</sup>	95,24 $\pm$ 8,22 <sup>az</sup>	29,66 $\pm$ 1,71 <sup>a</sup>	2,65 $\pm$ 0,30 <sup>ay</sup>	1,15 $\pm$ 0,05 <sup>a</sup>	25,21 $\pm$ 5,11 <sup>dt</sup>

Values are expressed as means  $\pm$  SD; n=6 for each treatment group. a:  $p > 0.05$   
 Statistical significance compared to the control group: b:  $p < 0.05$ , c:  $p < 0.01$ , d:  $p < 0.001$   
 Statistical significance compared to the  $\text{AlCl}_3$  group: y:  $p < 0.05$ , z:  $p < 0.01$ , t:  $p < 0.001$

**Table 2.** Concentrations ( $\mu\text{g/g}$ ) of Macro and Trace Elements in Liver of Rats Treated with  $\text{AlCl}_3$  and/or HES

Groups	Fe	Ca	Zn	Cu	Mn	Al
C	117,73 $\pm$ 3,15 <sup>a</sup>	91,53 $\pm$ 3,66 <sup>a</sup>	31,51 $\pm$ 1,21 <sup>a</sup>	2,22 $\pm$ 0,26 <sup>a</sup>	1,21 $\pm$ 0,08 <sup>a</sup>	4,73 $\pm$ 0,46 <sup>a</sup>
HES	125,28 $\pm$ 3,43 <sup>ay</sup>	103,42 $\pm$ 3,30 <sup>az</sup>	37,47 $\pm$ 1,20 <sup>by</sup>	2,68 $\pm$ 0,07 <sup>ay</sup>	1,23 $\pm$ 0,07 <sup>a</sup>	7,15 $\pm$ 0,44 <sup>at</sup>
A	144,94 $\pm$ 5,70 <sup>c</sup>	135,26 $\pm$ 11,70 <sup>d</sup>	32,50 $\pm$ 1,80 <sup>a</sup>	1,87 $\pm$ 0,53 <sup>a</sup>	1,14 $\pm$ 0,08 <sup>a</sup>	41,63 $\pm$ 5,35 <sup>d</sup>
A+HES	136,28 $\pm$ 8,61 <sup>b</sup>	110,71 $\pm$ 4,65 <sup>ay</sup>	35,96 $\pm$ 2,18 <sup>a</sup>	1,54 $\pm$ 0,06 <sup>b</sup>	1,05 $\pm$ 0,04 <sup>a</sup>	22,28 $\pm$ 3,10 <sup>dt</sup>

Values are expressed as means  $\pm$  SD; n=6 for each treatment group. a:  $p > 0.05$   
 Statistical significance compared to the control group: b:  $p < 0.05$ , c:  $p < 0.01$ , d:  $p < 0.001$   
 Statistical significance compared to the  $\text{AlCl}_3$  group: y:  $p < 0.05$ , z:  $p < 0.01$ , t:  $p < 0.001$

increase in Mn ( $p < 0.01$ ) levels of liver tissues and Zn levels of HES group increased when compared to control group ( $p < 0.05$ ).

## Discussion

A number of studies have shown that animals exposed to  $\text{AlCl}_3$  have increased Al concentrations in liver, plasma, urinary, kidney [14], testis and serum [15], hippocampus [16], and brain [17,18]. In the present study, there has been a significant increase in liver Al level after  $\text{AlCl}_3$  exposure. Yousef *et al.* reported that  $\text{AlCl}_3$  which is capable of causing marked alterations in some biochemical parameters has induced oxidative damage and inhibited the activities of antioxidant enzymes [19]. Wilhelm *et al.* reported that Al accumulation in the liver can be caused by Al exposure and at high concentrations this metal can have toxic effect upon the tissue [20]. In our results, Al levels increased as response to  $\text{AlCl}_3$  treatment, implying the increased oxidative damage to the tissues. It was observed that Fe level in liver of  $\text{AlCl}_3$  administered group was found to be increased compared to the control group. Some studies have shown that animals that are exposed to Al have had increased Fe concentrations in brain, urinary excretion [17] and testis [15]. Ward RJ *et al.* noted that in an animal model of Al

overload, increases in tissue iron in kidney, liver, heart and spleen, and also in various brain regions, such as frontal, temporal and parietal cortex and hippocampus, has been parallel to the increase at tissue Al content [21]. The ionic radii of  $\text{Al}^{3+}$  most closely resemble to those of  $\text{Fe}^{3+}$ , therefore the appearance of  $\text{Al}^{3+}$  in  $\text{Fe}^{3+}$  sites is possible. Al is known to be bound by  $\text{Fe}^{3+}$  carrying protein transferrin, thus reducing the binding of  $\text{Fe}^{2+}$ . The increase in free intracellular  $\text{Fe}^{2+}$  causes peroxidation of membrane lipids and thus causes membrane damage [22]. We observed that administration of EA and HES of  $\text{AlCl}_3$  treated animals decreased the Al levels in liver tissues. In addition, EA and HES combination caused a relative reduction in Fe levels. EA and HES are flavonoid-derived substances. Fernandez *et al.* reported that flavonoids could be very effective antioxidants which could also operate through antiradical and chelating mechanisms [23]. We have considered that the reason why EA and HES have decreased the level of Aluminum can be metal chelation and antioxidant effects of those flavonoid-derived substances. Moreover, in our previous study, EA decreased the high MDA levels in aluminum treated rat livers and EA also regulated GSH-PX enzyme activity which was decreased by the effect of aluminum [10].

Intracellular calcium ions have been assumed to increase its importance as a regulator of a variety of cellular processes, including muscle contraction, secretion and cell division [24, 25]. We found that Ca concentrations of the A group increased in the liver tissues, suggesting that Ca levels were significantly affected by the Al toxicity. Phenolic phytochemicals such as EA and flavanoids from fruits have been known as good chelators; therefore, these phytochemicals can effectively chelate the ions such as Ca in the extra cellular matrix or the cytosol and alter the net concentration of free Ca [26-29]. We observed that the increased liver Ca concentration due to Al exposure was reduced by EA and HES treatments. We think that EA and HES may have a corrective effect upon Ca concentration. Our results are in agreement with the literature [24-29].

In the current study, we have found an increase in Mn concentrations in liver tissues of EA administered rats. Esparza *et al.* demonstrated that Mn concentrations were higher in the melatonin- treated group than the one in Al-treated group in liver [30]. Moreover, in this study, it has been suggested that the protective effects of melatonin against cellular damage caused by Al-induced oxidative stress, together with its low toxicity, make melatonin worthy of investigation as a potential supplement to be included in the treatment of neurological disorders in which the oxidative effects must be minimized.

Furthermore, Mn acts as a cofactor of the antioxidant enzyme superoxide dismutase. It has been found that Mn deficiency stimulates lipid peroxidation [31]. However, we could not observe similar results in HES group.

Levels of Zn in HES-treated group were significantly higher than that in other groups ( $p < 0.05$ ) and there was not any statistical significant difference amongst of C, A, A+HES and A+EA groups ( $p > 0.05$ ). Zn is necessary for proper liver function and Zn deficiency has been related to the pathogenesis of multiple liver diseases [32-35]. Although liver Zn levels were observed to be increased by HES in HES treated groups, there were no statistically significant changes in EA administered groups. These effects of HES may be considered to be associated with liver enzymes.

It was found that Cu levels were relatively decreased in A group when compared to C group. The Cu levels of A+EA groups were increased in contrast to A group ( $p < 0.05$ ). Therefore, it can be considered that EA have a regulatory impact upon Cu concentrations. Gumus *et al.* found that EA has modulator effects on Cu and Zn levels in liver and serum of cholestatic rats [36]. Level of Cu in A+ HES-treated group was lower than the control group ( $p < 0.01$ ) and there was not any statistically significant difference among C, HES and A groups ( $p > 0.05$ ).

In conclusion, we observed that administration of  $AlCl_3$  to rats, increased liver tissue Al and Ca levels. EA and HES decreased Al and Ca levels in  $AlCl_3$  treated animals. Moreover, whereas EA has increased the levels

of Cu and Mn in EA treated individual group, HES has provided increase at Zn level. At present, aluminum found in food, water and environment may have harmful effects on human metabolism; none the less, we consider that this harmful effect can be minimized consuming EA and HES that exist in fruits and vegetables.

## Acknowledgments

This research was supported by the FUBAP project (No: 1687).

**Ethical Considerations:** The study protocol was approved by the National Institute of Health and Local Committee on Animal Research (Date: 27 July 2006, number: 3). All subjects were informed about the details of the study and the written consent of each subject was received.

**Conflict of Interest:** The authors declare no conflict of interest.

## References

- [1] Berthon G. Aluminum speciation in relation to aluminum bio-availability, metabolism and toxicity. *Coord. Chem. Rev* 2002; 228(2): 319-341.
- [2] Good PF, Perl DP, Bierer LM, Schidler J. Selective accumulation of aluminum and iron in the neurofibrillary tangles of Alzheimer's disease: a laser microprobe (LAMMA) study. *Annals of Neurology* 1992; 31(3): 286-292.
- [3] Yokel RA. Brain uptake, retention, and efflux of aluminum and manganese. *Environ Health Persp* 2002, 110 (5): 699-704.
- [4] Garg A, Garg S, Zaneveld LJD, Singla AK. Chemistry and pharmacology of the citrus bioflavonoid hesperidin. *Phytotherapy Res* 2001; 15(8): 655-669.
- [5] Galati EM, Trovato A, Kirjavainen S, Forestieri AM, Rossitto A, *et al.* Biological effects of hesperidin, a citrus flavonoid. (Note III). Antihypertensive and diuretic activity in rat. *Farmaco* 1996; 51(3):219-221.
- [6] Priyadarsini KI, Khopde SM, Kumar SS, Mohan H. Free radical studies of ellagic acid, a natural phenolic antioxidant. *J Agric Food Chem* 2002; 50(7):2200-2206.
- [7] Festa F, Aglitti T, Duranti G, Ricordy R, Perticone P, *et al.* Strong antioxidant activity of ellagic acid in mammalian cells in vitro revealed by the comet assay. *Anticancer Res* 2001; 21(6A):3903-3908.
- [8] Losso JN, Bansode RR, Trappey A, Bawadi HA, Truax R. In vitro antiproliferative activities of ellagic acid. *J Nutr Biochem* 2004; 15(11):672-678.
- [9] Girdhani S, Bhosle SM, Thulsidas SA, Kumar A, Mishra KP. Potential of radiosensitizing agents in cancer chemo-radiotherapy. *J Cancer Res Ther* 2005; 1(3):129-131.
- [10] Ozkaya A, Celik S, Yuce A, Sahin Z, Yilmaz O. The effects of ellagic acid on some biochemical parameters in the liver of rats against oxidative stress induced by aluminum. *Kafkas Univ Vet Fak Derg* 2010; 16 (2): 263-268.
- [11] Rotelli AE, Guardia T, Juarez AO, De La Rocha NE, Pelzer LE. Comparative study of flavonoids in experimental models of inflammation. *Pharmacological Res* 2003; 48(6): 601-606.
- [12] Ciftci H, Ozkaya A, Dayangac A, Olüçü A, Celik S, Sahin Z, *et al.* Effect of lipoic acid on the some elements in bra-

- in tissue of DMBA-induced guinea pigs. *Kafkas Univ Vet Fak Derg* 2009; 15(4): 569-573.
- [13] Ciftci H, Dayangac A, Ozkaya A, Cevrimli BS, Erdag R, Olcucu A, *et al.* Multi-element determination of macro and trace elements in kidney of DMBA and linalool applied guinea pigs by inductively-coupled plasma atomic emission spectrometry (ICP-AES). *Fresenius Environ Bul* 2011;20 (3):818-822.
- [14] Spencer AJ, Wood JA, Saunders HC, Freeman MS, Lote CJ. Aluminum deposition in liver and kidney following acute intravenous administration of aluminum-chloride or citrate in conscious rats. *Human & Experimental Tox* 1995; 14 (10):787-794.
- [15] Guo CH, Huang CJ, Chiou YL, Hsu GSW. Alteration of trace element distribution and testis ACE activity in lice with high peritoneal aluminum. *Biol Trace Elem Res* 2002; 86 (2): 145-157.
- [16] Gomez M, Esparza JL, Nogues MR, Cabre M, Domingo JL. Pro-oxidant activity of aluminum in the rat hippocampus: Gene expression of antioxidant enzymes after melatonin administration. *Free Radic Biol Med* 2005; 38(1): 104-111.
- [17] Chmielnicka J, Nasiadek M. Tissue distribution and urinary-excretion of essential elements in rats orally exposed to aluminum-chloride. *Biol Trace Elem Res* 1991; 31(2):131-138.
- [18] Yang MS, Wong HF, Yung KI. Determination of endogenous trace metal contents in various mouse brain regions after prolonged oral administration of aluminum chloride. *J of Tox and Environ Health-Part A-Current Issues* 1998; 55 (6):445-453.
- [19] Newairy ASA, Salama AF, Hussien HM, Yousef MI. Propolis alleviates aluminium-induced lipid peroxidation and biochemical parameters in male rats. *Food And Chemical Tox* 2009; 47(6):1093–1098.
- [20] Wilhelm M, Jaeger DE, Schüll-Cablitz H, Hafner D, Idel H. Hepatic clearance and retention of aluminium: Studies in the isolated perfused rat liver. *Toxicol. Lett.* 1996; 89(3): 257–263.
- [21] Ward RJ, Zhang Y, Crichton RR. Aluminium toxicity and iron homeostasis. *J of Inorganic Biochemistry* 2001; 87(1-2): 9-14.
- [22] Nehru B, Anand P. Oxidative damage following chronic aluminium exposure in adult and pup rat brains. *J. Trace Elem. Med. Biol* 2005; 19(2-3): 203–208.
- [23] Fernandez MT, Mira ML, Florencio MH, Jennings KR. Iron and copper chelation by flavonoids: an electrospray mass spectrometry study. *J of Inorganic Biochemistry* 2002; 92(2):105-111.
- [24] Panov AV, Andreeva L, Greenamyrea JT. Quantitative evaluation of the effects of mitochondrial permeability transition pore modifiers on accumulation of calcium phosphate: comparison of rat liver and brain mitochondria. *Arch. Biochem. Biophys* 2004; 424(1):44-52.
- [25] Valko M, Morris H, Cronin MTD. Metals, toxicity and oxidative stress. *Curr. Med. Chem* 2005;12: 1161–1208.
- [26] Hider RC, Liu ZD, Khodr HH. Metal chelation of polyphenols. *Methods Enzymol* 2001; 335:190-203.
- [27] Bors W, Michel C. Chemistry of the antioxidant effect of polyphenols. *Ann. N.Y.Acad. Sci* 2002; 957:57-69.
- [28] Cohen JE, Fields RD. Extracellular calcium depletion in synaptic transmission. *Neuroscientist* 2004; 10(1):12-17.
- [29] Vattem DA, Shetty K. Biological functionality of ellagic acid: A review. *J of Food Biochemistry* 2005; 29(3): 234-266.
- [30] Esparza JL, Gomez M, Romeu M, Mulero M, Sanchez DJ, *et al.* Aluminum-induced pro-oxidant effects in rats: protective role of exogenous melatonin. *J. Pineal Res* 2003; 35 (1):32–39.
- [31] Paynter DI. The role of dietary copper, manganese, selenium, and vitamin E in lipid peroxidation in tissues of the rat. *Biol. Trace Elem. Res* 1980; 2 (2): 121–135.
- [32] Rocchi E, Borella P, Borghi A, Paolillo F, Pradelli M, *et al.* Zinc and magnesium in liver cirrhosis. *Eur. J. Clin. Invest* 1994; 24(3): 149-155.
- [33] Poo JL, Rosas-Romero R, Rodriguez F, Silencio JL, Munoz R, *et al.* Serum zinc concentrations in two cohorts of 153 healthy subjects and 100 cirrhotic patients from Mexico City. *Dig. Dis* 1995; 13(2): 136-142.
- [34] Stamoulis I, Kouraklis G, Theocharis S. Zinc and the liver: an active interaction. *Dig. Dis. Sci* 2007; 52(7):1595-1612.
- [35] Davidson G, Kritas S, Butler R. Stressed mucosa. *Nestle Nutr. Workshop Ser. Pediatr. Program* 2007; 59:133–142.
- [36] Gumus M, Yuksel H, Evliyaoglu O, Kapan M, Boyuk A, *et al.* Effects of Ellagic Acid on Copper, Zinc, and Biochemical Values in Serum and Liver of Experimental Cholestatic Rats. *Biol Trace Elem Res* 2011; 143:(1), 386-393.