



Fractional calculus of modified special functions involving the generalized M-series in their kernels and illustrative examples

Enes Ata^a, İ Onur Kıymaz^a, Praveen Agarwal^{b,c,*}, Shilpi Jain^d, Shaher Momani^{c,e}

^a Department of Mathematics, Kırşehir Ahi Evran University, Kırşehir, 40100, Turkey

^b Department of Mathematics, Anand International College of Engineering, Jaipur 303012, India

^c Nonlinear Dynamics Research Center (NDRC), Ajman University, Ajman, United Arab Emirates

^d Department of Mathematics, Poornima College of Engineering, Jaipur 302022, India

^e Faculty of Science, The University of Jordan, Amman 11942, Jordan

ARTICLE INFO

Keywords:

Fractional derivatives and integrals
Fractional differential equations
Beta function
Gauss hypergeometric function
Confluent hypergeometric function

ABSTRACT

In this paper we apply the Riemann–Liouville, Erdelyi–Kober and Caputo fractional operators to the modified beta, modified Gauss hypergeometric and modified confluent hypergeometric functions in which the generalized M-series are included in their kernels. Furthermore, as examples, we obtain solutions of some fractional differential equations involving the above modified special functions.

1. Introduction and preliminaries

The concept of fractional calculus was first introduced in 1695 through correspondence between Leibniz and L'Hospital, as evidenced by sources such as Refs. 1–4. The letters discuss the formula

$$D^n f(x) = \frac{d^n f(x)}{dx^n}, \quad \text{for } n > 0$$

and the question of whether the number n can be extended to fractional numbers. The question's answer resulted in the creation of a new theory by scientists known as fractional calculus. Numerous studies have been conducted on fractional calculus, as seen in Refs. 5–12.

The Riemann–Liouville fractional integral (RLFI)¹ of order $\varepsilon \in \mathbb{C}$ is given by

$$\left[I_{0+}^\varepsilon f \right] (\rho) = \frac{1}{\Gamma(\varepsilon)} \int_0^\rho (\rho - \omega)^{\varepsilon-1} f(\omega) d\omega,$$

where $\rho > 0, \Re(\varepsilon) > 0$.

The Erdelyi–Kober fractional integral (EKFI)¹ for $\sigma > 0, \mu \in \mathbb{C}$ is defined by

$$\left[I_{0+;\sigma,\mu}^\varepsilon f \right] (\rho) = \frac{\sigma \rho^{-\sigma(\varepsilon+\mu)}}{\Gamma(\varepsilon)} \int_0^\rho (\rho^\sigma - \omega^\sigma)^{\varepsilon-1} \omega^{\sigma\mu+\sigma-1} f(\omega) d\omega, \quad (1)$$

where $\rho > 0, \Re(\varepsilon) > 0$.

In this paper, we will use the following Erdelyi–Kober fractional integral (EKFI) for $\sigma = 1$ in Eq. (1):

$$\left[I_{0+;\mu}^\varepsilon f \right] (\rho) = \frac{\rho^{-\varepsilon-\mu}}{\Gamma(\varepsilon)} \int_0^\rho (\rho - \omega)^{\varepsilon-1} \omega^\mu f(\omega) d\omega,$$

where $\rho > 0, \Re(\varepsilon) > 0$.

The Riemann–Liouville fractional derivative (RLFD)¹ of order $\varepsilon \in \mathbb{C}$ for $m-1 < \Re(\varepsilon) < m, m \in \mathbb{N}$ is given by

$$\left[D_{0+}^\varepsilon f \right] (\rho) = \frac{1}{\Gamma(m-\varepsilon)} \frac{d^m}{d\rho^m} \int_0^\rho (\rho - \omega)^{m-\varepsilon-1} f(\omega) d\omega,$$

where $\rho > 0, \Re(\varepsilon) > 0$.

The Caputo fractional derivative (CFD)¹ for $m-1 < \Re(\varepsilon) < m, m \in \mathbb{N}$ is defined by

$$\left[{}^c D_{0+}^\varepsilon f \right] (\rho) = \frac{1}{\Gamma(m-\varepsilon)} \int_0^\rho (\rho - \omega)^{m-\varepsilon-1} f^{(m)}(\omega) d\omega,$$

where $\rho > 0, \Re(\varepsilon) > 0$.

Special functions have been in the focus of attention of scientists due to their popularity in disciplines such as mathematics, physics, engineering. Many scientists have published quite a lot of studies in

* Corresponding author at: Department of Mathematics, Anand International College of Engineering, Jaipur 303012, India.

E-mail addresses: enesata.tr@gmail.com (E. Ata), iokiyamaz@ahievran.edu.tr (İ.O. Kıymaz), goyal.praveen2011@gmail.com (P. Agarwal), shilpijain1310@gmail.com (S. Jain), shahermmm@yahoo.com (S. Momani).

<https://doi.org/10.1016/j.padiff.2024.100720>

Received 6 January 2023; Received in revised form 31 March 2024; Accepted 14 May 2024

Available online 3 June 2024

2666-8181/© 2024 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

recent years on various generalizations of special functions, see for example Refs. 13–33 and the reference therein.

One of the popular special function is ${}_pM_q^\beta$ which defined for $\Re(\alpha) > 0$ as

$${}_pM_q^\beta(z) = {}_pM_q^\beta(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; z) = \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \frac{z^n}{\Gamma(\alpha n + \beta)}, \quad (2)$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$). If $p \leq q$ then (2) is convergent for all z . If $p = q + 1$ it is also convergent for $|z| < \delta = \alpha^\alpha$, but if $p > q + 1$ it is divergent. If $p = q + 1$, (2) can be convergent for $|z| = \delta$ depending on the conditions of the parameters³⁴. ${}_pM_q^\beta$ is also known as the generalized M-series.

Remark 1. Here, it is important to note that the generalized M-series is a special case of the Wright generalized hypergeometric function ${}_p\Psi_q(z)$; it is easy to find its relationship with various classical special and trigonometric functions that is why it is more general in nature. It is also important because its basic cases are followed by the Mittag-Leffler function and the hypergeometric functions, and all these functions have indeed found key applications in solving problems in applied sciences, chemistry, physics and biology.

The symbol $(\cdot)_n$ used to denote the Pochhammer symbol³⁵, which defined by

$$(\zeta)_n = \frac{\Gamma(\zeta + n)}{\Gamma(\zeta)} = \begin{cases} \zeta(\zeta + 1) \dots (\zeta + n - 1), & n = 1, 2, \dots, \\ 1, & n = 0. \end{cases} \quad (3)$$

The modified gamma and beta functions for $\Re(\rho) > 0$, $\Re(x) > 0$, $\Re(y) > 0$, $\Re(\alpha) > 0$ was introduced by Ata in Ref. 16, respectively, as follows:

$$\begin{aligned} M_{p,q}^{(\alpha,\beta)}(x; \rho) &= M_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; x; \rho) \\ &= \int_0^\infty \Delta^{x-1} {}_pM_q^\beta\left(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; -\Delta - \frac{\rho}{\Delta}\right) d\Delta \end{aligned}$$

and

$$\begin{aligned} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) &= M_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; x, y; \rho) \\ &= \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} {}_pM_q^\beta\left(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \frac{-\rho}{\Delta(1 - \Delta)}\right) d\Delta, \end{aligned}$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

In the same paper the author used the modified beta function to defined the modified Gauss and confluent hypergeometric functions for $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$, $\Re(\rho) > 0$, respectively, as follows:

$$\begin{aligned} M_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; \rho) &= M_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \chi_1, \chi_2; \chi_3; \rho) \\ &= \sum_{n=0}^{\infty} (\chi_1)_n \frac{M_{p,q}^{(\alpha,\beta)}(\chi_2 + n, \chi_3 - \chi_2; \rho)}{B(\chi_2, \chi_3 - \chi_2)} \frac{z^n}{n!}, \quad (|z| < 1) \end{aligned}$$

and

$$\begin{aligned} M_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) &= M_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \chi_2; \chi_3; z; \rho) \\ &= \sum_{n=0}^{\infty} \frac{M_{p,q}^{(\alpha,\beta)}(\chi_2 + n, \chi_3 - \chi_2; \rho)}{B(\chi_2, \chi_3 - \chi_2)} \frac{z^n}{n!}, \end{aligned}$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

If we take $\rho = 0$ and $p = q = \xi_1 = \eta_1 = \alpha = \beta = 1$ in the modified special functions given above, we get the classic forms of corresponding special functions³⁵. Additionally, the author called the modified gamma, modified beta, modified Gauss hypergeometric, and modified confluent hypergeometric functions as the M-gamma, M-beta, M-Gauss hypergeometric, and M-confluent hypergeometric functions, respectively.

It should be noted that the relationship between the classic gamma and beta functions, as described in Ref. 35, is as follows:

$$B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x + y)}, \quad (\Re(x) > 0, \Re(y) > 0). \quad (4)$$

The paper proposes the use of modified special functions involving generalized M-series, which contain more parameters. It is expected that this will expand the application areas.

The remainder of this paper is planned as follows: In Section 2, we apply Riemann–Liouville, Caputo and Erdelyi–Kober fractional operators to M-beta, M-Gauss hypergeometric and M-confluent hypergeometric functions. In Section 3, we obtain solutions of fractional differential equations including M-beta, M-Gauss hypergeometric and M-confluent hypergeometric functions.

2. Fractional calculus of modified special functions

In this section, we apply Riemann–Liouville fractional integral, Erdelyi–Kober fractional integral, Riemann–Liouville fractional derivative, and Caputo fractional derivative operators to M-beta, M-Gauss hypergeometric, and M-confluent hypergeometric functions.

2.1. Riemann–Liouville fractional integral (RLFI)

Theorem 2.1. Let $\Re(\epsilon) > 0$, $\Re(x) > 0$, $\Re(y) > 0$, $\Re(\alpha) > 0$. Then, applying the RLFI to the M-beta function yields the following formula:

$$\left[I_{0+}^\epsilon M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{{}_pM_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; x, y; \rho)}{\Gamma(1 + \epsilon)\rho^{-\epsilon}},$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the RLFI to the M-beta function, we have

$$\left[I_{0+}^\epsilon M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{1}{\Gamma(\epsilon)} \int_0^\rho (\rho - \omega)^{\epsilon-1} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) d\omega.$$

Using the definition of the M-beta function, we get

$$\begin{aligned} \left[I_{0+}^\epsilon M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) &= \frac{1}{\Gamma(\epsilon)} \int_0^\rho (\rho - \omega)^{\epsilon-1} \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \\ &\quad \times {}_pM_q^\beta\left(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \frac{-\omega}{\Delta(1 - \Delta)}\right) d\Delta d\omega. \end{aligned}$$

Using the definition of the generalized M-series, we obtain

$$\begin{aligned} \left[I_{0+}^\epsilon M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) &= \frac{1}{\Gamma(\epsilon)} \int_0^\rho (\rho - \omega)^{\epsilon-1} \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \\ &\quad \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \frac{\left(\frac{-\omega}{\Delta(1 - \Delta)}\right)^n}{\Gamma(\alpha n + \beta)} d\Delta d\omega. \end{aligned}$$

Taking $\omega = \rho u$ and using Eq. (4), we have

$$\begin{aligned} \left[I_{0+}^\epsilon M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) &= \rho^\epsilon \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \\ &\quad \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \frac{\left(\frac{-\rho}{\Delta(1 - \Delta)}\right)^n}{\Gamma(\alpha n + \beta)} \frac{\Gamma(n + 1)}{\Gamma(n + 1 + \epsilon)} d\Delta. \end{aligned}$$

Multiplying by $\frac{\Gamma(1 + \epsilon)}{\Gamma(1 + \epsilon)}$ and using Eq. (3), we get

$$\begin{aligned} \left[I_{0+}^\epsilon M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) &= \frac{\rho^\epsilon}{\Gamma(1 + \epsilon)} \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \\ &\quad \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n (1)_n}{(\eta_1)_n \dots (\eta_q)_n (1 + \epsilon)_n} \frac{\left(\frac{-\rho}{\Delta(1 - \Delta)}\right)^n}{\Gamma(\alpha n + \beta)} d\Delta. \end{aligned}$$

Considering the definition of the generalized M-series, we obtain

$$\begin{aligned} \left[I_{0+}^\epsilon M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) &= \frac{\rho^\epsilon}{\Gamma(1 + \epsilon)} \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \\ &\quad \times {}_{p+1}M_{q+1}^\beta\left(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; \frac{-\rho}{\Delta(1 - \Delta)}\right) d\Delta. \end{aligned}$$

Thus, we have the desired formula as:

$$\left[I_{0+}^\epsilon M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{{}_pM_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; x, y; \rho)}{\Gamma(1 + \epsilon)\rho^{-\epsilon}},$$

which completes the proof. $\square$

Theorem 2.2. Let $\Re(\epsilon) > 0$, $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$. Then, applying the RLFI to the M-Gauss hypergeometric function yields the following formula:

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{M_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; \chi_1, \chi_2; \chi_3; z; \rho)}{\Gamma(1 + \epsilon)\rho^{-\epsilon}},$$

where $\arg(1 - z) < \pi$ and $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the RLFI to the M-Gauss hypergeometric function, we have

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{1}{\Gamma(\epsilon)} \int_0^{\rho} (\rho - \omega)^{\epsilon-1} M_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \omega) d\omega.$$

Taking $\omega = \rho u$ and using Eq. (4), we get

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\rho^{\epsilon}}{B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1 - \Delta)^{\chi_3-\chi_2-1} (1 - z\Delta)^{-\chi_1} \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \left(\frac{-\rho}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1)}{\Gamma(\alpha n + \beta) \Gamma(n+1 + \epsilon)} d\Delta.$$

Multiplying by $\frac{\Gamma(1+\epsilon)}{\Gamma(1+\epsilon)}$ and using Eq. (3), we obtain

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\rho^{\epsilon}}{\Gamma(1 + \epsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1 - \Delta)^{\chi_3-\chi_2-1} (1 - z\Delta)^{-\chi_1} \times {}_pM_{q+1}^{\beta} \left(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; \frac{-\rho}{\Delta(1-\Delta)} \right) d\Delta.$$

Thus, we have the desired formula as:

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{M_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; \chi_1, \chi_2; \chi_3; z; \rho)}{\Gamma(1 + \epsilon)\rho^{-\epsilon}},$$

which completes the proof. $\square$

Theorem 2.3. Let $\Re(\epsilon) > 0$, $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$. Then, applying the RLFI to the M-confluent hypergeometric function yields the following formula:

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{M_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; \chi_2; \chi_3; z; \rho)}{\Gamma(1 + \epsilon)\rho^{-\epsilon}},$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the RLFI to the M-confluent hypergeometric function, we have

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{1}{\Gamma(\epsilon)} \int_0^{\rho} (\rho - \omega)^{\epsilon-1} M_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \omega) d\omega.$$

Taking $\omega = \rho u$ and using Eq. (4), we get

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\rho^{\epsilon}}{B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1 - \Delta)^{\chi_3-\chi_2-1} \exp(z\Delta) \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \left(\frac{-\rho}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1)}{\Gamma(\alpha n + \beta) \Gamma(n+1 + \epsilon)} d\Delta.$$

Multiplying by $\frac{\Gamma(1+\epsilon)}{\Gamma(1+\epsilon)}$ and using Eq. (3), we obtain

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\rho^{\epsilon}}{\Gamma(1 + \epsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1 - \Delta)^{\chi_3-\chi_2-1} \exp(z\Delta) \times {}_pM_{q+1}^{\beta} \left(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; \frac{-\rho}{\Delta(1-\Delta)} \right) d\Delta.$$

Thus, we have the desired formula as:

$$\left[I_{0+}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{M_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \epsilon; \chi_2; \chi_3; z; \rho)}{\Gamma(1 + \epsilon)\rho^{-\epsilon}},$$

which completes the proof. $\square$

2.2. Erdelyi-Kober fractional integral (EKFI)

Theorem 2.4. Let $\Re(\epsilon) > 0$, $\Re(1 + \mu) > 0$, $\Re(1 + \mu + \epsilon) > 0$, $\Re(x) > 0$, $\Re(y) > 0$, $\Re(\alpha) > 0$. Then, applying the EKFI to the M-beta function yields the following formula:

$$\left[I_{0+;\mu}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\Gamma(1 + \mu) M_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1 + \mu; \eta_1, \dots, \eta_q, 1 + \mu + \epsilon; x, y; \rho)}{\Gamma(1 + \mu + \epsilon)},$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the EKFI to the M-beta function, we have

$$\left[I_{0+;\mu}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\rho^{-\epsilon-\mu}}{\Gamma(\epsilon)} \int_0^{\rho} (\rho - \omega)^{\epsilon-1} \omega^{\mu} M_{p,q}^{(\alpha,\beta)}(x, y; \omega) d\omega.$$

Using the definition of the M-beta function, we get

$$\left[I_{0+;\mu}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\rho^{-\epsilon-\mu}}{\Gamma(\epsilon)} \int_0^{\rho} (\rho - \omega)^{\epsilon-1} \omega^{\mu} \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \times {}_pM_q^{\beta} \left(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \frac{-\omega}{\Delta(1-\Delta)} \right) d\Delta d\omega.$$

Using the definition of the generalized M-series, we obtain

$$\left[I_{0+;\mu}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\rho^{-\epsilon-\mu}}{\Gamma(\epsilon)} \int_0^{\rho} (\rho - \omega)^{\epsilon-1} \omega^{\mu} \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \left(\frac{-\omega}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1 + \mu)}{\Gamma(\alpha n + \beta)} d\Delta d\omega.$$

Taking $\omega = \rho u$ and using Eq. (4), we have

$$\left[I_{0+;\mu}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \left(\frac{-\rho}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1 + \mu)}{\Gamma(\alpha n + \beta) \Gamma(n+1 + \mu + \epsilon)} d\Delta.$$

Multiplying by $\frac{\Gamma(1+\mu)\Gamma(1+\mu+\epsilon)}{\Gamma(1+\mu)\Gamma(1+\mu+\epsilon)}$ and using Eq. (3), we get

$$\left[I_{0+;\mu}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\Gamma(1 + \mu)}{\Gamma(1 + \mu + \epsilon)} \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n (1 + \mu)_n}{(\eta_1)_n \dots (\eta_q)_n (1 + \mu + \epsilon)_n} \left(\frac{-\rho}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n + \beta)}{\Gamma(\alpha n + \beta)} d\Delta.$$

Considering the definition of the generalized M-series, we obtain

$$\left[I_{0+;\mu}^{\epsilon} M_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\Gamma(1 + \mu)}{\Gamma(1 + \mu + \epsilon)} \int_0^1 \Delta^{x-1} (1 - \Delta)^{y-1} \times {}_{p+1}M_{q+1}^{\beta} \left(\xi_1, \dots, \xi_p, 1 + \mu; \eta_1, \dots, \eta_q, 1 + \mu + \epsilon; \frac{-\rho}{\Delta(1-\Delta)} \right) d\Delta.$$

Thus, we have the desired formula as:

$$\left[I_{0+;\mu}^{\varepsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\Gamma(1+\mu) {}^M B_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1+\mu; \eta_1, \dots, \eta_q, 1+\mu+\varepsilon; x, y; \rho)}{\Gamma(1+\mu+\varepsilon)},$$

which completes the proof. $\square$

Theorem 2.5. Let $\Re(\varepsilon) > 0$, $\Re(1+\mu) > 0$, $\Re(1+\mu+\varepsilon) > 0$, $\Re(\alpha) > 0$, $\Re(\chi_3) > \Re(\chi_2) > 0$. Then, applying the EKFI to the M-Gauss hypergeometric function yields the following formula:

$$\left[I_{0+;\mu}^{\varepsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\Gamma(1+\mu) {}^M F_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1+\mu; \eta_1, \dots, \eta_q, 1+\mu+\varepsilon; \chi_1, \chi_2; \chi_3; z; \rho)}{\Gamma(1+\mu+\varepsilon)},$$

where $\arg(1-z) < \pi$ and $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the EKFI to the M-Gauss hypergeometric function, we have

$$\left[I_{0+;\mu}^{\varepsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\rho^{-\varepsilon-\mu}}{\Gamma(\varepsilon)} \int_0^{\rho} (\rho-\omega)^{\varepsilon-1} \omega^{\mu} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \omega) d\omega.$$

Taking $\omega = \rho u$ and using Eq. (4), we get

$$\left[I_{0+;\mu}^{\varepsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{1}{B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} (1-z\Delta)^{-\chi_1} \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \left(\frac{-\rho}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1+\mu)}{\Gamma(\alpha n + \beta)} \frac{\Gamma(n+1+\mu+\varepsilon)}{\Gamma(n+1+\mu+\varepsilon)} d\Delta.$$

Multiplying by $\frac{\Gamma(1+\mu)\Gamma(1+\mu+\varepsilon)}{\Gamma(1+\mu)\Gamma(1+\mu+\varepsilon)}$ and using Eq. (3), we obtain

$$\left[I_{0+;\mu}^{\varepsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\Gamma(1+\mu)}{\Gamma(1+\mu+\varepsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} (1-z\Delta)^{-\chi_1} \times {}_pM_{q+1}^{\beta} \left(\xi_1, \dots, \xi_p, 1+\mu; \eta_1, \dots, \eta_q, 1+\mu+\varepsilon; \frac{-\rho}{\Delta(1-\Delta)} \right) d\Delta.$$

Thus, we have the desired formula as:

$$\left[I_{0+;\mu}^{\varepsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\Gamma(1+\mu) {}^M F_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1+\mu; \eta_1, \dots, \eta_q, 1+\mu+\varepsilon; \chi_1, \chi_2; \chi_3; z; \rho)}{\Gamma(1+\mu+\varepsilon)},$$

which completes the proof. $\square$

Theorem 2.6. Let $\Re(\varepsilon) > 0$, $\Re(1+\mu) > 0$, $\Re(1+\mu+\varepsilon) > 0$, $\Re(\alpha) > 0$, $\Re(\chi_3) > \Re(\chi_2) > 0$. Then, applying the EKFI to the M-confluent hypergeometric function yields the following formula:

$$\left[I_{0+;\mu}^{\varepsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\Gamma(1+\mu) {}^M \Phi_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1+\mu; \eta_1, \dots, \eta_q, 1+\mu+\varepsilon; \chi_2; \chi_3; z; \rho)}{\Gamma(1+\mu+\varepsilon)},$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the EKFI to the M-confluent hypergeometric function, we have

$$\left[I_{0+;\mu}^{\varepsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\rho^{-\varepsilon-\mu}}{\Gamma(\varepsilon)} \int_0^{\rho} (\rho-\omega)^{\varepsilon-1} \omega^{\mu} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \omega) d\omega.$$

Taking $\omega = \rho u$ and using Eq. (4), we get

$$\left[I_{0+;\mu}^{\varepsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{1}{B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} \exp(z\Delta) \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \left(\frac{-\rho}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1+\mu)}{\Gamma(\alpha n + \beta)} \frac{\Gamma(n+1+\mu+\varepsilon)}{\Gamma(n+1+\mu+\varepsilon)} d\Delta.$$

Multiplying by $\frac{\Gamma(1+\mu)\Gamma(1+\mu+\varepsilon)}{\Gamma(1+\mu)\Gamma(1+\mu+\varepsilon)}$ and using Eq. (3), we obtain

$$\left[I_{0+;\mu}^{\varepsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\Gamma(1+\mu)}{\Gamma(1+\mu+\varepsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} \exp(z\Delta) \times {}_pM_{q+1}^{\beta} \left(\xi_1, \dots, \xi_p, 1+\mu; \eta_1, \dots, \eta_q, 1+\mu+\varepsilon; \frac{-\rho}{\Delta(1-\Delta)} \right) d\Delta.$$

Thus, we have the desired formula as:

$$\left[I_{0+;\mu}^{\varepsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{\Gamma(1+\mu) {}^M \Phi_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1+\mu; \eta_1, \dots, \eta_q, 1+\mu+\varepsilon; \chi_2; \chi_3; z; \rho)}{\Gamma(1+\mu+\varepsilon)},$$

which completes the proof. $\square$

2.3. Riemann–Liouville fractional derivative (RLFD)

Theorem 2.7. Let $0 < \Re(\varepsilon) < 1$, $\Re(x) > 0$, $\Re(y) > 0$, $\Re(\alpha) > 0$. Then, applying the RLFD to the M-beta function yields the following formula:

$$\left[D_{0+}^{\varepsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{{}^M B_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\varepsilon; x, y; \rho)}{\Gamma(1-\varepsilon)\rho^{\varepsilon}},$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the RLFD for $m = 1$ to the M-beta function, we have

$$\left[D_{0+}^{\varepsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{1}{\Gamma(1-\varepsilon)} \frac{d}{d\rho} \int_0^{\rho} (\rho-\omega)^{-\varepsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \omega) d\omega.$$

Using the definition of the M-beta function, we get

$$\left[D_{0+}^{\varepsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{1}{\Gamma(1-\varepsilon)} \frac{d}{d\rho} \int_0^{\rho} (\rho-\omega)^{-\varepsilon} \int_0^1 \Delta^{x-1} (1-\Delta)^{y-1} \times {}_pM_q^{\beta} \left(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \frac{-\omega}{\Delta(1-\Delta)} \right) d\Delta d\omega.$$

Using the definition of the generalized M-series, we obtain

$$\left[D_{0+}^{\varepsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{1}{\Gamma(1-\varepsilon)} \frac{d}{d\rho} \int_0^{\rho} (\rho-\omega)^{-\varepsilon} \int_0^1 \Delta^{x-1} (1-\Delta)^{y-1} \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \left(\frac{-\omega}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1)}{\Gamma(\alpha n + \beta)} d\Delta d\omega.$$

Taking $\omega = \rho u$ and using Eq. (4), we have

$$\left[D_{0+}^{\varepsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\rho^{-\varepsilon}}{\Gamma(1-\varepsilon)} \int_0^1 \Delta^{x-1} (1-\Delta)^{y-1} \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n}{(\eta_1)_n \dots (\eta_q)_n} \left(\frac{-\rho}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1)\Gamma(1-\varepsilon)}{\Gamma(\alpha n + \beta)} \frac{\Gamma(n+1-\varepsilon)}{\Gamma(n+1-\varepsilon)} d\Delta.$$

Using Eq. (3), we get

$$\left[D_{0+}^{\varepsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\rho^{-\varepsilon}}{\Gamma(1-\varepsilon)} \int_0^1 \Delta^{x-1} (1-\Delta)^{y-1} \times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n (1)_n}{(\eta_1)_n \dots (\eta_q)_n (1-\varepsilon)_n} \left(\frac{-\rho}{\Delta(1-\Delta)} \right)^n \frac{\Gamma(n+1)}{\Gamma(\alpha n + \beta)} d\Delta.$$

Considering the definition of the generalized M-series, we obtain

$$\left[D_{0+}^{\epsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{\rho^{-\epsilon}}{\Gamma(1-\epsilon)} \int_0^1 \Delta^{x-1} (1-\Delta)^{y-1} \times {}^M M_{q+1}^{\beta} \left(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; \frac{-\rho}{\Delta(1-\Delta)} \right) d\Delta.$$

Thus, we have the desired formula as:

$$\left[D_{0+}^{\epsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{{}^M B_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; x, y; \rho)}{\Gamma(1-\epsilon)\rho^{\epsilon}},$$

which completes the proof. $\square$

Theorem 2.8. Let $0 < \Re(\epsilon) < 1$, $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$. Then, applying the RLFD to the M-Gauss hypergeometric function yields the following formula:

$$\left[D_{0+}^{\epsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{{}^M F_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; \chi_1, \chi_2; \chi_3; z; \rho)}{\Gamma(1-\epsilon)\rho^{\epsilon}},$$

where $\arg(1-z) < \pi$ and $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the RLFD for $m = 1$ to the M-Gauss hypergeometric function, we have

$$\left[D_{0+}^{\epsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{1}{\Gamma(1-\epsilon)} \frac{d}{d\rho} \int_0^{\rho} (\rho-\omega)^{-\epsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \omega) d\omega.$$

Taking $\omega = \rho u$ and using Eq. (4), we get

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{\rho^{-\epsilon}}{\Gamma(1-\epsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} (1-z\Delta)^{-\chi_1} \\ &\times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n \left(\frac{-\rho}{\Delta(1-\Delta)}\right)^n}{(\eta_1)_n \dots (\eta_q)_n \Gamma(an + \beta)} \frac{\Gamma(n+1)\Gamma(1-\epsilon)}{\Gamma(n+1-\epsilon)} d\Delta. \end{aligned}$$

Using Eq. (3), we obtain

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{\rho^{-\epsilon}}{\Gamma(1-\epsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} (1-z\Delta)^{-\chi_1} \\ &\times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n (1)_n \left(\frac{-\rho}{\Delta(1-\Delta)}\right)^n}{(\eta_1)_n \dots (\eta_q)_n (1-\epsilon)_n \Gamma(an + \beta)} d\Delta. \end{aligned}$$

Considering the definition of the generalized M-series, we have

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{\rho^{-\epsilon}}{\Gamma(1-\epsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} (1-z\Delta)^{-\chi_1} \\ &\times {}^M M_{q+1}^{\beta} \left(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; \frac{-\rho}{\Delta(1-\Delta)} \right) d\Delta. \end{aligned}$$

Thus, we get the desired formula as:

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{{}^M F_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; \chi_1, \chi_2; \chi_3; z; \rho)}{\Gamma(1-\epsilon)\rho^{\epsilon}}, \end{aligned}$$

which completes the proof. $\square$

Theorem 2.9. Let $0 < \Re(\epsilon) < 1$, $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$. Then, applying the RLFD to the M-confluent hypergeometric function yields the following formula:

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{{}^M \Phi_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; \chi_2; \chi_3; z; \rho)}{\Gamma(1-\epsilon)\rho^{\epsilon}}, \end{aligned}$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the RLFD for $m = 1$ to the M-confluent hypergeometric function, we have

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{1}{\Gamma(1-\epsilon)} \frac{d}{d\rho} \int_0^{\rho} (\rho-\omega)^{-\epsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \omega) d\omega. \end{aligned}$$

Taking $\omega = \rho u$ and using Eq. (4), we get

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{\rho^{-\epsilon}}{\Gamma(1-\epsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} \exp(z\Delta) \\ &\times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n \left(\frac{-\rho}{\Delta(1-\Delta)}\right)^n}{(\eta_1)_n \dots (\eta_q)_n \Gamma(an + \beta)} \frac{\Gamma(n+1)\Gamma(1-\epsilon)}{\Gamma(n+1-\epsilon)} d\Delta. \end{aligned}$$

Using Eq. (3), we obtain

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{\rho^{-\epsilon}}{\Gamma(1-\epsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} \exp(z\Delta) \\ &\times \sum_{n=0}^{\infty} \frac{(\xi_1)_n \dots (\xi_p)_n (1)_n \left(\frac{-\rho}{\Delta(1-\Delta)}\right)^n}{(\eta_1)_n \dots (\eta_q)_n (1-\epsilon)_n \Gamma(an + \beta)} d\Delta. \end{aligned}$$

Considering the definition of the generalized M-series, we have

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{\rho^{-\epsilon}}{\Gamma(1-\epsilon)B(\chi_2, \chi_3 - \chi_2)} \int_0^1 \Delta^{\chi_2-1} (1-\Delta)^{\chi_3-\chi_2-1} \exp(z\Delta) \\ &\times {}^M M_{q+1}^{\beta} \left(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; \frac{-\rho}{\Delta(1-\Delta)} \right) d\Delta. \end{aligned}$$

Thus, we get the desired formula as:

$$\begin{aligned} \left[D_{0+}^{\epsilon} {}^M \Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{{}^M \Phi_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; \chi_2; \chi_3; z; \rho)}{\Gamma(1-\epsilon)\rho^{\epsilon}}, \end{aligned}$$

which completes the proof. $\square$

2.4. Caputo fractional derivative (CFD)

Theorem 2.10. Let $0 < \Re(\epsilon) < 1$, $\Re(x) > 0$, $\Re(y) > 0$, $\Re(\alpha) > 0$. Then, applying the CFD to the M-beta function yields the following formula:

$$\left[{}^c D_{0+}^{\epsilon} {}^M B_{p,q}^{(\alpha,\beta)}(x, y; \rho) \right] (\rho) = \frac{{}^M B_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; x, y; \rho)}{\Gamma(1-\epsilon)\rho^{\epsilon}},$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the CFD for $m = 1$ to the M-beta function and performing the required calculations, the proof is complete. $\square$

Theorem 2.11. Let $0 < \Re(\epsilon) < 1$, $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$. Then, applying the CFD to the M-Gauss hypergeometric function yields the following formula:

$$\begin{aligned} \left[{}^c D_{0+}^{\epsilon} {}^M F_{p,q}^{(\alpha,\beta)}(\chi_1, \chi_2; \chi_3; z; \rho) \right] (\rho) &= \frac{{}^M F_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1-\epsilon; \chi_1, \chi_2; \chi_3; z; \rho)}{\Gamma(1-\epsilon)\rho^{\epsilon}}, \end{aligned}$$

where $\arg(1-z) < \pi$ and $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the CFD for $m = 1$ to the M-Gauss hypergeometric function and performing the required calculations, the proof is complete. $\square$

Theorem 2.12. Let $0 < \Re(\varepsilon) < 1$, $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$. Then, applying the CFD to the M-confluent hypergeometric function yields the following formula:

$$\left[{}^c D_{0+}^\varepsilon M\Phi_{p,q}^{(\alpha,\beta)}(\chi_2; \chi_3; z; \rho) \right] (\rho) = \frac{M\Phi_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 - \varepsilon; \chi_2; \chi_3; z; \rho)}{\Gamma(1 - \varepsilon)\rho^\varepsilon},$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Applying the CFD for $m = 1$ to the M-confluent hypergeometric function and performing the required calculations, the proof is complete. $\square$

3. Illustrative examples

Let $\Re(\varepsilon) > 0$ and $m - 1 < \Re(\varepsilon) < m$ for $m \in \mathbb{N}$. If $1 \leq p \leq \infty$ and $f \in I_{a+}^\varepsilon(L_p)$, then Ref. 1:

$$\left[I_{a+}^\varepsilon D_{a+}^\varepsilon f \right] (\rho) = f(\rho), \quad (5)$$

holds almost everywhere on $[a, b]$ and where $I_{a+}^\varepsilon(L_p)$ as follows¹:

$$I_{a+}^\varepsilon(L_p) = \left\{ f : f(\rho) = \left[I_{a+}^\varepsilon \varphi(\rho) \right] (\rho), \varphi(\rho) \in L_p(a, b) \right\}.$$

In this section, we obtain solutions of fractional differential equations involving the modified beta, Gauss hypergeometric and confluent hypergeometric functions using Eq. (5).

Example 1. Let $1 < \Re(\varepsilon) < 2$, $\Re(x) > 0$, $\Re(y) > 0$, $\Re(\alpha) > 0$. We consider the fractional differential equation

$$\left[D_{0+}^\varepsilon f \right] (\rho) = M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; x, y; \frac{\rho}{\varepsilon}), \quad (6)$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Considering Eq. (5) for $a = 0$ and applying the RLFI to the fractional differential equation, we have

$$\left[I_{0+}^\varepsilon D_{0+}^\varepsilon f \right] (\rho) = \left[I_{0+}^\varepsilon M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; x, y; \frac{\rho}{\varepsilon}) \right] (\rho).$$

That is,

$$f(\rho) = \frac{1}{\Gamma(\varepsilon)} \int_0^\rho (\rho - \omega)^{\varepsilon-1} M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; x, y; \frac{\omega}{\varepsilon}) d\omega.$$

Taking $\omega = \rho u$ and performing the necessary calculations, we get the solution function as:

$$f(\rho) = \frac{M\Phi_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \varepsilon; x, y; \frac{\rho}{\varepsilon})}{\Gamma(1 + \varepsilon)\rho^{-\varepsilon}}. \quad \square$$

Corollary 3.1. If we use the generalized beta function defined by Chaudhry et al.²⁰ instead of the modified beta function in Eq. (6), we obtain the solution function as follows:

$$f(\rho) = \rho^\varepsilon \sum_{n=0}^{\infty} \frac{\Gamma(x - n)\Gamma(y - n)}{\Gamma(n + 1 + \varepsilon)\Gamma(x + y - 2n)} \left(-\frac{\rho}{\varepsilon} \right)^n.$$

Example 2. Let $1 < \Re(\varepsilon) < 2$, $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$. We consider the fractional differential equation

$$\left[D_{0+}^\varepsilon f \right] (\rho) = M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \chi_1, \chi_2; \chi_3; z; \frac{\rho}{\varepsilon}), \quad (7)$$

where $\arg(1 - z) < \pi$ and $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Considering Eq. (5) for $a = 0$ and applying the RLFI to the fractional differential equation, we have

$$\left[I_{0+}^\varepsilon D_{0+}^\varepsilon f \right] (\rho) = \left[I_{0+}^\varepsilon M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \chi_1, \chi_2; \chi_3; z; \frac{\rho}{\varepsilon}) \right] (\rho).$$

That is,

$$f(\rho) = \frac{1}{\Gamma(\varepsilon)} \int_0^\rho (\rho - \omega)^{\varepsilon-1} M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \chi_1, \chi_2; \chi_3; z; \frac{\omega}{\varepsilon}) d\omega.$$

Taking $\omega = \rho u$ and performing the necessary calculations, we get the solution function as:

$$f(\rho) = \frac{M\Phi_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \varepsilon; \chi_1, \chi_2; \chi_3; z; \frac{\rho}{\varepsilon})}{\Gamma(1 + \varepsilon)\rho^{-\varepsilon}}. \quad \square$$

Corollary 3.2. If we take the generalized Gauss hypergeometric function defined by Chaudhry et al.²¹ instead of the modified Gauss hypergeometric function in Eq. (7), we obtain the solution function as follows:

$$f(\rho) = \frac{\Gamma(\chi_3)\rho^\varepsilon}{\Gamma(\chi_1)\Gamma(\chi_2)\Gamma(\chi_3 - \chi_2)} \times \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \frac{\Gamma(\chi_1 + k)\Gamma(\chi_2 + k - n)\Gamma(\chi_3 - \chi_2 - n)}{\Gamma(1 + n + \varepsilon)\Gamma(\chi_3 + k - 2n)} \left(-\frac{\rho}{\varepsilon} \right)^n \frac{z^k}{k!}.$$

Example 3. Let $1 < \Re(\varepsilon) < 2$, $\Re(\chi_3) > \Re(\chi_2) > 0$, $\Re(\alpha) > 0$. We consider the fractional differential equation

$$\left[D_{\rho}^\varepsilon f \right] (\rho) = M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \chi_2; \chi_3; z; \frac{\rho}{\varepsilon}), \quad (8)$$

where $\xi_i, \eta_j \neq 0, -1, -2, \dots$ ($i = 1, \dots, p; j = 1, \dots, q$).

Proof. Considering Eq. (5) for $a = 0$ and applying the RLFI to the fractional differential equation, we have

$$\left[I_{0+}^\varepsilon D_{0+}^\varepsilon f \right] (\rho) = \left[I_{0+}^\varepsilon M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \chi_2; \chi_3; z; \frac{\rho}{\varepsilon}) \right] (\rho).$$

That is,

$$f(\rho) = \frac{1}{\Gamma(\varepsilon)} \int_0^\rho (\rho - \omega)^{\varepsilon-1} M\Phi_{p,q}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p; \eta_1, \dots, \eta_q; \chi_2; \chi_3; z; \frac{\omega}{\varepsilon}) d\omega.$$

Taking $\omega = \rho u$ and performing the necessary calculations, we get the solution function as:

$$f(\rho) = \frac{M\Phi_{p+1,q+1}^{(\alpha,\beta)}(\xi_1, \dots, \xi_p, 1; \eta_1, \dots, \eta_q, 1 + \varepsilon; \chi_2; \chi_3; z; \frac{\rho}{\varepsilon})}{\Gamma(1 + \varepsilon)\rho^{-\varepsilon}}. \quad \square$$

Corollary 3.3. If we take the generalized confluent hypergeometric function defined by Chaudhry et al.²¹ instead of the modified confluent hypergeometric function in Eq. (8), we obtain the solution function as follows:

$$f(\rho) = \frac{\Gamma(\chi_3)\rho^\varepsilon}{\Gamma(\chi_2)\Gamma(\chi_3 - \chi_2)} \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \frac{\Gamma(\chi_2 + k - n)\Gamma(\chi_3 - \chi_2 - n)}{\Gamma(1 + n + \varepsilon)\Gamma(\chi_3 + k - 2n)} \left(-\frac{\rho}{\varepsilon} \right)^n \frac{z^k}{k!}.$$

4. Conclusion

In this paper, we applied RLFI, EKFI, RLFD and CFD to M-beta, M-Gauss hypergeometric and M-confluent hypergeometric functions. We would also like to point out that the results of the RLFD and CFD applied to the M-beta, M-Gauss hypergeometric and M-confluent hypergeometric functions overlap. Then, we obtained solutions of fractional differential equations involving the M-beta, M-Gauss hypergeometric and M-confluent hypergeometric functions.

In Fig. 1, the approximate behavior of the solutions of fractional differential equations involving the modified beta function and the generalized beta function defined by Chaudhry et al.²⁰ is compared.

In Fig. 2, the approximate behavior of the solutions of fractional differential equations involving the modified Gauss hypergeometric function and the generalized Gauss hypergeometric function defined by Chaudhry et al.²¹ is compared.

In Fig. 3, the approximate behavior of the solutions of fractional differential equations involving the modified confluent hypergeometric function and the generalized confluent hypergeometric function defined by Chaudhry et al.²¹ is compared.




Fig. 1. The behavior of approximate solutions $f(\rho)$ of Example 1 and Corollary 3.1 for $\varepsilon = 3/2$, where $p = q = 1$, $x = y = 4$, $0 < \rho < 4$ and generalized M-series and exponential series indexes $n = 0, 1$.




Fig. 2. The behavior of approximate solutions $f(\rho)$ of Example 2 and Corollary 3.2 for $\varepsilon = 3/2$, where $p = q = 1$, $\chi_1 = \chi_2 = 2$, $\chi_3 = 4$, $z = 1/2$, $0 < \rho < 4$ and generalized M-series and exponential series indexes $n = 0, 1$ and modified Gauss hypergeometric series and generalized Gauss hypergeometric series indexes $k = 0, 1$.




Fig. 3. The behavior of approximate solutions $f(\rho)$ of Example 3 and Corollary 3.3 for $\varepsilon = 3/2$, where $p = q = 1$, $\chi_2 = 2$, $\chi_3 = 4$, $z = 1/2$, $0 < \rho < 4$ and generalized M-series and exponential series indexes $n = 0, 1$, modified confluent hypergeometric series and generalized confluent hypergeometric series indexes $k = 0, 1$.

Note that in Figs. 1, 2, and 3, the red line represents the approximate behavior of the solutions of fractional differential equations involving the generalized special functions defined by Chaudhry et al. and the blue, yellow and green lines represent the approximate behavior of the solutions of fractional differential equations involving the modified special functions.

Declaration of competing interest

The author declares that there is no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

Data availability

No data was used for the research described in the article.

Acknowledgments

This work was partly presented in the 6th International Conference on Computational Mathematics and Engineering Sciences (CMES-2022) which organized by Ordu University on May 20–22, 2022 in Ordu-Turkey. The authors are thankful to the referees for making valuable suggestions leading to the better presentations of this paper.

References

- Kilbas AA, Srivastava HM, Trujillo JJ. *Theory and applications of fractional differential equations*. Amsterdam: North-Holland Mathematics Studies 204; Elsevier Science B.V.; 2006.
- Podlubny I. *Fractional differential equations: an introduction to fractional derivatives, fractional differential equations, to methods of their solution and some of their applications*. Academic Press; 1999.
- Samko SG, Kilbas AA, Marichev OI. *Fractional integrals and derivatives: theory and applications*. Gordon and Breach Science Publishers; 1993.
- Debnath L, Bhatta D. *Integral transforms and their applications*. 3rd ed. Boca Raton, London, New York: CRC Press; 2015.
- Baleanu D, Jajarmi A, Mohammadi H, Rezapour S. A new study on the mathematical modelling of human liver with Caputo-Fabrizio fractional derivative. *Chaos, Solitons & Fractals*. 2020;134:109705.
- Jajarmi A, Baleanu D, Sajjadi SS, Nieto JJ. Analysis and some applications of a regularized Ψ -Hilfer fractional derivative. *J Comput Appl Math*. 2022;415:114476.
- Baleanu D, Ghassabzade FA, Nieto JJ, Jajarmi A. On a new and generalized fractional model for a real cholera outbreak. *Alexandria Eng J*. 2022;61:9175–9186.
- Deftlerli Ö, Baleanu D, Jajarmi A, Sajjadi SS, Alshaikh N, Asad JH. Fractional treatment: an accelerated mass-spring system. *Romanian Rep Phys*. 2022;74(4):122.
- Bansal MK, Kumar D, Khan I, Singh J, Nisar KS. Certain unified integrals associated with product of M-series and incomplete H-functions. *Mathematics*. 2019;7(12):1191.
- Nisar KS, Rahman G, Tomovski Z. On a certain extension of the Riemann–Liouville fractional derivative operator. *Commun Korean Math Soc*. 2019;34(2):507–522.
- Khan O, Khan N, Nisar KS, Saif M, Baleanu D. Fractional calculus of a product of multivariable Srivastava polynomial and multi-index Bessel function in the kernel F_3 . *AIMS Mathematics*. 2020;5(2):1462–1475.
- Nisar KS, Baleanu D, Qurashi MMA. Fractional calculus and application of generalized Struve function. *Springer Plus*. 2016;5:1–13.
- Abubakar UM. A study of extended beta and associated functions connected to Fox-Wright function. *J Fract Calc Appl*. 2021;12(13):1–23.
- Ata E, Kıymaz İO. A study on certain properties of generalized special functions defined by Fox-Wright function. *Appl Math Nonlinear Sci*. 2020;5(1):147–162.
- Ata E, Kıymaz İO. Generalized gamma, beta and hypergeometric functions defined by Wright function and applications to fractional differential equations. *Cumhuriyet Sci J*. 2022;43(4):684–695.
- Ata E. Modified special functions defined by generalized M-series and their properties. 2022 arXiv preprint <https://arxiv.org/abs/2201.00867v1>[math.CA].
- Ata E, Kıymaz İO. New generalized special functions with two generalized M-series at their kernels and solution of fractional PDEs via double Laplace transform. *Comput Methods Differ Equ*. 2024;12(1):31–43.
- Ata E, Kıymaz İO. On the generating functions of the newly defined generalized hypergeometric functions. *TWMS J Appl Eng Math*. 2024;14(1):64–78.
- Chaudhry MA, Zubair SM. Generalized incomplete gamma functions with applications. *J Comput Appl Math*. 1994;55(1):99–124.
- Chaudhry MA, Qadir A, Rafique M, Zubair SM. Extension of Euler's beta function. *J Comput Appl Math*. 1997;78(1):19–32.
- Chaudhry MA, Qadir A, Srivastava HM, Paris RB. Extended hypergeometric and confluent hypergeometric functions. *Appl Math Comput*. 2004;159(2):589–602.
- Choi J, Rathie AK, Parmar RK. Extension of extended beta, hypergeometric and confluent hypergeometric functions. *Honam Math J*. 2014;36(2):357–385.
- Çetinkaya A, Kıymaz İO, Agarwal P, Agarwal R. A comparative study on generating function relations for generalized hypergeometric functions via generalized fractional operators. *Adv Differential Equations*. 2018;2018(156):1–11.
- Goswami A, Jain S, Agarwal P, Araci S. A note on the new extended beta and Gauss hypergeometric functions. *Appl Math Inf Sci*. 2018;12(1):139–144.
- Kulip MAH, Mohsen FBF, Barahmah SS. Further extended gamma and beta functions in term of generalized wright function. *Elec J Uni of Aden for Basic and Appl Sci*. 2020;1(2):78–83.
- Lee DM, Rathie AK, Parmar RK, Kim YS. Generalization of extended beta function, hypergeometric and confluent hypergeometric functions. *Honam Math J*. 2011;33(2):187–206.

27. Mubeen S, Rahman G, Nisar KS, Choi J. An extended beta function and its properties. *Far East J Math Sci.* 2017;102(7):1545–1557.
28. Özergin E, Özarslan MA, Altun A. Extension of gamma, beta and hypergeometric functions. *J Comput Appl Math.* 2011;235(16):4601–4610.
29. Parmar RK. A new generalization of gamma, beta, hypergeometric and confluent hypergeometric functions. *Matematiche (Catania).* 2013;68(2):33–52.
30. Rahman G, Mubeen S, Nisar KS. A new generalization of extended beta and hypergeometric functions. *J Fract Calc Appl.* 2020;11(2):32–44.
31. Shadab M, Saime J, Choi J. An extended beta function and its applications. *Far East J Math Sci.* 2018;103(1):235–251.
32. Srivastava HM, Agarwal P, Jain S. Generating functions for the generalized Gauss hypergeometric functions. *Appl Math Comput.* 2014;247:348–352.
33. Şahin R, Yağcı O, Yağbasan MB, Kıymaz İO, Çetinkaya A. Further generalizations of gamma, beta and related functions. *J Inequal Spec Funct.* 2018;9(4):1–7.
34. Sharma M, Jain R. A note on a generalized M-series as a special function of fractional calculus. *Fract Calc Appl Anal.* 2009;12(4):449–452.
35. Andrews GE, Askey R, Roy R. *Special functions.* Cambridge: Cambridge University Press; 1999.