

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/332189952>

Three new copper (II) complexes with chiral Schiff bases: Synthesis, characterization, DNA binding and DNA-cleavage studies

Article in *Journal- Chemical Society of Pakistan* · January 2019

CITATIONS

0

READS

186

6 authors, including:



Turgay Tunc

Ahi Evran Üniversitesi

16 PUBLICATIONS 66 CITATIONS

SEE PROFILE



Nadir Demirel

Ahi Evran Üniversitesi

28 PUBLICATIONS 463 CITATIONS

SEE PROFILE



Aslihan Gunel

Ahi Evran Üniversitesi

16 PUBLICATIONS 105 CITATIONS

SEE PROFILE



Nurcan Karacan

Gazi University

58 PUBLICATIONS 1,081 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Ahi Evran University Scientific Research Foundation under grant No FEF.A3.16.015 [View project](#)



Two-component organogelators: combination of N ϵ -alkanoyl-L-lysine with various N-alkanoyl-L-amino acids: additional level of hierarchical control [View project](#)

Three New Copper (II) Complexes with CHIRAL SCHIFF BASES: Synthesis, Characterization, DNA Binding and DNA-Cleavage Studies

¹Turgay Tunç*, ²Nadir Demirel, ²Mahmut Emir, ²Aslıhan Günel, ²Rıfki Kadioğlu, ³Nurcan Karacan

¹Department of Chemistry Engineering and Process, Faculty of Engineering, University of Ahi Evran, Kırsehir.


²Department of Chemistry, Faculty of Science & Art, University of Ahi Evran, Kırsehir 40100, Turkey.

³Department of Chemistry, Faculty of Science, University of Gazi, Ankara, 06500, Turkey.

ttunc@ahievran.edu.tr*

(Received on 25th January 2018, accepted in revised form 26th September 2018)

Summary: New mononuclear copper (II) complexes (1, 2 and 3) were synthesized from Schiff bases (H₂L) of chiral amino alcohols. The structures of the copper complexes were proposed by a combination of elemental analyses, FTIR, LCMS, magnetic susceptibility and molar conductance measurement methods. Spectroscopic and analytical data of the complexes suggest four-coordinated structures. Geometry optimization carried out with DFT/6-31G (d,p) were proposed to be distorted square planar geometry for the complexes. The similarity between experimental and theoretical IR spectra confirms the proposed structures. The interaction of copper (II) complexes with calf thymus (CT-DNA) was investigated using absorption titration method. The results suggest that the complex **1** and **2** can bind to DNA by intercalation. Binding constants K_b were found to be 2.46×10^5 for **1**, 5.41×10^5 for **2** and 7.00×10^4 for **3**. Moreover, agarose gel electrophoresis assay demonstrates that all complexes were found to cleavage of plasmid penty/d-topo plasmid DNA. Complex **2** shows the best cleavage activity (5 μ M).



Keywords: Schiff base complexes, DNA binding, DNA cleavage, DFT, Copper (II) complexes.

Introduction

Although metal ions have either boon or bane for living organisms [1]. They are crucial for cellular activity thanks to play role as co-factor. Metal ion deficiency or overloading of these ions can cause pathological conditions such as anemia because of iron deficiency and Wilson's disease thanks to copper overloading in the liver. Treatments of this situation are required same strategies. If someone has Wilson's syndrome S/he can treat with chelating agent to remove poisoned metal ion [2]. On the other hand, metallodrugs are not only therapeutics but also they are used for diagnostic purpose and even they have been used as theranostic agents recently [3].

The second review by G. Pappalardo and co-authors is focused on the re-purposing of copper-chelating compounds for treating neurodegenerative disorders, such as Alzheimer's (AD) and Parkinson's (PD). Together with zinc and iron, copper plays a key role in neuronal functions and its levels are finely regulated to avoid the onset of various age-related

diseases. Their vocation is to trigger cellular effects by redox and/or by coordinative processes. Several complexes with different metals such as Pt, As, Au, Co, Tc, Gd, are approved by FDA and they are used as therapeutic, anticancer, antisiphillis, antiarthritic, antimicrobial and diagnostic agent, in single photon emission computed tomography respectively [4].

Copper, an endogenous element for several aerobic organisms, is less toxic to normal cells than cancer cells. For this reason, copper complexes possess anticancer activities [5-7]. In the past decades, Schiff bases copper complexes have attracted much attention in the improvement of novel therapeutic agents because they have a various pharmacological activity such as anticancer, antifungal, anti-inflammatory antioxidation, antibacterial, antimalarial, DNA nuclease activity and so forth [8-12]. Beside anticancer activity of metallodrugs containing copper as coordinated ion. They are attracted considerable attention in view of neurological diseases

*To whom all correspondence should be addressed.

such as ALS, Parkinson's Disease [13] and epilepsy [14] in recent years.

Chiral molecules exhibit different chemical and physical properties in the chiral environment. Therefore, chiral biological active compounds for instance pharmaceuticals, agrochemicals; nutrients are sold as single enantiomers [15]. Enantioselectivity is the important criterion for metal complex-DNA interactions and drug design because the best fitting complexes for DNA helical structure exhibit the highest binding affinity [16]. On the other hand, chiral ligands increase the chirality [17] of the complexes and improve the pharmacological behavior of metal complexes. Therefore, there is a great demand for enantiomeric metallodrugs.

Although a large number of research on the chiral metal complexes have been published so far [18-28] few studies have been carried out on chiral copper(II) complexes [29-36] and their DNA binding and biological properties [37-39]. In this regard, we have synthesized and characterized three new copper(II) complexes (**1-3**) bearing chiral Schiff bases of chiral amino alcohols. The CT-DNA binding and supercoiled penty/d-topo plasmid DNA cleavage activities of copper complexes were also evaluated.

Experimental


Materials and Physical Measurements

All solvents and reagents were purchased from Sigma-Aldrich and used without further purification. IR spectra were recorded on a Nicolet-6700 ATR-FT-IR spectrophotometer in the 4000 – 400 cm^{-1} region. UV-Vis spectra were recorded on Thermo Scientific Genesys 10S spectrophotometer. Elemental analyses for C, H and N were carried out with an LECO CHNS-932 auto elemental analyzer. Melting points were measured using a Thermo Fisher Scientific Electrothermal 9100 apparatus. ^1H (400 MHz) and ^{13}C (100 MHz) NMR spectra were recorded on a Bruker DPX-400 high performance digital FT-NMR spectrometer. Mass spectra were recorded with Thermo Scientific TSQ Quantum Access Max LC-MS spectrometers in methanol/acetonitrile mixture.

Synthesis of the compounds

Chiral Schiff bases L1 (1,4-Bis(phenoxy) ethane, (4,4'-(E,E)-(R)-hydroxy-2-phenylethylimino) methyl)), L2 (1,4-Bis-(4,4'-(E,E)-(R)-hydroxy-2-phenylethylimino) methyl) benzene), L3 (1,4-Bis-(4,4'

(E,E)-(1S,2R)-hydroxy-1,2-diphenylethylimino) methyl) benzene) were synthesized according to our previous reported procedure [40]. The Cu(II) complexes (**1**, **2** and **3**) were synthesized using the following general procedure (Scheme 1). A methanolic solution (10 mL) of the ligand (0.064 mmol) was added dropwise to methanolic solution (10 mL) of a copper (II) acetate (0.064 mmol, 11.7 mg). 2-3 drop of the trimethylamine was added to the mixture, then, refluxed for six hours at 60 °C. The solution was allowed to cool and waited for 2 days at room temperature. The green solid was collected by vacuum filtration and washed twice with 3–5 mL aliquots of ice-cold methanol and dried under vacuum.



Scheme-1: Synthesis route of the copper (II) complexes.

Synthesis of $[\text{CuL}^1]$ (**1**)

Yield: 0.031 g (88.9%), M.p.: 104-106 °C. LC-MS (m/z): 615.06 [M+H] (calc. 614.16). FTIR data (neat, v/cm^{-1}): 3034, 2896, 1680, 1599, 1570, 1505, 1424, 1395, 1309, 1251, 1145, 1047, 951, 921, 837, 802, 761. Anal. calcd. for $\text{C}_{34}\text{H}_{34}\text{CuN}_2\text{O}_5$: C, 66.49; H, 5.58; N, 4.56. Found: C, 65.97; H, 5.42; N, 4.75. Λ_m° : 12.11 $\text{ohm}^{-1}\text{cm}^2\text{mol}^{-1}$. μ_{eff} : 1.92 BM.

Synthesis of $[\text{CuL}^2]$ (**2**)

Yield: 0.036 g (82.9%), M.p.: 122-124 °C. LC-MS (m/z): 647.02 [M+H] (calc. 646.25). FTIR data

(neat, ν/cm^{-1}): 3086, 2938, 2874, 1678, 1600, 1570, 1504, 1420, 1381, 1305, 1243, 1158, 1103, 1048, 997, 886, 827, 796. Anal. calcd. for $\text{C}_{38}\text{H}_{34}\text{CuN}_2\text{O}_4$: C, 70.63; H, 5.30; N, 4.33. Found: C, 71.04; H, 5.17; N, 4.26. Λ_m° : $11.25 \text{ ohm}^{-1}\text{cm}^2\text{mol}^{-1}$. μ_{eff} : 1.85 BM.

Synthesis of $[\text{CuL}^3]$ (**3**)

Yield: 0.042 g (79.4%), M.p.: 123-125 °C. LC-MS (m/z): 799.02 [M+H] (calc. 798.44). FTIR data (neat, ν/cm^{-1}): 3102, 3025, 2966, 2944, 1741, 1678, 1599, 1569, 1503, 1421, 1374, 1306, 1223, 1159, 1040, 999, 904, 827. Anal. calcd. for $\text{C}_{50}\text{H}_{42}\text{CuN}_2\text{O}_4$: C, 75.22; H, 5.30; N, 3.51. Found: C, 75.11; H, 5.62; N, 3.39. Λ_m° : $14.45 \text{ ohm}^{-1}\text{cm}^2\text{mol}^{-1}$. μ_{eff} : 1.83 BM.

Computational method

All density functional theory (DFT) calculations were made using Gaussian 03 program package using the standard B3LYP functional [41]. For the geometric optimization of the copper (II) complexes (multiplicity =2, charge = 0) initial geometry was obtained by semi-empirical PM6 method. Then, obtained structures were reoptimized using B3LYP/6-31G (d,p) basis set for all atoms in the gas phase. The vibrational frequency calculations were also performed using B3LYP/6-31G(d,p) level. All the frequencies are found to be positive for the structures, indicating them to be a local minima. The choice of B3LYP/6-31G(d,p) level is based on its good performance found in earlier computational studies on copper(II) complexes.

DNA binding experiment

CT-DNA binding studies was conducted in Tris buffer (5 mM, pH 7.1). UV-Vis absorbance at 260 and 280 nm of the solution of CT-DNA in the buffer about 1.9:1 indicating that the DNA was free from protein [42]. The amount of DNA concentration was calculated by absorption spectroscopy using the molar extinction coefficient ($6600 \text{ M}^{-1} \text{ cm}^{-1}$) at 260 nm [43]. The intrinsic binding constant K_b for the interaction of our Cu (II) complexes with CT-DNA was calculated based on the absorption spectral changes after the addition of increasing DNA. The binding constant was calculated according to the following equation [44-45].

$$[\text{DNA}] / (\epsilon_a - \epsilon_f) = [\text{DNA}] / (\epsilon_b - \epsilon_f) + 1/K_b (\epsilon_b - \epsilon_f)$$

$[\text{DNA}]$ represent the concentration of DNA in base pairs; the apparent absorption coefficient ϵ_a , ϵ_f and ϵ_b correspond to $A_{\text{obs}}/[\text{M}]$, the extinction coefficient of the free and the extinction coefficient of the compound when fully bound to DNA, respectively. The plot of $[\text{DNA}] / (\epsilon_a - \epsilon_f)$ vs $[\text{DNA}]$ gave a straight line with a

slope of $1/(\epsilon_b - \epsilon_f)$ and an intercept of $1/K_b(\epsilon_b - \epsilon_f)$; K_b was determined from the ratio of the slope to intercept. UV-vis titration experiments were performed by keeping the concentration of complexes constant and changing the nucleic acid/nucleotide concentration. The absorbance (A) was recorded after consecutive additions of CT-DNA. CT-DNA was added to both the compound solution and the reference solution to avoid the absorbance of the CT-DNA itself.

DNA cleavage experiment

The DNA cleavage activity of our complexes was examined using supercoiled pentry/d-topo plasmid DNA. Plasmid DNA was purified by using Perfect Prep Endo Free Plasmid Maxi Kit from Qiagen according to manufacturer instructions [46]. After purification, plasmid DNA amount was measured at 260 nm by nanodrop spectrophotometer. Pure DNA preparation has a ratio of $A_{260}/A_{280} > 1.8$ and $A_{260}/A_{230} = 1.80$. A stock solution (500 μM) of the complexes was prepared. Plasmid DNA was treated with increasing complex concentrations (20-150 μM) and enough buffer solution for a final volume of 50 μL (22 μL plasmid DNA + required volume of complexes to achieved desired concentration and volume are completed to 50 μL). The samples were incubated in the dark for 12 h at 37 °C. The DNA cleavage experiments were analyzed by 1% agarose gel electrophoresis using 1XTris-Acetate-EDTA (TAE) running buffer. 20 μL of each reaction solution together with 10 μL of the loading dye was loaded into the wells and subjected to electrophoresis at 75 V for 90 min. The gel was stained with ethidium bromide and documented using a Gene Tools imaging system from Syngene.

Results and Discussion

Structure of the copper (II) complexes

The molar conductivities of the complexes dissolved in DMF ($1.00 \times 10^{-3} \text{ M}$) were measured at room temperature. The molar conductance values (Λ_m°) were found to be 12.11 for **1**, 11.25 for **2** and 14.45 for **3**. Obtained low conductivity data indicate that the copper(II) complexes are non-electrolytes [47].

The magnetic moments of the complexes were found to be 1.92 for **1**, 1.85 for **2** and 1.83 BM for **3** at room temperature. These values are quite close to the values expected for spin-only monomeric copper(II) complexes and are characteristic of $3d^9$ electronic configuration.

ESI-MS spectra of the copper(II) complexes were depicted in Fig 1. The spectra recorded in the

positive mode of the complexes (**1-3**) were exhibited indicating the presence of $[M+H]^+$ ions for **1,2** and **3**. These molecular ions confirm the stoichiometry of the copper(II) complexes as $[CuL]$ type. Briefly, analytical data of the complexes were in agreement with a 1:1 (Cu:L) molar ratio.

Synthesis and single crystal X-ray structure of several copper (II) complexes bearing Schiff base prepared from different chiral amino alcohols have been reported in the literature [48-51]. In these complexes, Schiff bases coordinate to copper(II) atom with imine nitrogen atom and unde protonated or deprotonated alkoxy-oxygen atom. It is known that different copper salts give rise to different geometry

[52]. In this study, copper(II) acetate and trimethylamine were used in the synthesis, and deprotonation occurred. 1H -NMR spectrum of the complex **1**, not given here due to poor resolution, confirm the deprotonation. In addition, disappearing of ν_{OH} stretching vibration (Fig. 2), a broad band between $3500 - 3550\text{ cm}^{-1}$ of the Schiff bases with complex formation also confirm the deprotonation. On the basis of these findings, four- coordinated copper(II) complexes bearing four-dentate Schiff bases ligands having two imine nitrogen atoms and two deprotonated alkoxy-oxygen atoms were proposed for (**1-3**).




Fig. 1: LC-MS spectra of the copper (II) complexes.

DFT calculation of the copper (II) complexes

The geometry optimization of the proposed complexes (**1-3**) was performed at B3LYP/3-21G(d) level for different starting geometries, and several local minima were found. Then, two low-energy conformers were selected and optimized with B3LYP/6-31G(d,p) level. Their optimized structures are shown in Fig. 3. Calculated electronic energies, some selected bond angles, and bond length were given in Table-1. The calculated bond length of Cu-N and Cu-O were in agreement with the single crystal X-ray data obtained from similar copper(II) complexes reported in the literature [47-52]. Initial geometries of (a) and (b) were square planar and

tetrahedral structures, respectively. However, after optimization, both got the distorted square planar geometry. As seen from Table 1, square planar (a) structures have lower energy than tetrahedral (b) structures for three complexes (**1-3**). O-Cu-O and N-Cu-N bond angles of (a) are wider than the corresponding (b). If the octahedral structure was taken as starting geometry with additional two phenolic oxygen atoms of Schiff base, after optimization, it returns to the *pseudo* square planar structure. In this structure, the Cu-O distance was calculated as 6.5 Å, indicating that there is no bond between Cu and phenolic oxygen atoms.




Table-1: Total electronic energies (Hartrees) and selected bond angles ($^{\circ}$) and bond length (\AA) for two structures of the copper(II) complexes optimized via DFT/B3LYP/6-31G(d,p)

	$E_{\text{electronic}}$ (a.u)	O-Cu-O	N-Cu-N	Cu-N	Cu-O
1a	-3442.0434	152.65	161.22	2.01	1.88
1b	-3442.0315	115.00	122.18	1.99	1.85
2a	-3519.2707	158.13	170.06	2.01	1.87
2b	-3519.2695	153.58	169.97	2.01	1.87
3a	-3981.3856	159.65	173.16	2.02	1.88
3b	-3981.3802	153.03	170.67	2.02	1.89




Fig. 3: Optimized structures of square planar (a) and tetrahedral (b) the copper(II) complexes calculated by B3LYP/6-31G(d,p).

Harmonic vibrational frequencies of ground state optimized geometries (a) were calculated using the DFT/B3LYP /6-31G(d,p). Calculated IR spectra (Table-2) showed good agreement with the experimental IR spectra (Fig. 2). The differences were related to the packing effects of the solid state structure, with respect to the isolated complexes at the gas phase. The second reason is that calculated values are harmonic frequencies while the experimental values contain anharmonic vibrational frequencies. However, the spectral patterns are quite similar to each other, confirming the proposed structures. It is well known that the calculated

wavenumbers are higher than their experimental equivalents, for this reason, empirical scaling factors are used to obtain considerably better agreement with the experimental data. We have used the scaling factors of 0.98-0.96. Afterward, we compared and assigned the vibrational modes by matching peak intensities and using scaled peak frequencies. Experimental and calculated wave numbers of selected vibrational modes (for intensity higher than 40 km/mol) of the complexes were given in Table-2.

Table-2: Experimental and calculated wavenumbers of selected vibrational modes of the complexes.

1		2		3		Assignments*
exp	calc	exp	calc	exp	calc	
-	3202	-	3199	3028	3204	CH(ar)
-	3199	-	3173	-	3198	CH(ar)
-	3147	-	3138	-	3151	CH(ar)
2935	3056	2943	3052	2945	3053	CH ₂
2836	2970	2825	2900	2826	2947	CH ₂
2752	2907	2734	2899	2734	2847	CH ₂
1686	1716	1678	1718	1678	1714	$\nu_{\text{C-N}}$
1602	1661	1604	1660	1604	1665	ring
1575	1659	1576	1659	1576	1660	ring
1509	1657	1507	1656	1507	1658	ring
1426	1496	1426		1426	1460	CH(ar)
1399	1470	1392		1392	1395	CH(ar)+ CH ₂
1359	1365	1378		1379	1361	CH(ar)+ CH ₂
1257	1286	1245	1286	1245	1287	CH ₂ +CH(ar)
1173	1207	1162	1202	1162	1202	CH(ar) bridge
1111	1130	1111	1132	1111	1132	$\nu_{\text{C-O(alkoxy)}}$ +CH(ar)
1078	1109	1080	1098	1080	1097	$\nu_{\text{C-O(alkoxy)}}$ + CH(ar)
1050	1082	1047	1062	1045	1060	$\nu_{\text{C-O(phenolic)}}$
-	-	1028	1044	1027	1043	CH(ar)
-	-	1000	1013	999	1012	CH(ar)+ CH ₂
954	964	-	-	-	-	C-N
924	934	891	902	891	902	CH ₂ (aliphatic)+
841	850	829	846	829	845	CH ₂ (aliphatic)
801	816	803	829	803	828	CH (ar-bridge)
769	788	797	819	797	819	CH (ar-bridge)
700	717	769	783	770	783	CH (ar-terminal)
654	662	647	646	647	659	ring(ar)
620	653	617	631	617	632	ring(ar)
514	554	552	574	552	574	$\nu_{\text{asCu-O}}$
-	-	508	522	508	522	skeleton

* ν : stretching, δ : in-plane bending, γ : out-of-plane bending

DNA Binding Studies

The absorption spectra of the titration of the copper(II) complexes (20 μM) with CT-DNA from 0 to 100 μM are shown in Fig. 4. Absorption band in the range 250 -300 nm (λ_{max} is 280 nm) correspond to $\pi \rightarrow \pi^*$ transition of the aromatic rings of the copper(II) complexes. As seen in Fig.4, the band height at 280 nm gradually decreases for 1 and 2, but increases for 3. On titration with CT-DNA, complex 1 exhibits slight hypochromic and narrow bathochromic effects; whereas, complex 2 displays significant hypochromic and large bathochromic effects; moreover, complex 3 shows slight hyperchromic and narrow hypsochromic effects. Hypochromism is caused by parallel stacking of base pairs in a double helix and refer to the presence of intercalation or at last partial intercalation for 2 and

1. The extent of the hypochromic is usually proportional to the strength of the interaction [53]. Whereas, hyperchromism is an indication of the reduction in the amount of parallel stacking in double helix resulting from 1,2-intrastrand crosslinking. The extent of the hyperchromism is indicative of the partial or non-intercalative binding modes [54].




Fig. 4: Absorption spectra of the copper(II) complexes (20 μ M) in buffer solution (5 mM Tris/ 50 mM NaCl, pH 7.5) upon addition of CT-DNA from 0 to 100 μ M. Arrow indicates the changes in absorbance upon increasing the DNA concentration. Inset: Plots of $-\frac{[DNA]}{[DNA]}$ vs. $[DNA]$ for the titration of CT-DNA. (1 and 3 cyan represents 0 M $[DNA]$ and 2 orange represents 0 M $[DNA]$). Directions of Arrows indicate increasing DNA concentration.)

Cis-platin and its analogs were reported to have hyperchromic and bathochromic shifts [55]. In order to compare the DNA-binding strengths of these complexes, the intrinsic binding constant K_b was determined from the changes in absorbance. The values of K_b were 2.46×10^5 (1), 5.41×10^5 (2) and 7.00×10^4 (3), respectively (Table-3). It was observed that the K_b value increased when the aromatic moiety was replaced by the aliphatic ether group on the bridge at 2. It was expected due to the strong stacking interaction between the aromatic chromophore and the base pairs of DNA. On the contrary, K_b value decreased when the number of phenyl substituent increased at 3. Much increase in molecular volume caused an opposite effect. The copper(II) complexes having $[Cu_8]^{8+}$ and $[Cu_4]^{8+}$ units of the chiral alkanol amine Schiff base ligands were reported to exhibit hyperchromism and low K_b values as 3.42×10^3 and 3.04×10^4 , respectively [49].

Table-3: Absorption Spectroscopic properties of the complexes binding to CT DNA.

Complexes	Change in absorbance	Shifting	Shifting $\Delta\lambda$	K_b/M^{-1}
1	Hypochromism	Bathochromic	+5	2.46×10^5
2	Hypochromism	Bathochromic	+25	5.41×10^5
3	Hyperchromism	Hypsochromic	5	7.00×10^4




Fig. 5: Gel electrophoresis pattern showing cleavage of plasmid DNA (300 ng) treated with complex 1 in a Tris buffer (5 mM Tris/ 50 mM NaCl, pH 7,5) at 37 °C with an incubation time of 12 h. Lane1: Control DNA; Lane 2: DNA + 1 (5 μ M), Lane 3: DNA+ 1 (10 μ M), Lane 4: DNA+ 1 (25 μ M), Lane 5: DNA+ 1 (50 μ M), Lane 6:DNA+1 (100 μ M), Lane 7: DNA+1 (150 μ M), Lane 8: DNA+1 (200 μ M) (b) Comparison profiles of the lanes.

Plasmid DNA cleavage

The DNA cleavage activity of the complexes for supercoiled penty/d-topo plasmid DNA was investigated by gel electrophoresis. When supercoiled DNA is subject to electrophoresis, relative migration will be observed for the intact supercoiled form (Form I). If cutting occurs on one strand (nicking), the supercoiled form will relax and generate a slower-moving singly nicked form (Form II). If both strands are cleaved, a double nicked linear form (Form III) that migrates between Form I and Form II will be generated. Plasmid DNA electropherogram traces of the complexes (1-3) with increasing concentration (5-200 μM) were given in Fig 5-7. DNA cleavage activity of the complexes is not concentration dependent. While the percentage conversion to Form I to Form II is mostly at line 2 (10 μM), the maximum conversion was observed at 150 μM (line 7) for complex 1 (Fig 5). Complex 2 exhibited the best cleavage at 5 μM in Fig. 6 (line 2), and the complex 3 shows the best conversion to Form II at 10 μM (Fig. 7, line 3). These results do not correspond to their binding capabilities.




Fig. 6: Gel electrophoresis pattern showing cleavage of plasmid DNA (300 ng) treated with complex 2 in a Tris buffer (5 mM Tris/ 50 mM NaCl, pH 7,5) at 37 °C with an incubation time of 12 h. Lane1: Control DNA; Lane 2: DNA + 2 (5 μM), Lane 3: DNA+ 2 (10 μM), Lane 4: DNA+ 2 (25 μM), Lane 5: DNA+ 2 (50 μM), Lane 6:DNA+2 (100 μM), Lane 7: DNA+2 (150 μM), Lane 8: DNA+2 (200 μM) (b) Comparison profiles of the lanes.




Fig. 7: Gel electrophoresis pattern showing cleavage of plasmid DNA (300 ng) treated with complex 3 in a Tris buffer (5 mM Tris/ 50 mM NaCl, pH 7,5) at 37 °C with an incubation time of 12 h. Lane1: Control DNA; Lane 2: DNA + 3 (5 μM), Lane 3: DNA+ 3 (10 μM), Lane 4: DNA+ 3 (25 μM), Lane 5: DNA+ 3 (50 μM), Lane 6:DNA+3 (100 μM), Lane 7: DNA+3 (150 μM), Lane 8: DNA+3 (200 μM) (b) Comparison profiles of the lanes.

Conclusion

Three new mononuclear copper(II) complexes possessing ONNO type Schiff bases of chiral amino alcohol were synthesized. On the basis of analytical and experimental data, the Schiff bases were coordinated to Cu(II) ion with four donor atoms, two imine nitrogens and two deprotonated alkoxy-oxygen atoms. Deprotonation is achieved by the addition of trimethylamine to the reaction mixture. Geometry optimization with B3LYP/6-31G(d,p) in the gas phase was proposed to be a distorted square planar geometry of the complexes. The similarity between experimental IR spectra and theoretical IR spectra obtained from calculated harmonic vibrational frequencies of the complexes confirm the proposed structures. The CT-DNA binding mode of these complexes has been evaluated using absorption titration method. Intercalation was proposed for complex 1 and 2, which exhibit hypochromic and bathochromic effects. Whereas, complex 3 displays slight hyperchromic effect and hypsochromic shift. The binding constants K_b of the complexes were determined in the following order 2 > 1 > 3. The number of the phenyl group play an

important role in the DNA binding values. K_b value increased when the aromatic moiety was replaced by the aliphatic ether group on the bridge at **2**. However, K_b value decreased when two more phenyl groups join the structure of **3**. Much increase in molecular volume caused an opposite effect. The concentration-dependent DNA cleavage experiments using plasmid pentry/d-topo plasmid DNA were also performed. Complex **2** shows the best DNA cleavage activity (5 μ M). We consider that there is no correlation between the cleavage activities and DNA binding capabilities. We tested the correlation between DNA cleavage and binding properties of compounds by Pearson Analysis by Excell and we concluded that no correlation is present between these two features with 95% confidence ($p < 0.05$).

Acknowledgments

This study was supported by Ahi Evran University Scientific Researchs Foundation under grant No FEF.A3.16.015

References

1. K. H. Thompson, C. Orvig, Boon and Bane of Metal Ions in Medicine, *Science*, **300**, 936 (2003).
2. I. Subramanian, Z. F. Vanek, J. M. Bronstein, Diagnosis and treatment of Wilson's disease. *Curr. Neurol. Neurosci. Rep.* **2**, 317 (2002).
3. H. D. Lu, L. Z. Wang, B. K. Wilson, S. A. McManus, J. Jumai'an, P. K. Padakanti, A. Alavi, R. H. Mach, and R. K. Prud'homme, Copper Loading of Preformed Nanoparticles for PET-Imaging Applications, *ACS Applied Materials & Interfaces*, **10**, 3191 (2018).
4. K. D. Mjos and C. Orvig. Metallodrugs in Medicinal Inorganic Chemistry. *Chem. Rev.* **114**, 4540 (2014)
5. E. I. Solomon, D. E. Heppner, E. M. Johnston, J. W. Ginsbach, J. Cirera, M. Qayyum, M. T. Kieber-Emmons, C. H. Kjaergaard, R. G. Hadt, L. Tian, Copper Active Sites In Biology, *Chem. Rev.*, **114**, 3659 (2014).
6. C. Santini, M. Pellei, V. Gandin, M. Porchia, F. Tisato, C. Marzano, Advances In Copper Complexes as Anticancer Agents, *Chem. Rev.*, **114**, 815 (2014).
7. F. Tisato, C. Marzano, M. Porchia, M. Pellei and C. Santini, Copper In Diseases and Treatments, and Copper-Based Anticancer Strategies, *Med. Res. Rev.*, **30**, 708 (2010).
8. H. Yu, Y. Yang, Q. Li, T. Ma, J. Xu, T. Zhu, J. Xie, W. Zhu, Z. Cao, K. Dong, J. Huang, L. Jia, Ternary Dinuclear Copper(II) Complexes of a Reduced Schiff Base Ligand with Diimine Coligands: DNA Binding, Cytotoxic Cell Apoptosis, and Apoptotic Mechanism, *Chem. Biol. Drug Des.*, **87**, 398 (2016).
9. A. M. Abu-Dief, I. M. A. Mohamed, a Review of Versatile Applications of Transition Metal Complexes Incorporating Schiff Bases, *Beni-Suef Univ. J. Basic Appl. Sci.*, **4**, 119 (2015).
10. A. Chakraborty, P. Kumar, K. Ghosh and P. Roy, Evaluation of a Schiff Base Copper Complex Compound as Potent Anticancer Molecule with Multiple Targets of Action, *Eur. J. Pharmacol.*, **647**, 1 (2010).
11. J. Lu, Q. Sun, J. L. Li, L. Jiang, W. Gu, X. Liu, J. L. Tian, S.P. Yan, Two Water-Soluble Copper(II) Complexes: Synthesis, Characterization, DNA Cleavage, Protein Binding Activities and in Vitro Anticancer Activity Studies, *J Inorg Biochem.*, **137**, 46 (2014).
12. M. Sarigul, P. Deveci, M. Kose, U. Arslan, H. Türk Dagi, M. Kurtoglu, New Tridentate Azo Azomethines, and Their Copper(II) Complexes: Synthesis, Solvent Effect on Tautomerism, Electrochemical and Biological Studies, *J. Mol. Struct.*, **1096**, 64 (2015).
13. V. Lanza, D. Milardi, G. Di Natale and G. Pappalardo, Repurposing of Copper(II)-chelating Drugs for the Treatment of Neurodegenerative Diseases, *Curr. Med. Chem.*, **25**, 525 (2018).
14. L. D. Grünspan, B. H. M. Mussulini, S. Baggio, P. R. dos Santos, F. Dumas, E. P. Rico, D. L. de Oliveira, S. Moura., Teratogenic and Anticonvulsant Effects of Zinc and Copper Valproate Complexes in Zebrafish. *Epilepsy Research*, **139**, 171 (2018).
15. C. Wolf, Dynamic Stereochemistry of Chiral Compounds. Principles and Applications, *Rsc Publishing*, (2007).
16. J. G. Liu, B. H. Ye, Q. L. Zhang, X. H. Zou, Q. X. Zhen, X. Tian and L. N. Ji, Enantiomeric Ruthenium(II) Complexes Binding to DNA: Binding Modes And Enantioselectivity, *J. Biol. Inorg. Chem.*, **5**, 119 (2000).
17. J. Crassous, Chiral Transfer in Coordination Complexes: Towards Molecular Materials, *Chem. Soc. Rev.*, **38**, 830 (2009).
18. G. Muller, Luminescent Chiral Lanthanide(III) Complexes as Potential Molecular Probes, *Dalt. Trans.*, **42**, 9692 (2009).
19. B. L. Fei, W. Xu, W. Gao, J. Zhang, Y. Zhao, J. Long, C. E. Anson, A. K. Powell, Biology DNA Binding and Cytotoxicity Activity of a Chiral Iron (III) Triangle Complex Based on a Natural Rosin Product, *J. Photochem. Photobiol. B Biol.*, **142**, 77 (2015).

20. Z. Dezhahang, M. R. Poopari, J. Cheramy, Y. Xu, Conservation Of Helicity In A Chiral Pyrrol-2-Yl Schiff-Base Ligand And Its Transition Metal Complexes, *Inorg. Chem.*, **54**, 4539 (2015).
21. N. H. Khan, N. Pandya, N. C. Maity, M. Kumar, R. M. Patel, R. I. Kureshy, S. H. R. H. Abdi, S. Mishra, S. Das, H. C. Bajaj, Influence of Chirality of V(V) Schiff Base Complexes on DNA, BSA Binding and Cleavage Activity, *Eur. J. Med. Chem.*, **46**, 5074 (2011).
22. R. Alul, M. B. Cleaver, J. S. Taylor 'II, Synthesis and Characterization of a Series of Achiral and Chiral Mono-and Bis[(Dien)Pt(II)]I Derivatives as Potential DNA and RNA Structure Probes and Anticancer Drugst, *Inorg. Chem.*, **31**, 3636 (1992).
23. F. Arjmand, M. Muddassir, I. Yousuf, Design and Synthesis of Enantiomeric (R)- and (S)-Copper(II) and Diorganotin(IV)-Based Antitumor Agents: Their in Vitro DNA Binding Profile, Cleavage Efficiency and Cytotoxicity Studies, *J. Photochem. Photobiol. B Biol.*, **136**, 62 (2014).
24. Z. Chu, W. Huang, L. Wang, S. Gou, Chiral 27-Membered [3+3] Schiff-Base Macrocycles and Their Reactivity with First-Row Transition Metal Ions, *Polyhedron.*, **27**, 1079 (2008).
25. A. Dhamija, S. A. Ikbal and S. P. Rath, Induction and Rationalization of Supramolecular Chirality in The Tweezer-Diamine Complexes: Insights from Experimental and DFT Studies, *Inorg. Chem.*, **55**, 13014 (2016).
26. J. Rich, M. Rodríguez, I. Romero, L. Vaquer, X. Sala, A. Llobet, M. Corbella, M. N. Collomb, X. Fontrodona, Mn(II) Complexes Containing the Polypyridylic Chiral Ligand (-)-Pinene[5,6]Bipyridine. Catalysts for Oxidation Reactions, *Dalt. Trans.*, **30**, 8117 (2009).
27. T. E. Kokina, L. A. Glinskaya, A. V. Tkachev, V. F. Plyusnin, Y. V. Tsoy, I. Y. Bagryanskaya, E. S. Vasilyev, D. A. Piryazev, L. A. Sheludyakova, S. V. Larionov, Chiral Zinc(II) and Cadmium(II) Complexes with a Dihydrophenanthroline Ligand Bearing (-)-A-Pinene Fragments: Synthesis, Crystal Structures and Photophysical Properties, *Polyhedron.*, **117**, 437 (2016).
28. H. L. Kwong, H. L. Yeung, C. T. Yeung, W. S. Lee, C. S. Lee, W. L. Wong, Chiral Pyridine-Containing Ligands in Asymmetric Catalysis, *Coord. Chem. Rev.*, **251**, 2188 (2007).
29. B. L. Fei, Z. X. Huang, W. S. Xu, D. D. Li, Y. Lu, W. L. Gao, Y. Zhao, Y. Zhang and Q. B. Liu, Chiral Copper(II) Complex Based on Natural Product Rosin Derivative as Promising Antitumour Agent, *J. Photochem. Photobiol. B Biol.*, **160**, 43 (2016).
30. M. K. Paul, G. Kalita, A. R. Laskar, T. D. Choudhury and N. V. S. Rao, Synthesis and Mesomorphism of New Chiral Imines and Copper(II) Complexes, *J. Mol. Struct.*, **1039**, 219 (2013).
31. E. Chinnaraja, R. Arunachalam, M. K. Choudhary, R. I. Kureshy, P. S. Subramanian, Binuclear Cu(II) Chiral Complexes: Synthesis, Characterization and Application in Enantioselective Nitroaldol (Henry) Reaction., *Appl. Organomet. Chem.*, **30**, 95 (2016).
32. R. A. Feldman and J. M. Fraile, Non-Covalent Immobilization of Chiral Copper Complexes on Al-Mcm41: Effect of The Nature of The Ligand, *Catal. Commun.*, **83**, 74 (2016).
33. C. Hee, C. Wei, J. Wei, I. Hong, K. Vei, M. Jamil and H. Ling, Enantiomeric Pair of Copper (II) Polypyridyl-Alanine Complexes: Effect of Chirality on Their Interaction with Biomolecules, *J. Inorg. Biochem.*, **160**, 1 (2016).
34. S. Muche, I. Levacheva, O. Samsonova, A. Biernasiuk, A. Malm, R. Lonsdale, Ł. Popiołek, U. Bakowsky, M. Holyńska, Synthesis, Structure and Stability of a Chiral Imine-Based Schiff-Based Ligand Derived from L-Glutamic Acid and Its [Cu4] Complex, *J. Mol. Struct.*, **1127**, 231 (2017).
35. W. C. Zhang, X. Tang, X. Lu, One-Dimensional Chiral Copper (II) Complexes with Novel Nano-Structures and Superior Antitumor Activity, *J. Inorg. Biochem.*, **156**, 105 (2016).
36. N. H. Khan, N. Pandya, K. J. Prathap, R. I. Kureshy, S. H. R. Abdi, S. Mishra, H. C. Bajaj, Chiral Discrimination Asserted by Enantiomers of Ni (II), Cu (II) And Zn (II) Schiff Base Complexes in DNA Binding, Antioxidant and Antibacterial Activities., *Spectrochim. Acta. A. Mol. Biomol. Spectrosc.*, **81**, 199 (2011).
37. X. Zhou, Y. Li, D. Zhang, Y. Nie, Z. Li, W. Gu, Copper Complexes Based on Chiral Schiff-Base Ligands: DNA/BSA Binding Ability, DNA Cleavage Activity, Cytotoxicity and Mechanism of Apoptosis, *Eur. J. Med. Chem.*, **114**, 244 (2016).
38. F. Arjmand, F. Sayeed and M. Muddassir, Synthesis of New Chiral Heterocyclic Schiff Base Modulated Cu(II)/Zn(II) Complexes: Their Comparative Binding Studies with Ct-DNA, Mononucleotides and Cleavage Activity, *J. Photochem. Photobiol. B Biol.*, **103**, 166 (2011).
39. V. C. Da Silveira, J. S. Luz, C. C. Oliveira, I. Graziani, M. R. Ciriolo, A. M. Da C. Ferreira, Double-Strand DNA Cleavage Induced by Oxindole-Schiff Base Copper(II) Complexes

- with Potential Antitumor Activity, *J. Inorg. Biochem.*, **102**, 1090 (2008).
40. M. Çolak, T. Aral, H. Hoşgören and N. Demirel, Synthesis of Novel Chiral Schiff-Base Ligands and Their Application in Asymmetric Nitro Aldol (Henry) Reaction, *Tetrahedron Asymmetry.*, **18**, 1129 (2007).
 41. M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, J. A. Montgomery, Jr., T. Vreven, K. N. Kudin, J. C. Burant, J. M. Millam, S. S. Iyengar, J. Tomasi, V. Barone, B. Mennucci, M. Cossi, G. Scalmani, N. Rega, G. A. Petersson, H. Nakatsuji, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, M. Klene, X. Li, J. E. Knox, H. P. Hratchian, J. B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R. E. Stratmann, O. Yazyev, A. J. Austin, R. Cammi, C. Pomelli, J. W. Ochterski, P. Y. Ayala, K. Morokuma, G. A. Voth, P. Salvador, J. J. Dannenberg, V. G. Zakrzewski, S. Dapprich, A. D. Daniels, M. C. Strain, O. Farkas, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. V. Ortiz, Q. Cui, A. G. Baboul, S. Clifford, J. Cioslowski, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, M. Challacombe, P. M. W. Gill, B. Johnson, W. Chen, M. W. Wong, C. Gonzalez, J. A. Pople, Gaussian 03 (Revision B.02), Gaussian, Inc., Wallingford Ct. (2004).
 42. M. E. Reichmann, S. A. Rice, C. A. Thomas, P. Doty, A Further Examination of The Molecular Weight and Size of Desoxyribose Nucleic Acid, *J. Am. Chem. Soc.*, **76**, 3047 (1954).
 43. L. H. Abdel-Rahman, R. M. El-Khatib, L. A. E. Nassr and A. M. Abu-Dief, Synthesis, Physicochemical Studies, Embryos Toxicity and DNA Interaction of Some New Iron(II) Schiff Base Amino Acid Complexes, *J. Mol. Struct.*, **1040**, 9 (2013).
 44. S. Sujarani, A. Ramu, Docking of Ethanamine Schiff Base Imines & Metal (II) Complexes, Cytotoxicity & DNA Interaction Studies, *J. Mol. Struct.*, **1079**, 353 (2015).
 45. A. Wolfe, G. H. Shimer, T. Meehan, Polycyclic Aromatic Hydrocarbons Physically Intercalate Into Duplex Regions of Denatured DNA., *Biochemistry.*, **26**, 6392 (1987).
 46. EndoFree Plasmid Purification Handbook 04/2015.
 47. W. J. Geary, The Use of Conductivity Measurements in Organic Solvents for The Characterisation of Coordination Compounds, *Coord. Chem. Rev.*, **7**, 81 (1971).
 48. M. Niu, Z. Li, H. Li, X. Li, J. Dou, S. Wang, DNA/Protein Interaction, Cytotoxic Activity and Magnetic Properties of Amino-Alcohol Schiff Base Derived Cu(II)/Ni(II) Metal Complexes: Influence of The Nuclearity and Metal Ions, *Rsc Adv.*, **5**, 37085 (2015).
 49. M. Niu, Z. Li, X. Li, X. Huang, Two Chiral Alkanolamine Schiff Base Cu(II) Complexes as Potential Anticancer Agents: Synthesis, Structure, DNA/Protein Interactions, and Cytotoxic Activity, *Rsc Adv.*, **6**, 98171 (2016).
 50. Y. A. N. Hui, DNA Interaction and Cytotoxic Activity of a Chiral Amino-Alcohol Schiff Base Derived Cu (II) Complex, *Chin. J. Struct. Chem.*, 465 (2016).
 51. M. Rezaeivala, R. Golbedaghi, M. Khalili, Coordination Chemistry of Some New Cu (II), Ni (II) and Co (II) Macroacyclic (N₂O₄) Schiff Base Complexes: X-Ray Crystal Structure of Cu (II) Complex, *Russ. J. Coord. Chem.*, **42**, 66 (2016).
 52. D. Ghosh, K. Bera, M. Kumar, S. H. R. Abdi, N. H. Khan, I. Kureshy, H. C. Bajaj, Asymmetric Allylation of Sulfonyl Imines Catalyzed by in Situ Generated Cu (II) Complexes of Chiral Amino Alcohol Based Schiff Bases, *Rsc Adv.*, **4**, 56424 (2014).
 53. J. Lu, W. Pan, R. He, S. Jin, X. Liao, B. Wu, P. Zhao, H. Guo, DNA-Binding and Photocleavage Studies of Metallofluorescein-Porphyrin Complexes of Zinc(II) and Copper(II), *Transition Met. Chem.*, **37**, 497 (2012).
 54. U. Chaveerach, A. Meenongwa, Y. Trongpanich, C. Soikum, P. Chaveerach, DNA Binding and Cleavage Behaviors of Copper(II) Complexes with Amidino-O-Methylurea and N-Methylphenyl-Amidino-O-Methylurea, and Their Antibacterial Activities, *Polyhedron.*, **29**, 731 (2010).
 55. K. Nakamoto, M. Tsuboi, G. D. Strahan, Drug-DNA Interactions: Structures and Spectra, *Methods Biochem Anal.*, **51**, 366 (2008).