

## Antifungal Activities of some *Salvia* Species Extracts on *Fusarium oxysporum* f. sp. *radicis- lycopersici* (Forl) Mycelium Growth *In-vitro*

Yılar<sup>1</sup>, M. and İ. Kadioğlu<sup>2</sup>

<sup>1</sup>Ahi Evran Univ., Fac. of Agric., Dept. of Plant Protection, Kirsehir, Turkey, melihyilar@gmail.com.

<sup>2</sup>Gaziosmanpasa Univ., Fac. of Agric., Dept. of Plant Protection, Tokat, Turkey.

(Received: December 15, 2015 and Accepted: January 23, 2016)

### ABSTRACT

The antifungal effects of essential oils and plant extracts (water, ethanol and methanol) of *Salvia cryptantha* Montbret et Aucher ex Benth, *Salvia tomentosa* Mill., *Salvia officinalis* L.(cultural form) grown in Tokat province was screened against *Fusarium oxysporum* f. sp. *radicis- lycopersici*. The essential oils and plant extracts of *S. officinalis*, *S. cryptantha* and *S. tomentosa* were determined to find the most efficient against *F. oxysporum* f. sp. *radicis- lycopersici* *in vitro*. Different volumes of either essential oils and plant extracts were mixed with the sterile PDA to obtain various concentrations. The supplemented PDA were inoculated with agar disc (5 mm in diameter) of *Fusarium oxysporum* f. sp. *radicis- lycopersici* pathogens (from 7 day-old PDA cultures) were inoculated on medium. They were incubated at 25±2 °C for 7 days. Then the blocking fungal development was calculated. The highest effects on the development of mycelium of *F. oxysporum* f. sp. *radicis- lycopersici* has shown blocking rate of 62,71% with a *S. officinalis* essential oils, this was followed by *S. tomentosa* and *S. cryptantha*. Similar results were observed in plants extracts. The highest effects on the development of mycelium of *F. oxysporum* f. sp. *radicis- lycopersici* showed *S. cryptantha* plant extracts and this *S. officinalis* and *S. tomentosa* has followed.

**Key words:** Antifungal activity, essential oils, *F. oxysporum* f. sp. *radicis- lycopersici*, *Salvia* species.

### INTRODUCTION

Diseases, pests and weeds corrupting products in agricultural areas causing significant losses. For this reason, pesticides are heavily used in order to reduce the losses caused by these pests. However, studies to this date have revealed that increased pesticide use causes several problems. Plant metabolites and plant-based medicines are thought to be less harmful to human health and the environment compared to synthetic pesticides and studies have been conducted to this end (Kordali *et al.*, 2007). Therefore, studies on the effects of various plant extracts and essential oils on plant diseases have become prominent. Lamiaceae family that involves numerous aromatic plants, is among these studied plant groups.

*Salvia* species, a member of Lamiaceae family, is one of the most important ones in this group. It is reported to include around 95 different members according to the most recent studies in Turkey (Celep *et al.*, 2009). It has an important place in the flora of Turkey and its endemism rate (51%) is also quite high (Davis, 1982 and Poyraz and Koca, 2006). Most of *Salvia* species are commonly used in food, drug, cosmetics and perfumery industry (Bağcı and Kocak, 2008). Lamiaceae plants, involving *Salvia* species are rich especially in terms of terpenoid compounds and also contain flavonoids, essential oils, phenolic compounds and some quinonoids (Durling *et al.*, 2007; Bisio *et al.*, 2011; Al-Qudah *et al.*, 2014). For this reason, A large number of studies, carried out on *Salvia* species suggested that it has numerous biological activities such as antibacterial activities (Kawahara *et al.*, 2004), antifeedant activities (Fraga

*et al.*, 2005), antioxidant activities (Lakhal *et al.*, 2013), cytotoxic activities (Lee *et al.*, 2010), antiviral activities (Tada *et al.*, 1994), antifungal activities (Abu-Darwish *et al.*, 2013), antimicrobial activities (Paknejadi *et al.*, 2012), and herbicidal activities (Bouajaj *et al.*, 2013 and Rowshan and Karimi, 2013).

In the present study, efficacy of plant extracts and essential oils of *Salvia* species against the important plant pathogen *Fusarium oxysporum* was studied.

### MATERIALS AND METHODS

#### Plant materials

*Salvia* species; *S. officinalis*, *S. tomentosa* and *S. cryptantha*, used in the experiment were collected from the province of Tokat, Turkey in 2012-13 vegetation periods by harvesting the shoot system in flowering phase. Harvested plants were dried on papers in a dark room, ground in an electric mill and kept in plastic containers to be used in the experiment.

#### Extraction of essential oils

Essential oils of the plants were obtained by hydro-distillation method using a Schilcher device. Distilled water was added to weighed plant samples (1:10 w/v) and boiled for 2 hrs. Obtained essential oils were maintained until used in the experiment (Telci *et al.*, 2006).

#### Preparation of water, methanol and ethanol extracts of plant samples

##### Water extracts

Dried herbal materials were powdered by grinding them in a plant grinding mill. 400 gr of ground plant material was placed in a glass container containing

1000 ml of distilled water and shaken for 24 hrs at 120 rpm in an orbital shaker and then solid residues were removed using filter papers. Solid residues were completely removed using centrifuge for 15 min at 5000 rpm.

### Methanol and ethanol extracts

100 gr from each plant material were put in 1 liter erlenmayers and methanol, and ethanol was added as 600 ml of each. Mixtures were shaken for 24 hrs at room temperature, at 120 rpm, in an orbital shaker. The extract was then filtered using paper filters. Methanol and ethanol were removed by evaporating at 32-40°C. Remaining extract was used to prepare a stock solution with distilled water (Kadioglu and Yanar 2004).

### Fungus culture

The plant pathogen fungus used in this study was obtained from stock cultures found at Phytopathology laboratories of Department of Plant Protection, Faculty of Agriculture, Gaziosmanpasa University, Turkey. Fungus culture was used after being developed for 7 days at 25±2°C in 60 mm Petri dishes containing 10 ml of Potato Dextrose Agar (PDA).

### In- vitro antifungal activity of plant essential oils and extracts

PDA prepared to be used in the experiment was autoclaved and chilled to 40°C. Essential oils were mixed with melted sterile PDA at the concentrations of 0, 100, 500, 1000 and 2000 ppm. PDA was poured into 60 mm Petri dishes (as 10 mm). Different plant extracts obtained (water, ethanol and methanol) were mixed with melted sterile PDA to have final concentrations of 1, 3, 7, 10 and 20%, and then poured into 60 mm Petri dishes (as 10 ml). Mycelium discs (5 mm in diameter) obtained from the 7-day fungus culture were placed in the centre of Petri dishes. After inoculation, fungus culture was left for incubation at 28°C for 7 days. Fungal development was recorded after 7 days (Hadizadeh *et al.*, 2009). Inhibition in the development was calculated using the following formula (Pandey *et al.*, 1982):

$$I = \frac{DC - DT}{DT} \times 100$$

Where:- I: Inhibition percentage compared to the control (Mycelium development), DC: Mycelium development in the control and DT: Mycelium development in essential oil applications.

PDA without essential oils and extracts was used as a negative control and synthetic Propineb fungicide (0.4 g/200 mL PDA) was used as a positive control. The experiments were repeated twice and replicated four.

### Statistical Analysis

Analysis of variance (ANOVA) was used to determine the significance levels of differences

between experiment treatments, and averages were compared using the DUNCAN test. Statistical analyses were carried out using the SPSS software.

## RESULTS AND DISCUSSION

Essential oils and plant extracts of the three different *Salvia* species (*S. officinalis*, *S. tomentosa* and *S. cryptantha*) were found to be significantly effective on *F. oxysporum* mycelium. Essential oil from *S. officinalis*, one of the *Salvia* species used in the trial had the highest impact on *F. oxysporum* f. sp. *radicis-lycopersici* mycelium development (62.71% blocking rate), followed by *S. cryptantha* (53.39%) and then *S. tomentosa* (29.44%) (Table 1). An increase in the effect of blocking effect of plant extracts on *F. oxysporum* was observed depending on the dosage increase and extract used. While, the water extract of *S. officinalis* had the highest effect, followed by plant essential oil, ethanol and methanol extracts. The water extract of *S. officinalis* had a blocking rate of (65.29%) at the highest dosage, (58.63%) for the ethanol extract and (53.84%) for the methanol extract (Table 1). Onaran *et al.* (2014) indicated that *Thymus fallax* Fish & Mey., *Origanum vulgare* L. and *Mentha dumetorum* Schult plant essential oils blocked *F. oxysporum* mycelium development to a significant degree. In another study Hadi *et al.* (2013) reported that *Mentha piperita* L. extracts blocked *F. oxysporum* spore germination and mycelium development.

Among *S. cryptantha* plant essential oil and extracts, the water extract had a complete blocking effect on *F. oxysporum* (100%), followed by the methanol and ethanol extracts (67.66 - 67.77%), and the essential oil (53.40%) (Table 1). The ethanol extract of *S. tomentosa* had the highest blocking rate on *F. oxysporum* mycelium development compared to the control with 77.64%. The methanol extract blocked *F. oxysporum* mycelium development with a rate of 62.68%, the water extract with 41.47% and the essential oil with 29.44% (Table 1). However, differences regarding this effect were identified depending on the dosage of application and plant extracts and essential oils used. It was reported in different studies that *Salvia* species had anti-fungal effect on *Fusarium* species. *Salvia sclarea* essential oil was effective on *F. tricinctum* and *F. sporotrichioides* species (Džamič *et al.*, 2008). *Salvia sclarea* essential oil was also effective on *F. oxysporum* f. sp. *dianthi* development (Pitarokili *et al.*, 2002), *Salvia tigrina* ethanol extract was effective on *F. oxysporum* (Dulger and Hacıoglu, 2008). It was also found that *S. officinalis* plant was effective on *Candida* spp. and *Aspergillus niger* (Badiee *et al.*, 2012 and Abu-Darwish *et al.*, 2013), *Plasmopara*

Table (1): Effects on mycelium growth rate of of different *Salvia* spp. on *F. oxysporum* f. sp. *radicis-lycopersici*

		Doses	<i>S. officinalis</i>	<i>S. cryptantha</i>	<i>S. tomentosa</i>			Doses	<i>S. officinalis</i>	<i>S. cryptantha</i>	<i>S. tomentosa</i>
Essential oil	Kontrol	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	Methanol extract	Kontrol	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40
	100µl	31.67 <sup>b</sup> ±0.69	35.57 <sup>b</sup> ±3.18	46.20 <sup>ab</sup> ±2.45	46.20 <sup>ab</sup> ±2.45		1%	26.24 <sup>b</sup> ±0.31	24.61 <sup>b</sup> ±1.17	39.10 <sup>b</sup> ±0.23	39.10 <sup>b</sup> ±0.23
	500 µl	29.72 <sup>b</sup> ±0.95	34.26 <sup>b</sup> ±1.46	42.45 <sup>b</sup> ±0.50	42.45 <sup>b</sup> ±0.50		3%	25.79 <sup>b</sup> ±0.43	23.72 <sup>b</sup> ±0.85	30.48 <sup>c</sup> ±1.28	30.48 <sup>c</sup> ±1.28
	1000 µl	27.09 <sup>b</sup> ±0.66	29.75 <sup>bc</sup> ±0.39	42.33 <sup>b</sup> ±0.41	42.33 <sup>b</sup> ±0.41		7%	25.75 <sup>b</sup> ±0.25	23.62 <sup>b</sup> ±0.80	27.29 <sup>c</sup> ±0.48	27.29 <sup>c</sup> ±0.48
	2000 µl	19.26 <sup>c</sup> ±0.69	24.07 <sup>c</sup> ±0.77	36.45 <sup>c</sup> ±1.49	36.45 <sup>c</sup> ±1.49		10%	23.88 <sup>b</sup> ±0.34	21.58 <sup>b</sup> ±0.95	26.58 <sup>c</sup> ±0.15	26.58 <sup>c</sup> ±0.15
	Propineb	0.00 <sup>d</sup> ±0.00	0.00 <sup>d</sup> ±0.0	0.00 <sup>d</sup> ±0.00	0.00 <sup>d</sup> ±0.00		20%	21.37 <sup>c</sup> ±0.37	16.70 <sup>c</sup> ±0.42	19.28 <sup>d</sup> ±0.10	19.28 <sup>d</sup> ±0.10
						Propineb	0.00 <sup>d</sup> ±0.00	0.00 <sup>d</sup> ±0.00	0.00 <sup>c</sup> ±0.00	0.00 <sup>c</sup> ±0.00	
Water extract	Kontrol	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	Ethanol extract	Kontrol	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40	51.66 <sup>a</sup> ±3.40
	1%	29.33 <sup>b</sup> ±1.12	36.09 <sup>b</sup> ±0.86	51.60 <sup>a</sup> ±0.78	51.60 <sup>a</sup> ±0.78		1%	39.76 <sup>b</sup> ±1.34	25.38 <sup>b</sup> ±0.64	31.09 <sup>b</sup> ±0.07	31.09 <sup>b</sup> ±0.07
	3%	28.60 <sup>b</sup> ±0.22	28.95 <sup>c</sup> ±0.91	47.60 <sup>b</sup> ±1.05	47.60 <sup>b</sup> ±1.05		3%	39.79 <sup>b</sup> ±0.37	23.04 <sup>bc</sup> ±0.35	23.67 <sup>c</sup> ±0.23	23.67 <sup>c</sup> ±0.23
	7%	27.81 <sup>b</sup> ±0.19	26.93 <sup>c</sup> ±0.10	35.89 <sup>c</sup> ±2.40	35.89 <sup>c</sup> ±2.40		7%	23.17 <sup>c</sup> ±1.02	19.59 <sup>cd</sup> ±0.49	22.75 <sup>c</sup> ±0.20	22.75 <sup>c</sup> ±0.20
	10%	26.88 <sup>b</sup> ±0.51	18.68 <sup>d</sup> ±0.53	31.55 <sup>cd</sup> ±0.57	31.55 <sup>cd</sup> ±0.57		10%	23.01 <sup>c</sup> ±0.23	18.76 <sup>d</sup> ±0.31	20.72 <sup>c</sup> ±0.19	20.72 <sup>c</sup> ±0.19
	20%	23.84 <sup>c</sup> ±0.25	0.00 <sup>e</sup> ±0.00	30.23 <sup>d</sup> ±1.12	30.23 <sup>d</sup> ±1.12		20%	17.93 <sup>c</sup> ±3.69	16.64 <sup>d</sup> ±0.30	11.55 <sup>d</sup> ±0.09	11.55 <sup>d</sup> ±0.09
	Propineb	0.00 <sup>d</sup> ±0.00	0.00 <sup>e</sup> ±0.00	0.00 <sup>e</sup> ±0.00	0.00 <sup>e</sup> ±0.00		Propineb	0.00 <sup>d</sup> ±0.00	0.00 <sup>e</sup> ±0.00	0.00 <sup>c</sup> ±0.00	0.00 <sup>c</sup> ±0.00

\*Means in the same column with the same letter were not significantly different by ANOVA (a = 0.05)

*viticola* (Dagostin *et al.*, 2010), *Alternaria* spp. (Mahmoudi and Ahmadi, 2013); while *S. cryptantha* and *S. tomentosa* showed antimicrobial and antibactericidal effects (Haznedaroglu *et al.*, 2001). These findings are similar to the obtained results. Biological activities of *Salvia* species are a result of the compounds contained by plants. Because it was found in numerous studies conducted on *Salvia* species that these plants were rich in camphor, linalool, eucalyptol (1,8-cineole), borneol compounds (Pandey 2009 and Okamoto *et al.*, 2011), and phenolics (Lu and Foo 2000 and Yumrutas *et al.*, 2011).

This study and similar previous studies showed that *Salvia* species are effective on *F. oxysporum* f. sp. *radicis-lycopersici* mycelium development. Considering the environmental damage caused by fungicides commonly used against plant diseases, obtained findings of this study may give some lights to future studies in this field.

#### ACKNOWLEDGEMENT

The authors thank Assistant Prof. Dr. Sabriye BELGUZAR, Dept. of Plant Protection, Fac. of Agriculture, Gop Univ., Tokat, Turkey and Assistant Prof. Dr. Abdurrahman ONARAN (Dept. of Plant Protection, Fac. of Agriculture, Ahi Evran Univ., Tokat, Turkey) for providing the stock culture of the fungus.

#### REFERENCES

Abu-Darwish, M. S., C. Cabral, I. V.Ferreira, M. J. Gonçalands, C. Cavaleiro, M. T. Cruz, T. H. Albdour and L., Salgueiro, 2013. Essential oil of common sage (*Salvia officinalis* L.) from Jordan: Assessment of safety in mammalian cells and its antifungal and anti-inflammatory potential.

- BioMed Research International. volume 2013, article ID 538940, 9 pages
- Al-Qudah, M., H. Al-Jaber, M. H. A. Zarga and S. T. A., Orabi 2014. Flavonoid and phenolic compounds from *Salvia palaestina* L., growing wild in Jordan and their antioxidant activities. *Phytochemistry*. 99: 115-120.
- Badiee, P., A. R. Nasirzadeh and M., Motaffaf, 2012. Comparison of *Salvia officinalis* L. essential oil and antifungal agents against *Candida* species. *Journal of Pharmaceutical Technology & Drug Research*. 1-5.
- Bağcı, E. and A., Koçak, 2008. Composition of essential oils of *Salvia palaestina* Benth and *S. tomentosa* Miller species, a chemotaxonomical approach. *Science and Eng. J. of Firat Univ.* 20 (1): 35-41.
- Bisio, A., G. Damonte, D. Fraternali, E. Giacomelli, A. Salis, G. Romussi, S. Cafaggi, D. Ricci, and N. D. Tommasi, 2011. Phytotoxic clerodane diterpenes from *Salvia miniata* Fernald (Lamiaceae). *Phytochemistry*. 72: 265-275
- Bouajaj, S., A. Benyamma, H. Bouamama, A. Romane, D. Falconieri, A. Piras, and B., Marongiu, 2013. Antibacterial, allelopathic and antioxidant activities of essential oil of *Salvia officinalis* L. growing wild in the Atlas Mountains of Morocco. *Natural Product Research*. 27 (18):1673-1676.
- Celep, F., M. Doğan, and A. Duran, 2009. A new record for the flora of Turkey: *Salvia viscosa* Jacq. (Labiatae). *Türk J. Bot.* 32: 57-60
- Dagostin, S., T. Formolo, O. Giovannini, and I., Pertot 2010. *Salvia officinalis* extract can protect grapevine against *Plasmopara viticola*. *Plant Disease*. 94(5):575-580.
- Davis, P. H. 1982. *Flora of Turkey and The East Aegean Island*, Vol. 7, Edinburgh University Press, Edinburgh.
- Dulger, B. and N. Hacıoğlu 2008. Antifungal Activity

- of Endemic *Salvia tigrina* in Turkey. *Tropical Journal of Pharmaceutical Research*. 7 (3): 1051-1054.
- Durling, N. E., O. J. Catchpole, J. B. Grey, R. F. Webby, K. A. Mitchell, L. Y. Foo, and N. B. Perry 2007. Extraction of phenolics and essential oil from dried sage (*Salvia officinalis*) using ethanol-water mixtures. *Food Chemistry*. 101: 1417-1424.
- Džamič, A., M. Sokovič, M. Ristič, S. Grujič-Jovanović, J. Vukojevič, and P. D. Marin 2008. Chemical composition and antifungal activity of *Salvia sclarea* (Lamiaceae) essential oil. *Arch. Biol. Sci., Belgrade*. 60(2): 233-237.
- Fraga, B. M., C. E. Díaz, A. Guadaño and A. González-Coloma 2005. Diterpenes from *Salvia broussonetii* transformed roots and their insecticidal activity. *J. Agric. Food Chem.* 53: 5200-5206.
- Hadi, M., B. Kashefi, A. Sobhanipur and M. Rezaarabsorkhi 2013. Study on effect of some medicinal plant extracts on growth and spore germination of *Fusarium oxysporum* schlecht. *In vitro*. *American-Eurasian J. Agric. & Environ. Sci.*, 13 (4): 581-588.
- Hadizadeh, I., B. Peivastegan and H. Hamzehzarghani 2009. Antifungal Activity of essential oils from some medicinal plants of Iran against *Alternaria alternata*. *American Journal of Applied Sciences*. 6(5): 857-861.
- Haznedaroğlu, M. Z., N. U. Karabay, and U. Zeybek 2001. Antibacterial activity of *Salvia tomentosa* essential oil. *Fitoterapia*. 72: 829-831.
- Kadioglu, I., Y. and Yanar 2004. Allelopathic effects of plant extracts against seed germination of some weeds. *Assian Journal of Plant Sciences*. 3(4): 472-475.
- Kawahara, N., T. Tamura, I. Mayumi, T. Hosoe, K. Kawai, S. Sekita, M. Satake and Y. Goda 2004. Diterpenoid glucosides from *Salvia greggii*. *Phytochemistry*. 65:2577-2581
- Kordali, Ş., S. Kotan, and A., Çakır, 2007. Screening of antifungal activities of 21 oxygenated monoterpenes in-vitro as plant disease control agents. *Allelopathy Journal*. 19(2):373-392.
- Lakhal, H., H. Ghorab, S. Chibani, A. Kabouche, Z. Semra, F. Smati, S. Abuhamdah and Z. Kabouche 2013. Chemical composition and biological activities of the essential oil of *Salvia officinalis* from Batna (Algeria). *Der Pharmacia Lettre*. 5 (3): 310-314
- Lee, W. Y. W., C. C. M. Cheung, K. W. K. Liu, K. P. Fung, J. Wong, P. B. S. Lai and J. H. K. Yeung 2010. Cytotoxic effects of tanshinones from *Salvia miltiorrhiza* on doxorubicin-resistant human liver cancer cells. *J.Nat. Prod.*73: 854-859.
- Lu, Y. and Y. Foo, 2000. Flavonoid and phenolic glycosides from *Salvia officinalis*. *Phytochemistry*. 55:263-267.
- Mahmoudi, E. and A. Ahmedi 2013. Evaluation of *Salvia officinalis* antifungal properties on the growth and morphogenesis of *Alternaria alternata* under in-vitro conditions. *Tech. J. Engin. & App. Sci.* 3 (17): 2062-2069
- Okamoto, Y., K. Yamaji and K. Kobayashi 2011. Allelopathic activity of camphor released from camphor tree (*Cinnamomum camphora*). *Allelopathy Journal*. 27 (1): 123-132.
- Onaran, A., M. Yılar, S. Belgüzar, Y. Bayan and H. Akşit 2014. Antifungal and bioherbicidal properties of essential oils of *Thymus fallax* Fish & Mey., *Origanum vulgare* L. and *Mentha dumetorum* Schult. *Asian J. Chem.* 26(16)/pp 5159-5164
- Paknejadi, M., F. Foroohi and M. Yousefzadi 2012. Antimicrobial activities of the essential oils of five *Salvia* species from Tehran province, Iran. *Journal of Paramedical Sciences*. 3(2): 12-18.
- Pandey, D. K. 2009. Allelochemicals in *Parthenium* in response to biological activity and the environment. *Indian Journal of Weed Science*. 41(3&4) : 111-123.
- Pandey, D. K., N. N. Tripathi, R. D. Tripathi, and S. N. Dixit 1982. Fungitoxic and phytotoxic properties of essential oil of *Hyptis suaandolens*. *Z. Pflanzenkrankheiten Pflanzenschutz*, 89: 344–349.
- Pitarokili, D., M. Couladis, N. Petsikos-Panayotarou, and O. Tzakou 2002. Composition and antifungal activity on soil-borne pathogens of the essential oil of *Salvia sclarea* from Greece. *J. Agric. Food Chem.* 50: 6688-6691.
- Poyraz, İ. E. and F. Koca 2006. Morphological investigations on some medicinal *Salvia* L. Species in Eskişehir. *Anadolu University Journal of Science and Technology*. 7(2): 443-450.
- Rowshan, V. and S. Karimi 2013. Essential oil composition and allelopathic affect of *Salvia macrosiphon* BOISS. on *Zea mays* L. *International Journal of Agriculture: Research and Review*. 3 (4): 788-794.
- Tada, M., K. Okuno, K. Chiba, E. Ohnishi and T. Yoshii 1994. Antiviral diterpenes from *Salvia officinalis*. *Phytochemistry*. 35: 539-541.
- Telci, I., E. Bayram, G. Yilmaz and B. Avci 2006. Variability in essential oil composition of Turkish basils (*Ocimum basilicum* L.). *Biochemical Systematics and Ecology*, 34, 489-497.
- Yumrutas, O., A. Sokmen and N. Ozturk 2011. Determination of *in vitro* antioxidant activities and phenolic compounds of different extracts of *Salvia verticillata* ssp. *verticillata* and *Salvia verticillata* ssp. *amasiaca* from Turkey's flora. *Journal of Applied Pharmaceutical Science*.1(10): 43-46.