

Cutaneous manifestations of COVID-19: A review of the published literature

Ömer Faruk Elmas¹ | Abdullah Demirbaş² | Kemal Özyurt¹ |
Mustafa Atasoy³ | Ümit Türsen⁴

¹Department of Dermatology, Kırşehir Ahi Evran University, Kırşehir, Turkey

²Department of Dermatology, Konya Numune State Hospital, Konya, Turkey

³Department of Dermatology, Health Science University, Kayseri City Hospital, Kayseri, Turkey

⁴Department of Dermatology, Mersin University, Mersin, Turkey

Correspondence

Kemal Özyurt, Department of Dermatology, Kırşehir Ahi Evran University, Kırşehir, Turkey.
Email: drkozyurt@gmail.com

Abstract

COVID-19 is a highly contagious respiratory tract infection caused by severe acute respiratory syndrome coronavirus 2. COVID-19 outbreak, which caused thousands of deaths, has been declared a pandemic by the World Health Organization in March 2020. The infection has been reported to demonstrate different types of cutaneous manifestations including urticarial, maculopapular, papulovesicular, purpuric, livedoid, and thrombotic-ischemic lesions. Given the high mortality rate of the infection, timely and accurate identification of relevant cutaneous manifestations may play a key role in the early diagnosis and management. In this study, we provide a review with a focus on the reported cutaneous manifestations of COVID-19.

KEYWORDS

coronavirus, COVID-19, cutaneous manifestations, skin

1 | INTRODUCTION

COVID-19 is a highly contagious respiratory tract disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which was first reported on December 1, 2019 from Wuhan, China.¹ COVID-19 outbreak has been declared a pandemic by the World Health Organization on March 11, 2020 and as of April 26, 2020, approximately 3 million cases were identified worldwide and more than 200 thousand deaths occurred.^{1,2} SARS-CoV-2 is transmitted through virus-containing droplets and contaminated objects. The incubation period of the infection is ranging between 2 and 14 days, with an average of 5 days.³ Fever, cough, sore throat, shortness of breath, muscle pain, nausea, diarrhea, and loss of taste and smell have been reported as common symptoms of COVID-19.⁴ In addition, organ dysfunction, progressive respiratory failure, and coagulation disorders, which are associated with high mortality, are also reported.⁵

Infectious diseases that have serious effects on public health may primarily or secondarily involve the skin. Therefore, timely and accurate identification of skin lesions in patients with fever and rash may significantly increase diagnostic accuracy.

Viral infections are common both in children and adults and may cause a wide variety of mucocutaneous manifestations. In the current

literature, COVID-19 has also been reported to demonstrate different types of cutaneous manifestations including urticarial, maculopapular, varicella-like, purpuric, livedoid, and thrombotic-ischemic rashes. In this study, we aimed to review the cutaneous findings that have been associated with COVID-19.

2 | URTICARIAL LESIONS

Infections are one of the common causes of urticaria. Bacterial (*Streptococcus*, *Mycoplasma*, etc.), parasitic (*Schistosomiasis*, *Strongyloidiasis*, *Filariasis*, etc.), fungal and viral infections (*Epstein-Barr virus*, *enterovirus*, *adenovirus*, *hepatitis B*, *hepatitis C*, etc.) can cause urticarial eruptions.⁶

In the most comprehensive study focused on the cutaneous manifestations of COVID-19, Casas Galvan et al⁷ reported that 19% of the 375 patients showed urticarial eruption mostly distributed in the trunk or disperse. Only a few cases were limited to the palmar region. Recalcati et al reported that 18 (20.4%) out of 88 patients with COVID-19 had skin findings and three of these showed urticarial lesions. The lesions predominantly distributed over the trunk, and the urticarial eruption was not related to disease severity.⁸ Hedou et al

observed urticarial eruption in two out of 103 patients with COVID-19. In one of these patients, the eruption occurred in the prodromal stage of the infection.⁹ Henry et al identified urticarial lesions limited to the face and acral regions in a 27-year-old COVID-19 positive female patient who did not have a high fever.¹⁰ In contrast, C van Damme et al reported two patients with widespread urticarial lesions with fever as the initial signs of COVID-19.¹¹

It is obvious that urticaria may be a cutaneous manifestation of a viral infection. In this context, during the pandemic, patients with urticarial lesions should be carefully evaluated for the possibility of COVID-19.

3 | MACULOPAPULAR LESIONS

Maculopapular eruption is a common cutaneous condition that can be caused by drugs, and bacterial and viral infections. It is often associated with a viral or bacterial infection when accompanied by symptoms such as fever, headache, muscle pain, and respiratory distress. Infectious mononucleosis, measles, scarlet fever, hand-foot-mouth disease, herpes, hepatitis B, hepatitis C, Zika virus, Ebola, and HIV are viral conditions that can cause maculopapular eruptions.¹²

Casas Galvan et al⁷ reported that 47% of 375 patients with COVID-19 had a maculopapular eruption. Some of them showed perifollicular distribution and scaling, while some were similar to pityriasis rosea. The authors also reported that a few cases showed infiltrated papular lesions, resembling erythema elevatum diutinum or erythema multiforme.

Recalcati et al⁸ identified maculopapular eruptions in 14 out of 18 patients with COVID-19, while Hedou et al⁹ reported similar eruptions in two patients. Mahe et al,¹³ Hunt et al,¹⁴ Estebanez et al,¹⁵ Alramthan et al,¹⁶ and Ahouache¹⁷ reported single cases of maculopapular eruptions in patients with COVID-19.

Jones et al¹⁸ reported a case of Kawasaki disease in a COVID-19 positive 6-month-old female patient who fulfilled the criteria for Kawasaki disease with fever, conjunctivitis, prominent tongue papilla, dry cracked lips, polymorphous maculopapular rash, and swelling of the hands and lower extremities. The authors pointed out the necessity of further studies to clarify the potential association of COVID-19 and Kawasaki disease.

4 | PAPULOVESICULAR ERUPTIONS

In addition to the primary cutaneous diseases and drug eruptions, many bacterial (staphylococcal skin infections, gonococcal bacteremia, and vibrio vulnificus) and viral infections (enteroviral infections, herpes simplex, herpes zoster, varicella zoster, HIV, and parvovirus B19) can be manifested with vesiculobullous and papulovesicular eruptions. Infection-related vesicular lesions can be localized or widespread. While herpes simplex and herpes zoster are usually presented with limited lesions, chickenpox often shows widespread papulovesicular eruption.¹²

Recalcati et al,⁸ who conducted the first study regarding the cutaneous findings of COVID-19, reported varicella-like lesions in one of 18 patients. Marzona et al¹⁹ also identified varicella-like eruption in 12 out of 22 patients with COVID-19 and they concluded that this type of eruption may be a specific manifestation of COVID-19. Casas Galvan et al⁷ described small monomorphic vesicles located on the trunk in 9% of 375 patients with COVID-19. The authors reported that the vesicular eruptions identified were unlike polymorphic vesicles of chickenpox.

On the other hand, Hedou et al⁹ described oral herpes reactivation in an intubated patient, while Tammamo et al²⁰ reported herpes-related vesicular lesions localized on the trunk in three patients.

5 | PURPURIC ERUPTIONS

Infection-related purpuric eruptions may occur due to vascular invasion of the infectious agent or disseminated intravascular coagulation caused by toxic vascular effects of the infection. Early diagnosis and treatment of purpuric eruptions are crucial due to high rates of morbidity and mortality.

Coxsackievirus A9, Echovirus, Epstein-Barr virus, Cytomegalovirus, Measles virus, Arbovirus, and Arenavirus are the known viral causes of purpuric eruptions.¹² Jimenez-Couche et al²¹ reported a purpuric macular eruption distributed over the flexural areas, appearing on the third day of hydroxychloroquine and lopinavir/ritonavir treatments in an 84-year-old female patient.

Magro et al²² observed purpuric lesions in three patients with severe COVID-19. Histopathological examination of the lesions showed pauci-inflammatory thrombogenic vasculopathy, and C5b-9 and C4d accumulation both in the normal and lesional skin. These interesting findings suggest that, in addition to hypercoagulation, complement activation may contribute to the pathogenesis of COVID-19.

6 | LIVEDO RETICULARIS

Livedo reticularis is a cutaneous finding consisting of a mottled reticulated vascular pattern that appears as a lace-like purplish discoloration. It can be associated with progressive ischemia with or without any evidence of systemic diseases. The color change is caused by a decrease in the blood flow of the arterioles feeding the cutaneous capillaries, resulting in blue discoloration due to deoxygenated blood. Systemic diseases-related livedo reticularis is defined as secondary livedo reticularis. Hematological, rheumatic, and cardiovascular diseases and infections can cause secondary livedo reticularis.²³ Cytomegalovirus and parvovirus B19 are the reported causes of secondary livedo reticularis.²⁴

Casas Galvan et al⁷ reported that 6% of 375 patients with COVID-19 demonstrated cutaneous lesions indicating varying degrees of occlusive vascular disease. Manalo et al²⁵ identified livedo reticularis like lesions in two patients with COVID-19 and hypothesized that the lesions might be associated with microthrombosis.

TABLE 1 Key points of the articles reported cutaneous manifestations of COVID-19

Article	Country	Type of the article	Key points
Rodriguez-Jimenez et al ⁷	Spain	Original article	<ul style="list-style-type: none"> - The study included a total of 375 patients showing cutaneous manifestations of COVID-19 - 47% showed maculopapular lesions - 19% of the patients showed urticarial lesions - 19% showed pseudo-chilblain - 9% showed small monomorphic vesicles - 6% showed livedo/necrosis
Recalcati ⁸	Italy	Case series	<ul style="list-style-type: none"> - The study included 18 patients with COVID-19. - 20.4% (n = 18) of the patients showed cutaneous manifestations. (Erythematous rashes [n = 14], urticaria [n = 3], and Varicella-like vesicular lesions [n = 1]) - The lesions were usually located on the chest - There was no relationship between cutaneous manifestations and the severity of COVID-19
Hedou et al ⁹	France	Case series	<ul style="list-style-type: none"> - The study included 103 patients. - 5 (4.9%) out of 103 patients had cutaneous manifestations. (Erythematous rash [n = 2], urticaria [n = 2], and oral herpes activation [n = 1]) - One of the patients had urticaria in prodromal stage. - The lesions disappeared with a median time of 48 h.
Henry et al ¹⁰	France	Single case report	<ul style="list-style-type: none"> - 27-y-old female with urticarial lesions distributed over the face and acral areas. - The lesions started before fever and moderately responded to antihistamine treatment.
Mahe et al ¹³	France	Single case report	<ul style="list-style-type: none"> - A 64-y-old female with type 2 diabetes with erythematous macular lesions distributed over the chest, axilla, and antecubital region. - The lesions appeared at the fourth day of the infection and disappeared on the following day.
Jimenez-Cauche et al ²¹	Spain	Single case report	<ul style="list-style-type: none"> - A 84-y-old female with flexural erythematous purpuric macules. - The lesions appeared 3 d after the treatment of hydroxychloroquine, lopinavir/ritonavir.
Hunt et al ¹⁴	United States	Single case report	<ul style="list-style-type: none"> - A 20-y-old male with widely distributed morbilliform rash sparing the face and mucosa.
Marzano et al ¹⁹	Italy	Multicentric case series	<ul style="list-style-type: none"> - Varicella-like eruption in 22 patients. - Mean age of 60. - The time interval between the symptoms of COVID-19 and skin manifestations was 3 d. - Mean duration of the eruption was 8 d. - The most common location was chest. - There was no face and mucosa involvement. - The histopathological examination of the lesions was in keeping with viral exanthem.
Estebanez et al ¹⁵	Spain	Single case report	<ul style="list-style-type: none"> - A 28-y-old female with itchy erythematous yellow papules on both heels at the 13th day of the infection. - The patient used paracetamol 10 d before the lesions started. - The lesions were unresponsive to topical corticosteroids.
Manalo et al ²⁵	United States	Two cases report	<ul style="list-style-type: none"> - The authors reported two patients with livedo reticularis like eruption. - Patient 1: 67-y-old male. The lesions appeared on the seventh day of the infection and disappeared after 19 h. Transient hematuria accompanied. - Patient 2: 47-y-old female. The lesions appeared following sun exposure on the 10th day of the infection and disappeared after 20 min. The patient had Hashimoto thyroiditis and portal vein thrombosis.
Mazzotta et al ²⁸	Italy	Single case report	<ul style="list-style-type: none"> - A 13-y-old boy with erythematous purple macules, ulceration, and crusts. - The lesions disappeared in a few days.
Alramthan et al ¹⁶	Qatar	Two cases report	<ul style="list-style-type: none"> - Two female patients aged 27 and 35 y. - Erythematous purple maculopapular lesions on the dorsal hands, and subungual erythema.

(Continues)

TABLE 1 (Continued)

Article	Country	Type of the article	Key points
Magro et al ²²	United States	Case series	<ul style="list-style-type: none"> - A series of five patients with purpuric cutaneous lesions. - The lesions histologically demonstrated a pauci-inflammatory thrombogenic vasculopathy with deposition of C5b-9 and C4d in both involved and normally appearing skin.
Tamaro et al ²⁰	Italy	Case series	<ul style="list-style-type: none"> - Three patients with herpetiform lesions on the trunk. - The lesions were considered as suggestive of the infections caused by the members of the Herpesviridae family.
C van Damme et al ¹¹	Belgium	Two cases report	<ul style="list-style-type: none"> - Two patients (aged 39 and 71 y) with concomitant fever and extensive urticarial lesions as initial findings of COVID-19.
Ahouache et al ¹⁷	France	Single case report	<ul style="list-style-type: none"> - A 57-year-old woman with fever lasting for 4 d. - Diffuse fixed erythematous maculopapular lesions over the limbs and trunk, with burning sensation over the palms that appeared 2 d before fever.
Kolivras et al ²⁹	Belgium	Single case report	<ul style="list-style-type: none"> - A 23-y-old male patient with acute-onset violaceous plaques on the toes and lateral feet. - The lesions appeared 3 d after a low-grade fever and cough. - The patient had a history of psoriasis, which was treated with secukinumab; however, the secukinumab was stopped 1 mo ago.
Jones et al ¹⁸	United States	Single case report	<ul style="list-style-type: none"> - A 6-mo-old female with COVID-19 and concurrent Kawasaki disease. - The patient fulfilled the criteria for Kawasaki disease with fever, conjunctivitis, prominent tongue papilla, dry cracked lips, polymorphous maculopapular rash, and swelling of the hands and lower extremities. - The patient was free of respiratory symptom and the clinical course was mild.

7 | THROMBOTIC-ISCHEMIC LESIONS

Infection-related cutaneous thrombotic and ischemic lesions may occur due to direct vascular invasion of the infectious agent, vascular occlusion, or disseminated intravascular coagulation. These types of lesions are one of the important manifestations of serious infectious conditions with high morbidity and mortality such as meningococemia, and staphylococcal and pneumococcal septicemias.^{26,27}

Casas Galvan et al⁷ identified asymmetrical acral erythema and edema with vesicles or pustules, which were described "pseudo-chilblain," in 19% of 375 patients with COVID-19. The lesions were also reported to have purpuric areas.

Mazzotta et al²⁸ reported erythematous purple-colored macular lesions developed on the toes of a 13-year-old boy with COVID-19. The lesions disappeared in a short time following the ulceration and subsequent crusting. The authors hypothesized that the lesions may be caused by acro-ischemia due to endothelial damage and micro-thrombi, induced by the virus.²⁸ Kolivras et al²⁹ presented a case of COVID-19 infection-induced chilblain in a patient with the previous history of psoriasis.

Purpuric lesions, livedo reticularis, and thrombotic-ischemic lesions reported in relation to the disease support the view that hypercoagulation may play an important role in the high mortality rate of the infection.

Key points of the articles reported that cutaneous manifestations of COVID-19 have been shown in Table 1.

8 | CONCLUSIONS

In the limited number of studies available to date, COVID-19 has been reported to be associated with different types of cutaneous manifestations. In this context, during the pandemic, the possibility of COVID-19 should be carefully evaluated, particularly in patients presenting with cutaneous findings reported. On the other hand, it should be kept in mind that the disease may also show the skin findings related to viral infections in general. Given the high mortality rate of the infection, timely and accurate identification of the relevant cutaneous manifestations may play a key role in the early diagnosis and management. It is obvious that more in-depth researches are needed to understand the relationship between COVID-19 and skin.

ORCID

Ömer Faruk Elmas  <https://orcid.org/0000-0002-5474-6508>

Abdullah Demirbaş  <https://orcid.org/0000-0002-3419-9084>

Kemal Özyurt  <https://orcid.org/0000-0002-6913-8310>

Ümit Türsen  <https://orcid.org/0000-0002-5807-6759>

REFERENCES

1. World Health Organization. Naming the coronavirus disease (COVID-19) and the virus that causes it; 2020. [https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-\(covid-2019\)-and-the-virus-that-causes-it](https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance/naming-the-coronavirus-disease-(covid-2019)-and-the-virus-that-causes-it)
2. Hui DS, Azhar EI, Madani TA, et al. The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health—The latest 2019

- novel coronavirus outbreak in Wuhan, China. *Int J Infect Dis*. 2020; 91:264.
3. Velavan TP, Meyer CG. The COVID-19 epidemic. *Trop Med Int Health*. 2020;25(3):278-280.
 4. Li YC, Bai WZ, Hashikawa T. The neuroinvasive potential of SARS-CoV2 may play a role in the respiratory failure of COVID-19 patients. *J Med Virol*. 2020;92(6):552-555.
 5. Singhal T. A review of coronavirus Disease-2019 (COVID-19). *Indian J Pediatr*. 2020;87(4):281-286.
 6. Wedi B, Raap U, Wieczorek D, Kapp A. Urticaria and infections. *Allergy Asthma Clin Immunol*. 2009;5(1):10.
 7. Galvan Casas C, Catala A, Carretero Hernandez G, et al. Classification of the cutaneous manifestations of COVID-19: a rapid prospective nationwide consensus study in Spain with 375 cases. *Br J Dermatol*. 2020.
 8. Recalcati S. Cutaneous manifestations in COVID-19: a first perspective. *J Eur Acad Dermatol Venereol*. 2020;34:e212-e213.
 9. Hedou M, Carsuzaa F, Chary E, Hainaut E, Cazenave-Roblot F, Masson Regnault M. Comment on "Cutaneous manifestations in COVID-19: a first perspective" by Recalcati S. *J Eur Acad Dermatol Venereol*. 2020.
 10. Henry D, Ackerman M, Sancelme E, Finon A, Esteve E. Urticarial eruption in COVID-19 infection. *J Eur Acad Dermatol Venereol*. 2020.
 11. van Damme C, Berlingin E, Saussez S, Accaputo O. Acute urticaria with pyrexia as the first manifestations of a COVID-19 infection. *J Eur Acad Dermatol Venereol*. 2020.
 12. Kang JH. Febrile illness with skin rashes. *Infect Chemother*. 2015;47(3):155-166.
 13. Mahe A, Birckel E, Krieger S, Merklen C, Bottlaender L. A distinctive skin rash associated with coronavirus disease 2019? *J Eur Acad Dermatol Venereol*. 2020.
 14. Hunt M, Koziatek C. A case of COVID-19 pneumonia in a young male with full body rash as a presenting symptom. *Clin Pract Cases Emerg Med*. 2020;4(2):219-221.
 15. Estebanez A, Perez-Santiago L, Silva E, Guillen-Climent S, Garcia-Vazquez A, Ramon MD. Cutaneous manifestations in COVID-19: a new contribution. *J Eur Acad Dermatol Venereol*. 2020.
 16. Alramthan A, Aldaraji W. Two cases of COVID-19 presenting in clinical picture resembling chilblains disease: first report from the Middle East. *Clin Exp Dermatol*. 2020.
 17. Ahouach B, Harant S, Ullmer A, et al. Cutaneous lesions in a patient with COVID-19: are they related? *Br J Dermatol*. 2020.
 18. Jones VG, Mills M, Suarez D, et al. COVID-19 and Kawasaki disease: novel virus and novel case. *Hosp Pediatr*. 2020;10(5):537-540.
 19. Marzano AV, Genovese G, Fabbrocini G, et al. Varicella-like exanthem as a specific COVID-19-associated skin manifestation: multicenter case series of 22 patients. *J Am Acad Dermatol*. 2020.
 20. Tammaro A, Adebajo GAR, Parisella FR, Pezzuto A, Rello J. Cutaneous manifestations in COVID-19: the experiences of Barcelona and Rome. *J Eur Acad Dermatol Venereol*. 2020.
 21. Jimenez-Cauhe J, Ortega-Quijano D, Prieto-Barrios M, Moreno-Arrones OM, Fernandez-Nieto D. Reply to "COVID-19 can present with a rash and be mistaken for dengue": petechial rash in a patient with COVID-19 infection. *J Am Acad Dermatol*. 2020.
 22. Magro C, Mulvey JJ, Berlin D, et al. Complement associated microvascular injury and thrombosis in the pathogenesis of severe COVID-19 infection: a report of five cases. *Transl Res*. 2020.
 23. Sajjan VV, Lunge S, Swamy MB, Pandit AM. Livedo reticularis: a review of the literature. *Indian Dermatol Online J*. 2015;6(5):315-321.
 24. Arslan F, Batirel A, Mert A, Ozer S. Cytomegalovirus (CMV)-related cutaneous necrotizing vasculitis: case report and literature review. *Braz J Infect Dis*. 2012;16(5):482-485.
 25. Manalo IF, Smith MK, Cheeley J, Jacobs R. A dermatologic manifestation of COVID-19: transient livedo reticularis. *J Am Acad Dermatol*. 2020.
 26. Thompson MJ, Ninis N, Perera R, et al. Clinical recognition of meningococcal disease in children and adolescents. *Lancet*. 2006;367(9508):397-403.
 27. Launay E, Gras-Le Guen C, Martinot A, et al. Suboptimal care in the initial management of children who died from severe bacterial infection: a population-based confidential inquiry. *Pediatr Crit Care Med*. 2010;11(4):469-474.
 28. Mazzotta F, Troccoli T. Acute acro-ischemia in the child at the time of COVID-19. *Eur J Pediatr Dermatol*. 2020.
 29. Kolivras A, Dehavay F, Delplace D, et al. Coronavirus (COVID-19) infection-induced chilblains: a case report with histopathological findings. *JAAD Case Reports*. 2020;6:489-492.

How to cite this article: Elmas ÖF, Demirbaş A, Özyurt K, Atasoy M, Türsen Ü. Cutaneous manifestations of COVID-19: A review of the published literature. *Dermatologic Therapy*. 2020;33:e13696. <https://doi.org/10.1111/dth.13696>