

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/346943684>

The effect of gamma radiation and magnetic field on seed germination and seedling growth at low temperature in sorghum x sudangrass hybrids

Article in *Maydica* · December 2020

CITATIONS

7

READS

423

4 authors:



Ramazan Beyaz

Kırşehir Ahi Evran Üniversitesi

55 PUBLICATIONS 566 CITATIONS

SEE PROFILE



Yasin Özgen

Ankara University

17 PUBLICATIONS 178 CITATIONS

SEE PROFILE



Aslınur Çavdar

Ankara University

3 PUBLICATIONS 56 CITATIONS

SEE PROFILE



Mustafa Yıldız

Ankara University

125 PUBLICATIONS 1,232 CITATIONS

SEE PROFILE

The effect of gamma radiation and magnetic field on seed germination and seedling growth at low temperature in sorghum x sudangrass hybrids

Ramazan Beyaz^{1*}, Yasin Ozgen², Aslinur Cavdar³, Mustafa Yildiz²

¹ Kırşehir Ahi Evran University, Faculty of Agriculture, Department of Soil Science and Plant Nutrition, 40100-Bağbaşı, Kırşehir, Turkey

² Ankara University, Faculty of Agriculture, Department of Field Crops, Dışkapı, Ankara, Turkey

³ Ankara University, Biotechnology Institute, Dışkapı, Ankara, Turkey

* Corresponding author: E-mail: ramazanbeyaz@gmail.com

Keywords: *Sorghum x sudangrass hybrid, low-temperature, gamma radiation, magnetic field, germination, seedling growth*

Abstract

This study was aimed to provide seed germination and seedling growth of sorghum x sudangrass (*Sorghum bicolor* (L.) Moench) hybrids at low temperature by using gamma radiation and magnetic field strength. Seeds of two hybrids ("Aneto" and "Sugar Graze") were treated with different gamma doses (0-control, 100 Gy) and magnetic field strengths (0-control and 75 mT) for 72 hours while some were applied with both gamma radiation + magnetic field. Germination trials were carried out for 10 days in an incubator at 15°C temperature. Germinated seeds from each treatment were transferred to pots (5 pots with 4 seeds each) and cultured for 28 days in growth chamber at 15°C temperature. In this study, germination percentage, seedling growth percentage, seedling height, root length, seedling fresh and dry weights, and water content of seedling were determined. Lowest results in all parameters in both hybrids were observed in the control treatment without gamma radiation and magnetic field strength application. The highest values in all parameters studied were recorded in seeds treated both by gamma radiation+magnetic field strength applied together in "Aneto", and in seeds treated with magnetic field strength in "Sugar Graze". As a result of the research, gamma radiation and magnetic field strength applied to seed increased seed germination and seedling growth at a low temperature of 15°C.

Introduction

Sorghum x sudangrass (*Sorghum bicolor* (L.) Moench) hybrids produce quality forage that grows rapidly, tolerates heat and drought and high yielding. Sorghum x sudangrass (*Sorghum bicolor* (L.) Moench) hybrids are the plant most adapted to high temperature climate (Zerbini and Thomas, 2003; Ali et al, 2009; Borghi et al, 2013). Seed germination and plant growth take place between 25-35°C. Sorghum x sudangrass (*Sorghum bicolor* (L.) Moench) hybrids are generally sensitive to low-temperature stress (range of 20°C to about 0°C) (Yu et al, 2004). Sorghum suffers chilling injury when subjected to non-freezing temperatures below 10–15°C (Peacock, 1982). Maize, as an allied species of sorghum, is more tolerant to low temperature. However, several studies have been reported to improve maize tolerance against low-temperature (or cold stress). Thus, cold tolerance is required to be studied in sorghum.

Gamma irradiation along with other high energy rays is a type of electromagnetic waves which owe high penetration capacity into molecules and can bring about

ionization of the subject material by removing their electrons (UNSC, 2000). Exposure of plant materials specifically seeds to gamma irradiation results in mutagenic changes in living cells by several means. Since gamma radiation owes high energy and greater penetrability into exposed cells and tissues, DNA of the subject material may undergo severe alterations.. These alterations can either result from direct physical strikes of gamma irradiation on DNA or due to production of reactive O₂ species such as hydrogen peroxides, hydroxyl ions and other active atomic oxygen which can further interact with DNA and other cellular components and biomolecules resulting in the ionization, functional changes of proteins and enzymes and overall metabolic activities (Mittler, 2002; Wi et al, 2005). Gamma-rays can be employed to develop new varieties that can be readily adapted in a short period in different locations with varying agro-climatic and growing conditions, and where available resources are limited. Low doses of gamma rays have been reported to increase cell proliferation, germination, cell growth, enzyme activity, stress resistance, and crop yields (Baek et al, 2005; Kim et al, 2005).

All organisms live under the influence of the Earth's magnetic field (MF), also termed the geo MF (GMF), (5×10^{-6} Tesla (T)) with geographical variations in its intensity ranges from 25 to 65 μ T, its inclination, and declination (Maus et al, 2010). Biophysical methods like magnetic and electromagnetic stimulation can be promising and environmentally friendly methods in the future of agriculture. Since the stimulation of plants with magnetic field to increase the quantity and quality of seed germination, seedling development and yields of different species by influencing physiological and biochemical mechanisms has caught the interest of many scientists all over the world.

Many studies have reported the positive impact of gamma irradiation and magnetic field strength on seed germination and seedling growth in rice (Florez et al, 2004), maize (Florez et al, 2007), *Festuca arundinaceae* Schreb. and *Lolium perenne* L. (Carbonell et al, 2008), *Asparagus officinalis* L. (Soltani et al., 2006) for growing under adverse environmental conditions (Javed et al, 2011; Baghel et al, 2018; Radhakrishnan 2019; Majeed et al, 2018; Beyaz, 2020). There are limited reports on the effects of low-temperature on seed germination and seedling growth in sorghum hybrids (Ercoli et al, 2004; Yu et al, 2004; Maulana and Tesso, 2013; Tari et al, 2013; Antony et al, 2019). Therefore, this study was aimed to investigate the effects of gamma irradiation and magnetic field strength applied alone or together on seed germination and seedling growth in sorghum hybrids at 15°C temperature.

Material and methods

Plant Materials

Seeds of two sorghum x sudangrass (*Sorghum bicolor* (L.) Moench) hybrids ("Aneto" and "Sugar Graze") were used as plant material in this study. The seeds were obtained from the Field Crops Central Research Institute (Ankara, Turkey) and Ulusoy Seed Company (Ankara, Turkey).

Gamma irradiation

Seeds were irradiated with different doses (0-control and 1000 Gy) of ^{60}Co γ rays at 0.8 kGy h^{-1} at the Turkish Atomic Energy Authority, Sarayköy Nuclear Research and Training Center, Sarayköy, Ankara. Fricke and alanine dosimeters were used for dose mapping and determination of dose rates of gamma source. Seeds were irradiated along with a dosimeter for 75 Gy to be sure that ionization was uniform. The seeds were placed on a plate and rotated 360 degrees in a cylindrical radiation field for the gamma irradiation treatment

Magnetic field generation

A MF system consists of two Helmholtz coils forming an electromagnet mounted on a wooden frame. The number of turns of copper wire per coil was 3000. The mean MF in the center of the coils ranged from 50 to 500 mT. Coils which were placed horizontally, were connected to a power supply (0-12 A, ref. 13506-93, PHYWE, Göttingen, Germany). Moreover, intensity through the coils was measured by an ammeter. GD anode was placed between coils. The accuracy and uniformity of MF strengths were detected by a digital teslameter (ref. 13610-93, PHYWE, Göttingen, Germany).

Magnetic field treatment

Seeds of two hybrids ("Aneto" and "Sugar graze") placed in the middle of the gap between the coils were exposed to 75 mT MF strength for 72 h. The MF strength used in the study (75 mT) were determined according to the protocols described by Aycan et.al. (2018) and Bahadir et al (2018). All trials were carried out in laboratory condition where temperature and humidity were controlled

Growth conditions and morphological observations

Seeds of two hybrids ("Aneto" and "Sugar Graze") were treated with different gamma doses (0-control, 100 Gy) and magnetic field strengths (0-control and 75 mT) for 72 hours. One hundred seeds were treated with gamma and magnetic field separately while 100 seeds were applied with gamma radiation + magnetic field strength together. Five replicates of 25 seeds for each treatment for both hybrids were placed between filter papers that were watered with 20 ml of tap water, and put in plastic bags. Water content of the filter papers were controlled every day and they were watered with the same amount of water when it was needed. Plastic bags were incubated at 15°C in the dark for 10 days for seed germination evaluation. Germinated seeds from each treatment were transferred to pots (5 pots with 4 seeds each) and cultured for 28 days in growth chamber at 15°C temperature. Seed germination (%), seedling growth (%), seedling height (cm), root length (cm), fresh and dry weight seedling contents (g), and seedling water content in weight and percentage were recorded on the 28th day after germinated seeds were transferred to soil. Dry weights were measured after drying samples at 105°C for 2.5 h.

Statistical Analysis

The experiment was conducted according to the "Completely Randomized Block Design" concept. Five replicates were tested, and each replicate contain-

Table 1 - The effects of gamma radiation and magnetic field strength on seed germination and seedling growth at a low temperature (15°C) in sorghum x sudangrass hybrids

Hybrids	Treatments	Germination (%)	*Seedling Growth (%)	Seedling Height (cm)	Seedling Root Length (cm)	Seedling Fresh Weight (g)	Seedling Dry Weight (g)	Seedling Water Content (g) – Percent (%)
"Aneto"	Control	43.33±1.7c	38.33±1.7b	13.27±1.6b	5.50±0.3b	0.11±0.009c	0.013±0.003b	0.10±0.007c – 91.00
	Gamma	46.67±1.7bc	38.33±1.7b	16.50±1.8a	7.83±0.8a	0.17±0.006b	0.017±0.003b	0.15±0.007ab – 88.00
	Magnetic Field	53.33±1.6b	43.33±1.6b	16.50±0.3a	9.00±1.2a	0.17±0.003b	0.023±0.003ab	0.15±0.006ab – 88.00
	Gamma + Magnetic Field	75.00±2.9a	68.33±3.3a	17.67±1.2a	9.33±1.2a	0.21±0.009a	0.033±0.003a	0.18±0.009a – 86.00
"Sugar Graze"	Control	48.33±1.7d	41.67±1.7d	13.33±1.7c	4.50±0.3c	0.08±0.01c	0.01±0.00c	0.07±0.009c – 87.50
	Gamma	65.00±2.9c	63.33±1.7c	14.83±0.7c	5.83±0.4c	0.12±0.003b	0.02±0.003c	0.10±0.006b – 83.33
	Magnetic Field	86.67±3.3a	81.67±1.7a	21.67±0.4a	11.33±0.7a	0.18±0.007a	0.04±0.003a	0.14±0.009a – 77.78
	Gamma + Magnetic Field	76.67±1.7b	71.67±3.3b	18.33±0.6b	7.50±0.3b	0.13±0.006b	0.03±0.003b	0.10±0.007b – 77.00

Each value is the mean of 3 replications. The study was set in two parallels. Values represent mean ± standard error of the mean. Values within a column followed by different letters are significantly different at the 0.01 level for each cultivar. *Seedling growth percentage means developed seedlings out of total germinated seed.

ned 25 seeds for seed germination percentage, and 4 seedlings for seedling growth percentage, seedling height, root length, fresh and dry weights of seedling, and seedling water content evaluations. The experiments were repeated twice. In all statistical analyses, 'SPSS Statistics ver. 22 (IBM Corp, Chicago, IL, USA)' was used. Analysis of Variance (One-way ANOVA) was applied for each experiment, followed by Duncan's multiple range test at the 1% level to find significant differences among treatments (Snedecor and Cochran, 1967).

Results and discussion

The effects of gamma radiation and magnetic field strength on seed germination percentage, seedling growth percentage, seedling height, root length, fresh and dry weights of seedling, and water content of seedling of two sorghum x sudangrass hybrids ("Aneto" and "Sugar Graze") are reported in Table 1. The differences in all parameters among treatments (control, gamma irradiation, magnetic field strength and gamma+magnetic field strength) in both hybrids were statistically significant ($p < 0.01$) (Table 1).

In hybrid "Aneto", the lowest data for all parameters were recorded in control treatment where there were no gamma irradiation and magnetic field strength. On the other hand, the highest data were observed in the treatment of gamma radiation + magnetic field strength applied to seeds together. The highest values recorded were 75.00%, 68.33%, 17.67 cm, 9.33 cm, 0.21 g, 0.033 g and 0.18 g in germination percentage, seedling growth percentage, seedling height, root length, fresh and dry weights of seedling, and water

content of seedling, respectively (Table 1). The effect of magnetic field strength on seed germination percentage, seedling growth percentage, root length and dry weight of seedling resulted higher than that observed by gamma radiation. For other parameters (seedling height, fresh weight of seedling and water content of seedling), values were similar in both gamma radiation and magnetic field strength treatments as 16.50 cm, 0.17 g and 0.15 g, respectively. Water content in percentage in the treatment of gamma radiation + magnetic field strength applied to seeds together was found as 86.00% which is the lower than other treatments (Table 1).

In hybrid "Sugar Graze", the lowest results in all parameters were again obtained from control treatment. The highest values were obtained from the treatment of magnetic field strength as 86.67%, 81.67%, 21.67 cm, 11.33 cm, 0.18 g, 0.04 g and 0.14 g in germination percentage, seedling growth percentage, seedling height, root length, fresh and dry weight of seedling, and water content of seedling, respectively. Water content in percentage in the treatment of magnetic field strength was recorded as 77.78.00% which is the lower than control and gamma treatments but a little bit higher than gamma radiation + magnetic field strength treatment.

The lower results in water content in percentage showed that the highest metabolic activity and mitotic cell division were in seedlings developed from gamma + magnetic field treatment.

In our study, the highest results were obtained from treated seeds (by gamma radiation and magnetic field alone or together) in all parameters in both hybrids. Seed germination and seedling growth are the critical



Fig. 1 - Seedlings of hybrid "Aneto" developed from seeds treated by gamma radiation + magnetic field (on the left) and from non-treated seeds (control) (on the right)

and/or main indicators for the resistance of plants to adverse environmental conditions. Beyaz et al (2016) and Bahadır et al (2018) confirmed the positive effect of gamma irradiation and magnetic stimulation on seed germination and seedling growth in *Lathyrus chrysanthus* Boiss. Payez et al (2013) reported that electromagnetic field increase seed germination and growth of wheat seedlings. Shine et al (2011) reported the beneficial effect of pre-sowing magnetic treatment for improving germination parameters and biomass accumulation in soybean. It has also been reported that decreased germination and seedling growth in stressed rice seedlings is due to decreased mobilization of starch and α -amylase activity (Lin and Kao, 1995).

From our findings, it can be concluded that the lower levels of all parameters of non-treated (control) seeds were directly due to a decreased water uptake from the soil and consequently, a reduced mobilization of nutrients in the plant. Treatment of seeds with gamma radiation and magnetic field either alone or together increased permeability of the seed coat and so enabled water, all solutes and nutrients to transfer into the tissue more easily, providing all cells with high metabolic activity. In our case, increases in fresh and dry weights of seedlings developed from treated seeds at the end of the study were chiefly due to an increase in the absorption of water and other components from the soil. In our study, seedlings from non-treated seeds (control) were found smaller than the ones from treated ones in both hybrids. Dale (1988) states that fresh weight increase is mainly due to cell enlargement by water absorption, cell vacuolation, and turgor-derived wall expansion. The increase in dry weight is closely related to cell division and new material synthesis (Sunderland, 1960). Dry weight increase in seedlings developed from treated seeds is due to an increase in carbohydrate metabolism resulting from increased water uptake.

It has been stated that water stress alters the level of plant hormones (Morgan, 1990). The decreased cytokinin and gibberellic acid and increased abscisic acid content under stress conditions (Boucaud and Unger, 1976) suggest that growth reduction under stress conditions could be the result of the stress-induced changes in membrane permeability and water uptake due to altered endogenous hormonal levels (Ilan, 1971). Hsia (1973) has reported that the inhibition of growth under water stress conditions is the result of inhibition of cell division. Baghel et al (2017) reported that static magnetic field pretreatment (SMF) mitigated the adverse effects of water stress in soybean. Beyaz (2020) noted that low-dose of gamma radiation eliminating negative effects both salt and drought stress in common vetch (*Vicia sativa* L.). Wang et al (2018) also revealed that gamma ray pretreatment could improve the salt tolerance of barley seedlings. Selim and El-Nady (2011) stated that magnetic treatments act as a protective factors against water deficit in tomato.

Rochalska and Orzeszko-Rywka (2005) reported that a low frequency MF (16 Hz) can be used as a method of post-harvest seed improvement for different plant species, especially for seeds of temperature sensitive species germinating at low temperatures.

In the present study, magnetic field treatment alone was seen more effective than gamma radiation on seed germination and seedling growth in sorghum x sudangrass (*Sorghum bicolor* (L.) Moench) hybrids at a low temperature of 15°C. Lower results in gamma radiation treatment could be attributed to the dose applied. A dose-dependent enhancement of germination capacity of seeds and seedling growth by magnetic field treatment has also been reported in wheat seeds (Shine et al 2011). Singh et al (2012) showed that seed germination and seedling growth increased with gamma radiation in a dose-dependant manner in wheat plant. Bahadır et al (2018) reported that magnetic field (MF) strength and different exposure time periods have different effect on seed germination and seedling growth. Moreover, Marcu et al (2013) and Majeed et al (2017) reported that germination and growth response of different plants to gamma radiation treatment generally depends on the quantity of radiation dose, exposure time and plant species.

In the current study, the results showed that magnetic field + gamma radiation applied together also gave positive effects on seed germination and seedling growth in both hybrids. Our results were verified with the study of Alikamanoğlu et al 2007) who reported that magnetic field changes the effect of gamma radiation in a positive way as a result of combined treatment of magnetic field and gamma radiation in *Paulownia tomentosa*

tissue culture. Furthermore, Pietruszewski et al (2017) reported that the application of electromagnetic field (EMF) and electromagnetic radiation (EMR) stimulation could be a good tool for improving the yield and plant growth for agricultural production.

References

To our knowledge, this is the first study reporting that seed germination and seedling growth at low temperature in sorghum x sudangrass hybrids could be increased by gamma radiation and magnetic field treatments using either alone or both treatments. The effects of gamma radiation and magnetic field on other adverse environmental conditions such as salinity and drought in sorghum hybrids should be addressed in future studies.

Conflict of interest

Authors declare no conflict of interest

References

- Abdul M, Muhammad Z, Ullah R, Ali H, 2009. Genetic diversity and assessment of drought tolerant sorghum landraces based on morpho-physiological traits at different growth stages. *Plant Omics J* 2(5): 214-227.
- Ali MA, Niaz S, Abbas A, Sabir W, Jabran K, 2009. Genetic diversity and assessment of drought tolerant sorghum landraces based on morpho-physiological traits at different growth stages. *Plant Omics J* 2(5): 214-227.
- Alikamanoğlu S, Yaycılı O, Atak Ç, Rzakoulieva A, 2007. Effect of magnetic field and gamma radiation on paulownia tomentosa tissue culture. *Biotechnol Biotechnol Equip* 21(1): 49-53.
- Antony RM, Kirkham MB, Tod TC, Bean SR, Wilson JD, Armstrong PR, Elizabeth M, Brabec DL, 2019. Low-temperature tolerance of maize and sorghum seedlings grown under the same environmental conditions. *J Crop Improv* 33(3): 287-305.
- Aycan M, Beyaz R, Bahadir A, Yildiz M, 2018. The effect of magnetic field strength on shoot regeneration and Agrobacterium tumefaciens-mediated gene transfer in flax (*Linum usitatissimum* L.). *Czech J Genet Plant* 55(1): 20-27.
- Baek MH, Kim JH, Chung BY, Kim JS, Lee IS, 2005. Alleviation of salt stress by low dose g-irradiation in rice. *Biologia Plantarum* 49 (2): 273-276.
- Baghel L, Kataria S, Guruprasad KN, 2018.

Effect of static magnetic field pretreatment on growth, photosynthetic performance and yield of soybean under water stress. *Photosynthetica* 56: 718-730.

- Bahadir A, Beyaz R, Yildiz M, 2018. The effect of magnetic field on in vitro seedling growth and shoot regeneration from cotyledon node explants of *Lathyrus chrysanthus* Boiss. *Bioelectromagnetics* 39(7): 547-555.
- Beyaz R, Kahramanogullari CT, Yildiz C, Darcin ES, Yildiz M, 2016. The effect of gamma radiation on seed germination and seedling growth of *Lathyrus chrysanthus* Boiss. under in vitro conditions. *J Environ Radioact* 162: 129-133.
- Beyaz,R., 2020. Impact of gamma irradiation pretreatment on the growth of common vetch (*Vicia sativa* L.) seedlings grown under salt and drought stress. *Int J Radiat Biol* 96(2): 257-266.
- Borghi E, Crusciol CA, Nascente C, Sousa AS, Martins VV, Mateus PO, Costa C, 2013. Sorghum grain yield, forage biomass production and revenue as affected by intercropping time. *Eur J Agron* 51: 130-139.
- Boucaud J, Unger IA, 1976. Hormonal control of germination under saline conditions of three halophyte taxa in genus Suaeda. *Physiol Plant* 36: 197-200.
- Carbonell MV, Martinez E, Florez M, Maqueda R, Lopez-Pintor A, Amaya JM, 2008. Magnetic field treatments improve germination and seedling growth in *Festuca arundinacea* Schreb. and *Lolium perenne* L. *Seed Sci Technol* 36: 31-37.
- Dale JE, 1988. The control of leaf expansion. *Annu. Rev. Plant Physiol* 39: 267-295.
- Ercoli L, Mariotti M, Masoni A, Arduini I. 2004. Growth Responses of Sorghum Plants to Chilling Temperature and Duration of Exposure. *Eur J Agron* 21: 93-103.
- Florez M, Carbonell MV, Martínez E, 2004. Early sprouting and first stages of growth of rice seeds exposed to a magnetic field. *Electromagnetics* 23: 157-166.
- Florez M, Carbonell MV, Martínez E, 2007, Exposure of maize seeds to stationary magnetic fields: effects on germination and early growth. *Environ Exp Bot* 59: 68-75.
- Hsia TC, 1973. Plant responses to water stress. *Annu Rev Plant Physiol* 24: 219-270.
- Ilan I, 1971. Evidence for hormonal regulation of the selectivity of ion uptake by plant cells. *Physiol Plant* 25: 230-233.
- Javed N, Ashraf M, Akram N, Al-Qurainy F, 2011. Alleviation of adverse effects of

- drought stress on growth and some potential physiological attributes in maize (*Zea mays*) by seed electromagnetic treatment. *Photochem Photobiol* 87: 1354-1362.
- Kim JH, Chung BY, Kim JS, Wi SG, 2005. Effects of in Planta gamma-irradiation on growth, photosynthesis, and antioxidative capacity of red pepper (*Capsicum annuum* L.) plants. *J Plant Biol* 48: 47-56.
- Lin CC, Kao CH, 1995. NaCl stress in rice seedlings: Starch mobilization and the influence of GA3 on seedling growth. *Bot Bull Acad Sin* 36: 169-173.
- Majeed A, Muhammad Z, Ullah R, Ali H, 2018. Gamma irradiation : effect on germination and general growth characteristics of plants—a review. *Pak J Bot* 50(6): 2449-2453.
- Marcu D, Damian G, Cosma C, Cristea V, 2013. Gamma radiation effects on seed germination, growth and pigment content, and ESR study of induced free radicals in maize (*Zea mays*). *J Biol Phys* 39(4): 625-634.
- Marks N, Szecówka PS, 2010. Impact of variable magnetic field stimulation on growth of aboveground parts of potato plants. *Int Agrophysics* 24: 165-170.
- Maulana F, Tesso T T, 2013. Cold temperature episode at seedling and flowering stages reduces growth and yield components in Sorghum. *Crop Science* 53: 564-574.
- Maus S, Macmillan S, McLean S, Hamilton B, Thomson A, Nair M, Rollins C, 2010. The US/UK world magnetic model for 2010–015. http://www.ngdc.noaa.gov/geomag/WMM/data/WMM2010/WMM2010_Report.pdf. Accessed 5 May 2015.
- Mittler R, 2002. Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7: 405-410.
- Morgan PW, 1990. Effects of abiotic stresses on plant hormone systems. In: Koziowski TT (ed) *Stress Responses in Plants: Adaptation and Acclimation Mechanism*, Wiley-Liss, New York.
- Payez A, Ghanati F, Behmanesh M, Abdolmaleki P, Hajnorouzi A, Rajabbeigi E, 2013. Increase of seed germination, growth and membrane integrity of wheat seedlings by exposure to static and a 10-KHz electromagnetic field. *Electromagn Biol Med* 32(4): 417-429.
- Peacock JM, 1982. Response and tolerance of sorghum to temperature stress. In: House, L.R., et al (Eds.), *Sorghum in the Eighties*. Proceedings of the International Symposium on Sorghum, Patancheru, India, November 2–7, 1981. International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Hyderabad, India, pp. 143-159.
- Pietruszewski S, Muszyński S, Dziwulska A, 2007. Electromagnetic fields and electromagnetic radiation as non-invasive external stimulants for seeds (selected methods and responses). *Int. Agrophysics* 21: 95-100.
- Radhakrishnan R, 2019. Magnetic field regulates plant functions, growth and enhances tolerance against environmental stresses. *Physiol Mol Biol Plants* 25(5): 1107-1119.
- Selim AFH, El-Nady MF, 2011. Physio-anatomical responses of drought stressed tomato plants to magnetic field. *Acta Astronautica* 69: 387-396.
- Shine MB, Guruprasad KN, Anand A, 2011. Enhancement of germination, growth, and photosynthesis in soybean by pre-treatment of seeds with magnetic field. *Bioelectromagnetics* 32: 474-484.
- Singh B, Ahuja S, Singhal RK, Babu PV, 2012. Effect of gamma radiation on wheat plant growth due to impact on gas exchange characteristics and mineral nutrient uptake and utilization. *J Radioanal Nucl Chem* 298(1): 249-257.
- Snedecor GW, Cochran WG, 1980. *Statistical Methods*. Seventh Edition. Ames Iowa: The Iowa State University Press; 507 p.
- Soltani F, Kashi A, Arghavani M, 2006. Effect of magnetic field on *Asparagus officinalis* L. seed germination and seedling growth. *Seed Sci Technol* 34: 349-353.
- Sunderland N, 1960. Cell division and expansion in the growth of the leaf. *J Exp Bot* 11: 68-80.
- Tari I, Laskay Z, Takács Z, Poór Pi, 2013. Response of Sorghum to abiotic stresses: A Review. *J Agron Crop Sci* 199: 264-274.
- Wang X, Ma R, Cao Q, Shan Z, Jiao Z, 2018. Enhanced tolerance to salt stress in highland barley seedlings (*Hordeum vulgare* ssp. *vulgare*) by gamma irradiation pretreatment. *Acta Physiol Plant* 40: 174.
- Wi SG, Chung BY, Kim JH, Baek MH, Yan DH, Lee JW, Kim JS, 2005. Ultrastructural changes of cell organelles in Arabidopsis stems after gamma irradiation. *J Plant Biol* 48 (2): 195-200.
- Yua J, Tuinstra MR, Claassen MM, Gordon WB, Witt MD, 2004. Analysis of cold tolerance in sorghum under controlled environment conditions. *Field Crops Res* 85: 21-30.
- Zerbini IE, Thomas D, 2003. Opportunities for improvement of nutritive value in sorghum and pearl millet residues in South Asia through genetic enhancement. *Field Crops Res* 84: 3-15.